



WHITE PAPER

Clinical Nutrition for

Surgical and Injury Recovery

HEALFAST, INC.
SINCE ©2017

VISIT: WWW.HEALFASTPRODUCTS.COM
EMAIL: INFO@HEALFASTPRODUCTS.COM



Introduction to Clinical Nutrition for Surgery and Injury

Decades of scientific literature demonstrate that adequate nutrition and optimal outcomes are linked, not only to complications and major morbidity, but also to mortality. Undernourished pre-operative nutritional status has been linked consistently to an increase in post-operative complications including surgical site infections, longer hospital stays, slower wound healing, higher rate of mortality, amongst other complications.

Over 15 different meta analysis of trials suggest that perioperative administration of nutritional formulae contribute to a decreased rate of these same postoperative complications and consequently to a decreased length of stay in the hospital. Nutritional supplementation, however, is often underutilized in the U.S. healthcare system. There are a variety of potential reasons for this trend including questionable safety of some available ingredients, lack of nutritional training amongst providers, lack of resources and time, financial complexity in instituting proper nutrition care, concerns over lack of regulation in the over-the-counter market, and potential interactions with peri-operative and peri-trauma processes and prescribed medications with some supplement ingredients.

However, research has repeatedly demonstrated that utilizing nutritional management in the healthcare setting has high capacity for potential benefits and we believe that nutritional supplementation can address a significant unmet need by providing nutritional products that contain elements targeted to address specific clinical processes. Nutritional therapy also has the potential to be a cost-effective intervention to improve patient outcomes and avoid costly complications at low risk to the patient

if formulated properly. Supplements formulated for clinical use, however, must utilize elements, dosages, and routes of administration that are validated by evidence-based research while avoiding compounds with documented adverse effects or interactions in the peri-operative and peri-injury periods.

We note that given the variability in quality in the current supplement market, nutritional supplements manufactured for clinical uses need to utilize high grade, high fidelity ingredients, and be responsibly sourced and properly verified for purity and content. These supplements also need to be compounded by a qualified and reputable manufacturer in an FDA registered facility that utilizes Good Manufacturing Practice (GMP) standards and undergo extensive testing preferably by a third-party lab independent of the manufacturer.

We believe that the advantage of clinical nutrition to the patient cannot be ignored. Current challenges in the industry can be addressed with meticulous evidence-based formulation and education. This White Paper introduces to the reader the topic of clinical nutrition, it's challenges, and the scientific evidence. We aim to describe the clinical need to optimize nutrition in the peri-operative and peri-traumatic periods and to describe the current challenges and concerns in the field of oral nutritional supplementation.

We will then review the ingredients with the strongest scientific publication support that may be beneficial for a healing patient while providing a minimal risk profile.



In Summary:

- There is a clear link between optimized nutrition and clinical outcomes
- Large amount of scientific evidence suggests that perioperative administration of nutritional formulae contribute to a decreased rate of complications
- Nutritional supplementation is under-utilized in the healthcare
- There is a lack of high quality, medical grade nutritional supplements on the market
- There is large variability and safety in over-the-counter supplements
- Supplements made for clinical use need to be based on rigorous scientific evidence, utilizing elements, forms, and dose of ingredients with scientific backing
- Supplements for clinical use should undergo extensive facility testing, and be produced in accordance with FDA and GMP standards

The Need to Optimize Nutrition in a Clinical Setting

Nutritional status can have a profound effect on recovery in surgical and injury patients. This idea was first reported in 1930s, when a study showed that undernourished patients undergoing ulcer surgery had a risk of dying 10 times higher when compared with well-nourished individuals.¹

In 1990s, a follow up prospective study of 500 patients that included 200 surgical patients, admitted to university hospital, found that **40% of patients were undernourished at the time of admission.**² In another, separate eleven-year retrospective study, malnutrition rates were estimated to be up to 62% among the inpatient population.³ These findings conclude that malnutrition or nutritional insufficiency is very common in clinical setting across the healthcare settings.

Undernourished pre-operative nutritional status has since been linked consistently to an increase in post-operative complications, such as surgical site infection, longer hospital stays, slow wound healing, higher rate of mortality, and other surgical outcomes.^{4,5} On the other hand, there have also been studies that have shown that oral nutritional supplementation can reduce these very same complications, including length of stay (LOS), patient episode cost, and readmission rates.³

Surgical intervention itself is a complex set of processes which can result in derangements in physiological and metabolic responses, and can include decreased immunity, increased inflammation, high levels of stress hormones, increased breakdown of tissue and decreased creation of new tissue.

These changes contribute to increased protein and energy requirements after surgical intervention. When combined with prolonged preoperative and postoperative fasting and bowel preps, these factors may even increase the risk of poor post-operative nutrition status and lead to worse outcomes in relatively nourished individuals.⁶

Repleting patients with key nutrients prior to surgery and supporting their nutrition following surgery, may help patients prepare for the stress of surgical intervention and aid in a quicker post-operative recovery. **In the eleven-year retrospective study** mentioned above, oral nutritional supplementation prior to the procedure reduced LOS by 2.3 days, decreased episode cost by \$4,734, and reduced probability of readmissions within 30 days by up to 6.9%.³





Nutrition Support Is Often Overlooked by Healthcare Providers

Decades of literature demonstrate that adequate nutrition and optimal outcomes are clearly linked, not only to complications and major morbidity, but also to mortality. In a recent Swiss-Austrian survey of 176 surgical departments, the faculty was asked to give opinions regarding potential benefits and use of nutrition in surgical practice.

- Nearly 80% of the senior leadership agreed that attention to nutrition would decrease surgical complications
- And 59% felt that attention to nutrition could shorten hospital stay.
- Despite this overwhelming affirmation that appropriate nutrition therapy can decrease complications and hospital stay, only 20% of the surgeons surveyed implemented routine screening.
- When asked why they did not implement routine screening, 33% responded that logistical issues were too complex
- And 49% felt that financial issues prevented adequate screening and intervention.¹⁹⁹

In fact, surgeons and anesthesiologists routinely advise their patients to stop nutritional supplements at least two weeks prior to scheduled surgery.⁷ This is due, in part, to the popularity of complementary and alternative medical treatments such as herbal remedies and their potential to have serious and potentially harmful side effects when combined with medications commonly prescribed in the perioperative and peri-trauma periods. However, there are many micro and macronutrients that are of critical importance in aiding the healing process that do not have these significant interactions and complications similar to those of herbal alternatives. Interestingly, only a few products using these ingredients exist in the market today. This is especially alarming and warrants closer evaluation given the quantity of scientific research highlighting the importance and benefits of peri-operative nutritional support.^{8,9}

15 different meta analysis of trials suggest that peri-operative administration of nutritional formulae have contributed to a decreased rate of postoperative complications and consequently to a decreased length of stay in the hospital.¹⁷⁶⁻¹⁹⁸

In the peri-operative time frame, physicians rarely have time or adequate training to advise their patients on when or how to apply nutrition to support recovery. Nutritional supplementation, however, is a cost-effective therapy, making it inexpensive to provide with the potential to improve patient outcomes and avoid costly complications, at low risk to the patient.¹

Accounting for the high incidence of malnutrition and undernutrition, specifically among surgical and injury patients, improving patient outcomes with supplementation would come at a low cost to healthcare providers.¹ In effect, this could help decrease health care spending while improving quality of care provided.¹

Given upcoming regulations, such as Medicare Access and CHIP Reauthorization Act of 2015 (MACRA) and Merit-based Incentive Payment System (MIPS), higher emphasis will be placed on providing quality care and reducing complications. Although there is typically little financial incentive, training, or personnel needed for lifestyle modifications and nutrition management, in light of these new policies and incentives healthcare practitioners may be further encouraged to seek out opportunities to reduce complications and potentially in-hospital length of stay. One high potential intervention involves nutritional management.





Key Nutrients to Support Injury and Surgical Healing

Surgery and injury elicits a series of reaction including release of stress hormones and inflammatory mediators, such as cytokines. Cytokine response to infection and injury has major impact on metabolism and includes catabolism of glycogen, fat, and protein with release of glucose, fatty acids, and amino acids into the circulation. The release of these substrates diverts them from their normal purpose of maintaining peripheral protein (such as muscle) mass, to the tasks of healing and immune response.¹⁶²⁻¹⁶³

The consequence of catabolism is loss of muscle mass, which is crucial to both a short and long term functional recovery. Nutritional therapy may thus provide substrate and the energy for optimal healing and recovery and may slow down catabolic processes that lead to reduction in patient functional status.

Largely, surgical and injury related tissue healing is an anticipated drain on a patient's nutritional stores. Preloading with appropriate vitamins, minerals, amino acids, antioxidants, anti-inflammatories, probiotics, and trace minerals can better optimize a patient for the physiologic and metabolic stress that co-occurs with these processes.

We, however, must note in our review that certain elements are best purposed pre-operatively, and some are most needed post-operatively; thus it is valuable to have differentiate the unique pre and post-operative regimens that also take into consideration commonly used peri-operative processes, medications, events and anesthetics.

Reviewed below are the most studied nutrients that have a role in the peri-operative and peri-traumatic period. They are listed without particular order and serve as brief introductions since including all evidence on these ingredients would extend the review beyond the scope of this White Paper.

Additionally, there are several ingredients that may be potentially beneficial in the perioperative period, namely nucleotides and omega acids, however, current studies have had mixed results, and some even suggested these to be harmful in certain circumstances. More research on these ingredients are required to further determine their roles in clinical nutrition and thus they are not discussed below..

Although not discussed explicitly below, it should be noted that more than a single form of a specific compound may exist and that all forms may not necessarily be equal in terms of physiological effect. For example, a compounds such as copper or zinc have several specific formulations with different bioavailabilities and absorption profiles resulting in different physiological actions. Some ingredients, such as the probiotics, have specific subgroups, in this case, strains, of which some are helpful to the healing and recovery process and some are not. These details are relatively well studied, but are also beyond the scope of this review.



Vitamin C

Vitamin C supplementation perioperatively or during peri-injury period is highly recommended since Vitamin C promotes healing by various mechanisms including partaking an essential role in collagen formation, providing support for the immune system, and acting as a tissue antioxidant and free-radical scavenger.

According to the U.S. Centers for Disease Control and Prevention's most recent National Report on Biochemical Indicators of Diet and Nutrition in the U.S. Population, approximately 6% of the US populations is already deficient in Vitamin C.¹² Surgical intervention further compounds the problem by lowering blood Vitamin C concentration. This reduction in Vitamin C levels is postulated to occur due to increased oxidative stress and as well as redistribution. Decreased Vitamin C levels are significant particularly due to the fact that Vitamin C acts as an antioxidant thereby protecting cells from damage caused by free radicals which abound in times of tissue stress. Further, Vitamin C is needed for normal immune system function, iron absorption and production of collagen matrix, which are all critical processes in wound healing. Due to its many biochemical functions, Vitamin C is considered not only an essential micronutrient for maintaining health, but also an important therapeutic supplement in a variety of clinical conditions, including surgical and injury processes.^{10, 11}

In a small study of patients who underwent uncomplicated gastrectomy for gastric cancer, blood concentration of Vitamin C decreased post-operatively and the reduction remained significant for seven days following surgery.¹³ It was further noted that if there were associated post-operative complications Vitamin C levels are reduced even further. Although normal baseline physiological functioning requires Vitamin C for optimal metabolism, it was shown that doses much higher than the recommended daily allowance may be needed to normalize plasma and tissue Vitamin C concentration in post-operative patients.

As the body cannot synthesize Vitamin C, the main source of Vitamin C is from consumption of fruits, vegetables and plant foods.¹⁴ Intake of these foods to the level need for recovery is often impossible in the peri-operative or peri-injury period. We note that relatively high doses of supplemental Vitamin C, combined with other trace elements, have been shown to dramatically accelerate wound healing, in some as much as **50%**.

Most multivitamin preparations available in the U.S. contain approximately 200 mg of Vitamin C. However, in uncomplicated surgery patients, more than 500 milligrams per day of Vitamin C supplementation may be required to support wound healing. Further, Vitamin C can have additional impact on other physiological functions, for example, in cardiac surgery patients, Vitamin C supplementation may prevent post-operative atrial fibrillation.^{15,16} Recently, it was shown that oral Vitamin C supplementation, in association with beta-blockers, is even more effective in preventing post-operative atrial fibrillation than beta-blockers alone.^{17,18}

Roles of Vitamin C are truly multifaceted. In 2016, a systematic review and meta analysis in showed that there is even evidence that preoperative dose of Vitamin C can be used as an adjunct for reducing postoperative morphine consumption, and that there is high-level evidence supporting perioperative Vitamin C supplementation for complex regional pain syndrome (CRPS I) prevention after extremity surgery.¹⁶⁴

Overall, the importance of Vitamin C to patients are clear in that well controlled clinical trials in sepsis, trauma and major burns testing high-dose Vitamin C have shown clinical benefit with nearly non-existent risk profile.¹⁶⁵

Citrus Bioflavonoids

Citrus Bioflavonoids are a collection of compounds typically found in citrus fruits which have a great variety of benefits, including functions as strong antioxidants, with potential properties such as anti-tumor agents and cardiovascular support. Importantly, they have a strong ability to improve Vitamin C absorption, which in itself a very strong benefit. These compounds additionally provide a broad spectrum of documented benefits in cases of heart disease, cancer, vision disorders, allergies, and viral infections. Certain flavonoids, notably diosmin and hesperidin, have been used routinely in Europe for many years to treat varicose veins, hemorrhoids, edema, bruising, and wounds.

Two separate European studies indicate that oral application of flavonoids to patients with leg ulcers (a common complication of chronic venous insufficiency) greatly increases healing rates. When this formulation was given for two months to a group of individuals who were suffering from leg ulcers, their ulcers healed faster and to a greater degree than those of a control group that received a placebo instead.¹⁹ A larger, randomized study recently completed in Poland confirmed these results and highlighted the supportive role that flavonoids play in the wound-repair process.²⁰

Another recent study demonstrated that infected wounds heal more quickly if treated with flavonoids.²¹ This study, performed on guinea pigs, showed that either topical or oral application of diosmin/hesperidin was effective in accelerating the healing of infected wounds which resulted in the suggestion that may also have antibacterial or immune supportive functions.

Diosmin and hesperidin may have many other beneficial properties in wound healing.²² On a molecular level, flavonoids have been shown to inhibit the synthesis of prostaglandins, a feature that likely contributes to their anti-inflammatory effect.²³ The combination of all these properties at the site of a wound is important in creating an environment that favors tissue repair.

Vitamin B Complex

Vitamin B Complex is a series of vitamins that work together in many metabolic reactions. There is ample and decisive evidence that deficiency in any vitamins in the B complex can significantly decrease capacity for tissue regeneration and healing and induce many deleterious metabolic effects. Vitamin B complex is thought to have many benefits and is particularly known for its integral role in energy productions and has just recently been recognized for its benefits in the healing process. It is integral to all major wound healing processes.⁴¹

Vitamin B complex is thought to speed up wound healing, increase protein synthesis and the number of repair cells at the wound, prevent excessive inflammation, protect against oxidative stress, and increase tissue blood flow.^{37, 38, 39, 40}





Vitamin B Complex: Multiple Mechanisms of Action

Vitamin	Name	Molecular Function
Vitamin B ₁	Thiamine	Thiamine plays a central role in the release of energy from carbohydrates. It is involved in RNA and DNA production, as well as nerve function. ¹⁶⁶
Vitamin B ₂	Riboflavin	Riboflavin is involved in release of energy in the in several metabolic reactios as well as the catabolism of fatty acids. ¹⁶⁷
Vitamin B ₃	Niacin	Niacin plays animportant role in energy transfer reactions in the metabolism of glucose, fat and alcohol. ¹⁶⁸ It is also a oenzyme in lipid and nucleic acid synthesis. ¹⁶⁹
Vitamin B ₅	Pantothenate	Pantothenic acid is involved in the oxidation of fatty acids and carbohydrates. Coenzyme A, which can be synthesised from pantothenic acid, is involved in the synthesis of amino acids, fatty acids, ketone bodies, cholesterol, ¹⁷⁰ phospholipids, steroid hormones, neurotransmitters, and antibodies. ¹⁷¹
Vitamin B ₆	Pyridoxine	Pyridoxine serves as a cofactor in many enzyme reactions mainly in amino acid metabolism including biosynthesis of neurotransmitters.
Vitamin B ₈	Biotin	Biotin plays a key role in the metabolism of lipids, proteins and carbohydrates. It is a critical co-enzyme involved in the synthesis of fatty acids and in gluconeogenesis; It is also involved in synthesis of several amino acids
Vitamin B ₉	Folate	Folate acts as a co-enzyme in the form of tetrahydrofolate (THF), which is needed for normal cell division, especially in times of tissue growth, which are times of rapid growth. Folate also aids in the production of red blood cells. ¹⁷³
Vitamin B ₁₂	Cobolamine	Pantothenic acid is involved in the oxidation of fatty acids and carbohydrates. Coenzyme A, which can be synthesised from pantothenic acid, is involved in the synthesis of amino acids, fatty acids, ketone bodies, cholesterol, ¹⁷⁰ phospholipids, steroid hormones, neurotransmitters, and antibodies. ¹⁷¹



Vitamin A

Vitamin A is required in cell growth (especially tissue and bone formation), which is very important in the healing process that takes place after surgery and injuries. It is known to activate production of connective tissue, including collagen (promotes tissue synthesis), and helps new blood vessels grow, which is necessary for nourishing newly formed tissue. It increases collagen cross-linking and wound breaking strength.^{44, 45, 46, 47}

Vitamin A increases the initial inflammatory response in wounds, thus facilitating the migration of cells that assist with wound healing at the site. Importantly, it reverses corticosteroid-induced inhibition of cutaneous wound healing.^{42, 43}

Vitamin A also enhances resistance to infection by stimulating the body's immune function.

Bromelain

Another well-known and popular healing adjunct is Bromelain, an extract from pineapple.⁵⁸⁻⁶⁰ Various studies indicate that Bromelain reduces swelling, bruising, pain after surgery and physical injuries, as well as healing time. Bromelain is often used in conjunction with Quercetin.⁶¹⁻⁸⁷

A wide range of therapeutic benefits have been claimed for bromelain; such as reversible inhibition of platelet aggregation, angina pectoris, bronchitis, sinusitis, surgical traumas, thrombophlebitis, pyelonephritis and enhanced absorption of drugs, particularly of antibiotics.⁸⁸⁻¹⁰¹

Recent results from preclinical and pharmacological studies recommend bromelain as an orally given drug for complementary tumor therapy. Especially promising are reports on animal experiments claiming an antimetastatic efficacy as inhibition of growth and invasiveness of tumor cells.¹⁰¹⁻¹⁰⁹

Bromelain also has potential to eliminate burn debris and to accelerate wound healing and in several studies Bromelain supplementations significantly increased the resorption rate of hematomas.¹¹⁰⁻¹¹⁹

Quercetin

Quercetin, a plant pigment naturally found in foods such as onions, is a so called bioflavonoid. It promotes decreased histamine release in the body and thus acts like an anti-inflammatory as well as an anti-oxidant. Usually after surgery inflammation is a common response, and often unfavorable response if excessive or prolonged.

Quercetin supplementation is also known to enhance wound healing, as Quercetin incorporates into the collagen matrices.

Quercetin, particularly in conjunction with Bromelain may help reduce bruising, swelling and scarring.^{25, 26}

Probiotics

Probiotics are widely known for their specific healing benefits regarding certain surgeries, as well as other broader beneficial effects in general.

Surgery patients often receive antibiotic treatment. The use of perioperative broad-spectrum antibiotics has become the standard of care for most major surgeries. Antibiotics are used to reduce postoperative infection risk. This practice has been validated in the literature and is now considered an essential part of the surgical process. While considered essential to surgical success and safety, antibiotic use does contribute to postoperative morbidity. Significant gastrointestinal flora disruption can occur, leading to decreased resistance to pathogens known to cause antibiotic associated diarrhea, such as *Clostridium Difficile*. Studies have shown that the use of antibiotics known to disturb the gastrointestinal flora is associated with clinical symptoms, such as diarrhea, **in as many as 30% of patients.**^{27, 28}

Research has shown that the use of probiotics can help to restore gut microecology through various mechanisms including: receptor competition, competition for nutrients, inhibition of epithelial and mucosal adherence of pathogens, introduction of lower colonic pH favoring the growth of nonpathogenic species, stimulation of immunity, and/or production of antimicrobial substances.^{29,30}

In fact it is common practice in Britain for hospitals to routinely use probiotic supplements to reduce the incidence of antibiotic-resistance and hospital-acquired infections. Studies show that probiotic supplementation leads to a significant reduction of bacterial infection rates. Several probiotics strains also have well documented efficacy data in treatment of antibiotic associated diarrhea.

Other studies show that the incidence of postoperative infections (including Pneumonia and Cholangitis), the duration of postoperative hospital stay, and the length of antibiotic therapy were all lower among patients receiving probiotics.

Probiotics, when used following surgery, can accelerate recovery through improved immunity, as well as reduce the risk of postoperative infection and antibiotic-associated diarrhea. The existing body of evidence supports the use of Lactobacillus species, either alone or in combination with other genera.³¹





Amino Acids: Arginine and Glutamine

Amino acids play a crucial role in protein synthesis and muscle tissue repair. Specifically, L-arginine is an amino acid that has been shown to stimulate wound healing, while L-glutamine restores cellular energy stores. In clinical studies, patients given L-arginine supplementation after major surgery benefited from a faster recovery of immunological parameters and fewer infectious complications.³²

Randomized clinical trials of patients undergoing surgery for hip fracture demonstrated that providing pre- and post-operative energy-protein supplements resulted in better recovery plasma proteins and fewer post-operative complications, even among patients with normal pre-operative nutritional status.^{33, 34}

In addition, a recent study showed that amino acid supplementation assisted with surgical recovery in patients who had undergone total knee arthroplasty, as demonstrated by attenuated muscle atrophy and accelerated return to functional mobility.³⁵

Glutamine

Glutamine is the most abundant amino acid in the body and accounts for 60% of the intracellular amino acid pool. This amino acid is considered to be conditionally essential as a deficiency can occur rapidly after injury. Glutamine is used as an energy source after the stress response. In addition, glutamine is the primary fuel source for rapidly dividing cells like epithelial cell during healing.

Glutamine has potent antioxidant activity and also has direct immunological function by stimulating white cell proliferation through its use as energy. Glutamine has anticatabolic and anabolic properties also and is the rate-limiting agent for new protein synthesis.⁴⁸⁻⁵¹

Because of its many roles in the wound, it is of particular concern when there is a rapid fall in both intracellular and extracellular glutamine levels, to a deficiency state, in the presence of a major wound.⁵²

Glutamine protects against excessive inflammatory injury by inducing the expression of heat shock proteins, which provide cellular protection in states of inflammation, injury, and stress. Furthermore, glutamine can modulate and preserve gut function, which is compromised in severe stress.⁵³

Glutamine, besides being an important metabolic fuel, can also be considered an important nutritional mediator of the inflammatory response. It does appear to decrease wound infection and it does improve healing in experimental studies. Further, prospective, controlled, and randomized clinical trials show decreased mortality and infectious morbidity in burn patients given glutamine supplements.⁵⁴⁻⁵⁷

Arginine

As a single agent, Arginine is the best-studied component of immunonutrition, with copious evidence showing that Arginine is beneficial to wound healing.¹²⁰⁻¹²²

Arginine, while normally a non-essential amino acid, becomes an essential nutrient during periods of physiological stress. However, during these periods Arginine is rapidly depleted, and this can affect the immune system response in injured states and during other disease processes.¹²³⁻¹²⁵

Arginine is critical for collagen synthesis, improving our immune function, and wound healing abilities.¹²⁵⁻¹²⁸

Numerous studies show that Arginine supplementation, administered either orally or parenterally, accelerates wound healing by increasing collagen deposition in wounds.¹²⁸⁻¹³¹

Not only does Arginine increase collagen accumulation, but it also causes increased activity of peripheral blood lymphocytes. It improves epithelial reconstitution after intestinal injuries and also has been shown to nearly double improvements in pressure ulcer healing times.¹²⁵⁻¹³⁴

Metabolically, Arginine is a precursor to Proline, and is thus recruited specifically for collagen synthesis. It is also a precursor for ornithine, which is critical for polyamine synthesis, and for nitric oxide. In fact, Arginine is the sole metabolic substrate for the synthesis of Nitric Oxide, which plays a role in wound collagen synthesis and regulation of blood flow.¹³⁵

For example, inhibition of NO synthesis during healing results in weaker wound recovery and decreased collagen synthesis. This is commonly found in cases of diabetes and malnutrition.¹²⁵⁻¹²⁹





Trace Minerals

Zinc

Zinc is a cofactor of many enzymes, including some that are needed to repair wounds. It is also an antioxidant and has significant antibacterial properties. Even a mild deficiency of zinc can interfere with optimal recovery from everyday tissue damage and more serious trauma.

The hypermetabolic state associated with surgery and injuries leads to a marked increase in loss of zinc, and a risk for a zinc deficiency state has adverse effects on the healing process including a decrease in epithelial rate, wound strength, and decreased collagen strength.

One controlled trial found the healing time of a surgical wound was reduced by 43% with oral supplementation zinc. Zinc also plays a vital role in the immune system and can reduce inflammation, bacterial growth, and scar tissue creation, whereas a deficiency can certainly worsen recovery.²⁴

Selenium

Selenium has strong antioxidant roles. There is some suggestion that it may expedite wound healing in burn patients. A prospective, randomized controlled trial investigated the effect of large intravenous doses of trace elements (copper, selenium, and zinc) on patients with major burns and found that cutaneous concentrations of those trace elements were increased, antioxidant status (as measured by normalization of plasma glutathione peroxidase level) was improved, and wound healing was benefited as measured by decreased graft requirement.¹³⁶ The antioxidant properties of selenium benefit wound healing.^{137, 138, 139} Further, selenium is commonly deficient in patients in the hospital setting.¹⁴⁰

Copper

Copper has a complex role in various cells, modulating several cytokines and growth factor mechanisms of action, and it is essentially involved in all stages of the wound healing process.¹⁴¹⁻¹⁵⁶ Old and recent literature shows that the activity of PDGF, VEGF, and Angiogenin is copper-dependent.

Copper peptides, specifically, are small protein fragments that, along with copper, stimulate collagen production and can improve the effectiveness of antioxidants and can escalate the amount of protein synthesis in collagen and elastin thus also supporting wound healing processes.

Magnesium

Magnesium has vast metabolic impact in regards to recovery.¹⁵⁷ It aids in muscle relaxation, soreness, and cramping.^{158, 159} Although magnesium is not a primary analgesic in itself, it enhances the analgesic actions of more established analgesics as an adjuvant agent.

The role of magnesium for perioperative analgesia has been investigated by many authors and it was found in a randomized control trial that administration of magnesium pre-operatively can reduce pain scores and decrease amount of opioids needed to control pain.¹⁶⁰ This finding has been repeated and generalized in multiple studies in both IV and oral formulations. Magnesium can also improve endothelial function, especially in patients with atherosclerosis.¹⁶¹





Addressing the Need for Clinical Nutrition

There is a significant amount of clinical data to support that poor nutritional status is linked to increases in post-operative complication and adverse outcomes in surgical patients. Literature supports that nutritional therapy can reduce these same risks and improve outcomes. Extensive clinical research has validated the supportive role of nutritional elements in peri-operative or peri-injury time frames.

Despite the relative abundance of scientific research, there is a striking lack of high-quality, pharmaceutical-grade nutritional supplements and medicinals that are specifically designed to support the nutritional status of these patients.

Nutritional supplements can address this significant unmet need by providing nutritional products that contain elements targeted to address specific clinical and medical conditions, while using elements, dosages, and routes of administration that have been validated by evidence-based research and avoiding compounds that have documented adverse effects or interactions in the peri-operative and peri-injury periods.

With variability in quality in the current supplement industry, great care must be taken in manufacturing a supplement for clinical use. The supplement for this purpose must utilize high grade, high fidelity ingredients, responsibly sourced and properly verified for purity and content. These supplements also need to be compounded by a qualified manufacturer in an FDA registered facility that utilizes Good Manufacturing Practice (GMP) standards and need to undergo extensive testing for heavy metals, pesticides, and microbial contaminants, preferably by a third-party lab independent of the manufacturer.

Furthermore, optimal interventions for nutrition therapy should be comprehensive, spanning the entire perioperative or peri-injury period, and target various parts of the process with specific nutrients that benefit the underlying set of processes. Recent advances in perioperative care have emphasized the concept of “prehabilitation” to provide pre-operative intervention such as physical therapy and nutritional support to best prepare the patient prior to surgery. In fact, research has shown that adding nutritional formulas during the week of surgery enhances recovery and return to baseline function.³⁶

The breadth of research in the field of clinical nutrition needs to be applied. Current evidence suggests that many nutrients can be compounded at relatively low cost, provide a favorable safety profile, all with significant potential to improve patient care, optimize recovery, and reduce complications.



Clinical References

1. Studley, HO. Percentage of weight loss: a basic indicator of surgical risk in patients with chronic peptic ulcer. *JAMA* 1936;106:458.
2. McWhirter JP, Pennington CR. Incidence and recognition of malnutrition in hospital. *BMJ* 1994;308(6934):945.
3. Philipson T, Snider J, Lakdawalla D, Stryckman B, Goldman D. Impact of Oral Nutritional Supplementation On Hospital Outcomes. *Clinical Nutrition*. 2013;32. doi:10.1016/s0261-5614(13)60017-5.
4. Burden S, Todd C, Hill J, Lal S. Preoperative nutrition support in patients undergoing gastrointestinal surgery. *Cochrane Database Syst Rev* 2012 Nov 14;11:CD008879. doi: 10.1002/14651858.CD008879.pub2.
5. Meguid MM, Laviano A. Malnutrition, outcome and nutritional support: time to revisit the issues. *Ann Thorac Surg* 2001;71:766-8.
6. Souba WW, Wilmore D. Diet and nutrition in the care of the patient with surgery, trauma, and sepsis. In: Shils M, Olson J, Shike M, Ross AC, editors. *Modern nutrition in health and disease*. 9th ed. Baltimore, MD: Williams & Wilkins; 1999. p. 1589-618.
7. Rispler DT, Sara J. The impact of complementary and alternative treatment modalities on the care of orthopaedic patients. *Journal of the American Academy of Orthopaedic Surgeons* 2011;19(10):634-643.
8. Akbarshahi H, et al. Perioperative nutrition in elective, gastrointestinal surgery – potential for improvement? *Dig Surg* 2008;25:165-174.
9. Lugli AK, Wykes L, Carli F. Strategies for perioperative nutrition support in obese, diabetic and geriatric patients. *Clinical Nutrition* 1 2008;27:16-24.
10. Philipson T, Snider J, Lakdawalla D, Stryckman B, Goldman D. Impact of Oral Nutritional Supplementation On Hospital Outcomes. *Clinical Nutrition*. 2013;32. doi:10.1016/s0261-5614(13)60017-5.
11. McWhirter JP, Pennington CR. Incidence and recognition of malnutrition in hospital. *BMJ* 1994;308(6934):945.
12. National Institutes of Health. Vitamin C – Fact Sheet for Consumers. Available at <https://ods.od.nih.gov/factsheets/VitaminC-Consumer/>.
13. U.S. Centers for Disease Control and Prevention. Second National Report on Biochemical Indicators of Diet and Nutrition in the U.S. Population 2012, Executive Summary. Available at http://www.cdc.gov/nutritionreport/pdf/exesummary_web_032612.pdf.
14. Fukushima R, Yamazaki E. Vitamin C requirement in surgical patients. *Current Opinion in Clinical Nutrition and Metabolic Care* 2010;13(6):669-676.
15. National Institutes of Health. Vitamin C – Fact Sheet for Consumers. Available at <https://ods.od.nih.gov/factsheets/VitaminC-Consumer/>.
16. Akbarshahi H, et al. Perioperative nutrition in elective, gastrointestinal surgery – potential for improvement? *Dig Surg* 2008;25:165-174.
17. Akbarshahi H, et al. Perioperative nutrition in elective, gastrointestinal surgery – potential for improvement? *Dig Surg* 2008;25:165-174.
18. Eslami M, et al. Oral ascorbic acid in combination with beta-blockers is more effective than beta-blockers alone in the prevention of atrial fibrillation after coronary artery bypass grafting. *Tex Heart Inst J* 2007;34:268-274.
19. Guilhou JJ, Fevrier F, Debure C, et al. Benefit of a 2-month treatment with a micronized, purified flavonoidic fraction on venous ulcer healing. A randomized, double-blind, controlled versus placebo trial. *Int J Microcirc Clin Exp* 1997;17 Suppl 1: 21-6.
20. Gliński W, Chodyncka B, Roszkiewicz J, et al. Effectiveness of a micronized purified flavonoid fraction (MPFF) in the healing process of lower limb ulcers. *Minerva Cardioangiolo* 2001;49:107-14.
21. Hasanoglu A, Cengiz A, Suleyman O, Kali K, Senol M, Ertas E. Efficacy of micronized flavonoid fraction in healing of clean and infected wounds. *Int J Angiol* 2001;10:41-4.
22. Struckmann JR, Nicolaidis AN. Flavonoids. A review of the pharmacology and therapeutic efficacy of Dalon 500 mg in patients with chronic venous insufficiency and related disorders. *Angiology* 1994;45:419-28.
23. Manthey JA. Biological properties of flavonoids pertaining to inflammation. *Microcirc* 2000;7:S29-34.
24. Desneves KJ, et al. Treatment with supplementary arginine, vitamin C and zinc in patients with pressure ulcers: a randomized controlled trial. *Clinical Nutrition* 2005;24:979-987.
25. MedlinePlus. Bromelain. Available at <http://www.nlm.nih.gov/medlineplus/druginfo/natural/895.html>.
26. WebMD. Quercetin. Available www.webmd.com/vitamins-supplements/ingredientmono-294-quercetin.aspx?activeingredientid=294&activeingredientname=quercetin.
27. Barbut F, Meynard JL. Managing antibiotic associated diarrhea. *BMJ*. 2002;324(7350):1345-1346
28. Rolfe RD. The role of probiotic cultures in the control of gastrointestinal health. *J Nutr*. 2000;130:(2S suppl) 396S-402S
29. Cremonini F, Di Caro S, Santarelli L, et al. Probiotics in antibiotic-associated diarrhoea. *Dig Liver Dis*. 2002;34:(suppl 2) S78-S80
30. Susanne Hempel, Sydne J. Newberry, et al. Probiotics for the prevention and treatment of antibiotic-associated diarrhea, a systematic review and meta-analysis. *JAMA*. 2012; 307(18):1959-1969.
31. Gibson GR & Roberfroid MB (1995) Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *J Nutr*; 125:1401 – 1412.
32. Tepaske R. Immunonutrition. *Curr Opin Anaesthesiol* 1997;10:86-91.
- 33– Botella-Carretero JI, et al. Perioperative oral nutritional supplements in normal or mildly undernourished geriatric patients submitted to surgery for hip fracture: a randomized clinical trial. *Clinical Nutrition* 2010;29(5):574-579.
34. Botella-Carretero JI, et al. Effects of oral nutritional supplements in normally or mildly undernourished geriatric patients after surgery for hip fracture: a randomized clinical trial. *J Parenter Enteral Nutr* 2008;32:120-128.
35. Dreyer HC, et al. Essential amino acid supplementation in patients following total knee arthroplasty. *J Clin Invest* 2013;123(11):4654-4666. 5– McFarland LV. Epidemiology, risk factors and treatments for antibiotic-associated diarrhea. *Dig Dis*. 1998;16(5):292-307
36. Martindale RG, et al. Perioperative nutrition: what is the current landscape? *J Parenter Enteral Nutr* 2013;37(5 Suppl):5S-20S.
37. Weimann BI, Hermann D. Calcium D-Pantothenate. *Int J Vitam Nutr Res*. 1999 Mar;69(2):113-9.
38. Neiva RF, Al-Shammari K, Nociti FH Jr, et al. Effects of vitamin-B complex supplementation on periodontal wound healing *J Periodontol*, 76 (2005), pp. 1084-1091
39. Mochizuki S, Takano M, Sugano N, et al. The effect of B vitamin supplementation on wound healing in type 2 diabetic mice. *Journal of Clinical Biochemistry and Nutrition*. 2016;58(1):64-68. doi:10.3164/jcbs.14-122.
40. Benderitter M, Hadj-Saad F, Lhuissier M, Maupoil V, Guillard JC, Rochette L. Effects of exhaustive exercise and vitamin B6 deficiency on free radical oxidative process in male trained rats. *Free Radic Biol Med*. 1996;21:541–549.
41. Lakshmi R, Lakshmi AV, Bamji MS. Skin wound healing in riboflavin deficiency. *Biochem Med Metab Biol*. 1989 Dec;42(3):185-91.
42. Ehrlich HP, Hunt TK: Effects of cortisone and vitamin A on wound healing. *Ann Surg*. 1968;167:324- 328



43. Ehrlich HP, Tarver H, Hunt TK: Effects of vitamin A and glucocorticoids upon inflammation and collagen synthesis. *Ann Surg.* 1973;177:222- 227
44. Greenwald DP, Sharzer LA, Padawer J, Levenson SM, Seifter E: Zone II flexor tendon repair: effects of vitamin A, E, beta-carotene. *J Surg Res.* 1990;49:98- 102 Article
45. Seifter E, Crowley LV, Rettura G et al. Influence of vitamin A on wound healing in rats with femoral fracture. *Ann Surg.* 1975;181:836- 841 Article
46. Niu XT, Cushman B, Reisner A, Levenson SM, Demetriou AA Effect of dietary supplementation with vitamin A on arterial healing in rats. *J Surg Res.* 1987;42:61- 65 Article
47. Phillips JD, Kim CS, Fonkalsrud EW, Zeng H, Dindar H: Effects of chronic corticosteroids and vitamin A on the healing of intestinal anastomoses. *Am J Surg.* 1992;163:71- 77
48. Wischemeyer PE., Kahana M., Wolfson R., Ren H., Musch MM., and Chang EB.: Glutamine induces heat shock protein and protects against endotoxin shock in the rat. *J Appl Physiol* 2001; 90:2403.
49. Wischemeyer PE.: Glutamine and heat shock protein expression. *Nutrition* 2002; 18:225.
50. Aquino VM., Harvey AR., Garvin JH., Godder KT., Nieder ML., Adams RH., Jackson GB., and Sandler ES.: A double-blind randomized placebo-controlled study of oral glutamine in the prevention of mucositis in children undergoing hematopoietic stem cell transplantation: a pediatric blood and marrow transplant consortium study. *Bone Marrow Transplant* 2005; 36:611.
51. Ward E., Smith M., Henderson M., Reid U., Lewis I., Kinsey S., Allgar V., Bowers D., and Picton SV.: The effect of high-dose enteral glutamine on the incidence and severity of mucositis in paediatric oncology patients. *Eur J Clin Nutr* 2009; 63:134.
52. Beale RJ., Sherry T., Lei K., Campbell-Stephen L., McCook J., Smith J., Venetz W., Altheid B., Stehle P., and Schneider H.: Early enteral supplementation with key pharmac nutrients improves sequential organ failure assessment score in critically ill patients with sepsis: outcome of a randomized, controlled, double-blind trial. *Crit Care Med* 2008; 36:131
53. Sabine Ellinger. (2014) Micronutrients, Arginine, and Glutamine: Does Supplementation Provide an Efficient Tool for Prevention and Treatment of Different Kinds of Wounds?. *Advances in Wound Care* 3:11, 691-707
54. Peck MD, Chang Y. Nutritional support for burn injuries. *J Nutr Biochem* 1999;10:380–96.
55. Berger, M.M. Antioxidant micronutrients in major trauma and burns: evidence and practice. *Nutr Clin Pract.* 2006;21:438–449.
56. Gottschlich MM, Jenkins M, Warden GD et al. Differential effects of three dietary regimens on selected outcome variables in burn patients. *JPEN J Parenter Enteral Nutr.* 1990;14:225- 236
57. Saffle JR, Wiebke G, Jennings K, Morris E, Barton RG Randomized trial of immuno-enhancing enteral nutrition in burn patients. *J Trauma.* 1997;42:793- 802
58. Mondal S, Bhattacharya S, Pandey JN, Biswas M. Evaluation of acute anti-inflammatory effect of Ananas Comosus leaf extract in Rats. *Pharmacology online.* 2011;3:1312–1315.
59. Taussig SJ, Batkin S. Bromelain, the enzyme complex of pineapple (*Ananas comosus*) and its clinical application: an update. *Journal of Ethnopharmacology.* 1988;22(2):191–203.
60. Heinicke RM, Gortner WA. Stem bromelain: a new protease preparation from pineapple plants. *Economic Botany.* 1957;11(3):225–234.
61. Livio M, Gaetano GD, Donati MB. Effect of bromelain on fibrinogen level, prothrombin complex and platelet aggregation in the rat—a preliminary report. *Drugs under Experimental and Clinical Research.* 1978;1:49–53.
62. Neubauer RA. A plant protease for potentiation of and possible replacement of antibiotics. *Experimental Medicine and Surgery.* 1961;19:143–160.
63. Renzini G, Varego M. Die resorption von tetrazyklin ingenenwart von Bromelain bei oraler application. *Arzneimittel-Forschung Drug Research.* 1972;2:410–412.
64. Maurer HR. Bromelain: biochemistry, pharmacology and medical use. *Cellular and Molecular Life Sciences.* 2001;58(9):1234–1245.
65. Tochi BN, Wang Z, Xu SY, Zhang W. Therapeutic application of pineapple protease (Bromelain): a review. *Pakistan Journal of Nutrition.* 2008;7(4):513–520.
66. Taussig SJ. The mechanism of the physiological action of bromelain. *Medical Hypotheses.* 1980;6(1):99–104.
67. Hale LP. Proteolytic activity and immunogenicity of oral bromelain within the gastrointestinal tract of mice. *International Immunopharmacology.* 2004;4(2):255–264.
68. Ley CM, Tsiami A, Ni Q, Robinson N. A review of the use of bromelain in cardiovascular diseases. *Journal of Chinese Integrative Medicine.* 2011;9(7):702–710.
69. Chobotova K, Vernallis AB, Majid FAA. Bromelain activity and potential as an anti-cancer agent: current evidence and perspectives. *Cancer Letters.* 2010;290(2):148–156.
70. Castell JV, Friedrich G, Kuhn CS, Poppe GE. Intestinal absorption of undegraded proteins in men: presence of bromelain in plasma after oral intake. *American Journal of Physiology.* 1997;273(1):G139–G146.
71. Bhattacharyya BK. Bromelain: an overview. *Natural Product Radiance.* 2008;7(4):359–363. Rowan AD, Buttle DJ. Pineapple cysteine endopeptidases. *Methods in Enzymology.* 1994;244:555–568
72. Yoshioka K Izutsa S, Asa Y, Takeda Y. Inactivation kinetics of enzyme pharmaceuticals in aqueous solutions. *Pharmaceutical Research.* 1991;4:480–485.
74. Napper AD, Bennet SP, Borowski M, et al. Purification and characterization of multiple forms of the pineapple-stem-derived cysteine proteinases ananain and comosain. *Biochemical Journal.* 1994;301(3):727–735.
74. Cooreman W. Bromelain. In: Ruysen R, Lauwers A, editors. *Pharmaceutical Enzymes Properties and Assay Methods.* Gent, Belgium: E. Story-Scientia Scientific Publishing Co.; 1978. pp. 107–121. Filippova IY, Lysogorskaya EN, Oksenoit ES, Rudenskaya GN, Stepanov VM.
75. L-Pyroglutamyl-L-phenylalanyl-L-leucine-p-nitroanilide: a chromogenic substrate for thiol proteinase assay. *Analytical Biochemistry.* 1984;143(2):293–297.
76. Seifert J, Ganser R, Brendel W. Absorption of a proteolytic enzyme originating from plants out of the gastro-intestinal tract into blood and lymph of rats. *Zeitschrift fur Gastroenterologie.* 1979;17(1):1–8.
77. Shiew PS, Fang YL, Majid FAA. In vitro study of bromelain activity in artificial stomach juice and blood. *Proceedings of the 3rd International Conference on Biotechnology for the Wellness Industry;* 2010; PWTC;
78. Neumayer C, Fügl A, Nanobashvili J, et al. Combined enzymatic and antioxidative treatment reduces ischemia-reperfusion injury in rabbit skeletal muscle. *Journal of Surgical Research.* 2006;133(2):150–158.
79. Heinicke RM, van der Wal L, Yokoyama M. Effect of bromelain (Ananase) on human platelet aggregation. *Experientia.* 1972;28(10):844–845.
80. King DE, Ellis TM, Everett CJ, Mainous AG. Medication use for diabetes, hypertension, and hypercholesterolemia from 1988–1994 to 2001–2006. *Southern Medical Journal.* 2009;102(11):1127–1132.
81. Secor ER, Jr., William FC, Michelle MC, et al. Bromelain exerts anti-inflammatory effects in an ovalbumin-induced murin model of allergic disease. *Cellular Immunology.* 2005;237:68–75.[PMC free article]
82. Juhasz B, Thirunavukkarasu M, Pant R, et al. Bromelain induces cardioprotection against ischemia-reperfusion injury through Akt/FOXO pathway in rat myocardium. *American Journal of Physiology.* 2008;294(3):H1365–H1370.
83. Brien S, Lewith G, Walker A, Hicks SM, Middleton D. Bromelain as a treatment for osteoarthritis: a review of clinical studies. *Evidence-Based Complementary and Alternative Medicine.* 2004;1(3):251–257.
84. Bodi T. The effects of oral bromelains on tissue permeability to antibiotics and pain response to bradykinin: double blind studies on human subjects. *Clinical Medicine.* 1966;73:61–65.



85. Kumakura S, Yamashita M, Tsurufuji S. Effect of bromelain on kaolin-induced inflammation in rats. *European Journal of Pharmacology*. 1988;150(3):295–301.
86. Cohen A, Goldman J. Bromelain therapy in rheumatoid arthritis. *Pennsylvania Medical Journal*. 1964;67:27–30.
87. Barth H, Guseo A, Klein R. In vitro study on the immunological effect of bromelain and trypsin on mononuclear cells from humans. *European Journal of Medical Research*. 2005;10(8):325–331.
88. Lehmann PV. Immunomodulation by proteolytic enzymes. *Nephrology Dialysis Transplantation*. 1996;11(6):953–955.
89. Desser L, Rehberger A, Kokron E, Paukovits W. Cytokine synthesis in human peripheral blood mononuclear cells after oral administration of polyenzyme preparations. *Oncology*. 1993;50(6):403–407
90. Desser L, Rehberger A, Paukovits W. Proteolytic enzymes and amylase induce cytokine production in human peripheral blood mononuclear cells in vitro. *Cancer Biotherapy*. 1994;9(3):253–263.
91. Eckert K, Grabowska E, Stange R, Schneider U, Eschmann K, Maurer HR. Effects of oral bromelain administration on the impaired immunocytotoxicity of mononuclear cells from mammary tumor patients. *Oncology Reports*. 1999;6(6):1191–1199.
92. Engwerda CR, Andrew D, Murphy M, Mynott TL. Bromelain activates murine macrophages and natural killer cells in vitro. *Cellular Immunology*. 2001;210(1):5–10.
93. Engwerda CR, Andrew D, Ladhams A, Mynott TL. Bromelain modulates T cell and B cell immune responses in vitro and in vivo. *Cellular Immunology*. 2001;210(1):66–75.
94. Mynott TL, Ladhams A, Scarmato P, Engwerda CR. Bromelain, from pineapple stems, proteolytically blocks activation of extracellular regulated kinase-2 in T cells. *Journal of Immunology*. 1999;163(5):2568–2575.
95. Secor ER, Jr., Singh A, Guernsey LA, et al. Bromelain treatment reduces CD25 expression on activated CD4+ T cells in vitro. *International Immunopharmacology*. 2009;9(3):340–346.
96. Leipner J, Iten F, Saller R. Therapy with proteolytic enzymes in rheumatic disorders. *BioDrugs*. 2002;15(12):779–789.
97. Lotz-Winter H. On the pharmacology of bromelain: an update with special regard to animal studies on dose-dependent effects. *Planta Medica*. 1990;56(3):249–253.
98. Livio M, De Gaetano G, Donati MB. Effect of bromelain on fibrinogen level, prothrombin complex factors and platelet aggregation in rat: a preliminary report. *Drugs under Experimental and Clinical Research*. 1978;4:21–23.
99. De-Guili M, Pirota F. Bromelain: interaction with some protease inhibitors and rabbit specific antiserum. *Drugs under Experimental and Clinical Research*. 1978;4:21–23.
100. Taussig SJ, Batkin S. Bromelain, the enzyme complex of pineapple (*Ananas comosus*) and its clinical application: an update. *Journal of Ethnopharmacology*. 1988;22(2):191–203.
101. Mynott TL, Guandalini S, Raimondi F, Fasano A. Bromelain prevents secretion caused by *Vibrio cholerae* and *Escherichia coli* enterotoxins in rabbit ileum in vitro. *Gastroenterology*. 1997;113(1):175–184.
102. Chandler DS, Mynott TL. Bromelain protects piglets from diarrhoea caused by oral challenge with K88 positive enterotoxigenic *Escherichia coli*. *Gut*. 1998;43(2):196–202.
103. Mynott TL, Luke RKJ, Chandler DS. Oral administration of protease inhibits enterotoxigenic *Escherichia coli* receptor activity in piglet small intestine. *Gut*. 1996;38(1):28–32.
104. Bézé R, Lopes MTP, Salas CE, Hernández M. In vivo antitumoral activity of stem pineapple (*Ananas comosus*) bromelain. *Planta Medica*. 2007;73(13):1377–1383.
105. Taussig SJ, Szekerczes J, Batkin S. Inhibition of tumour growth in vitro by bromelain, an extract of the pineapple plant (*Ananas comosus*) *Planta Medica*. 1985;6:538–539.
106. Tysnes BB, Maurer HR, Porwol T, Probst B, Bjerkvig R, Hoover F. Bromelain reversibly inhibits invasive properties of glioma cells. *Neoplasia*. 2001;3(6):469–479.
107. Huang JR, Wu CC, Hou RCW, Jeng KC. Bromelain inhibits lipopolysaccharide-induced cytokine production in human THP-1 monocytes via the removal of CD14. *Immunological Investigations*. 2008;37(4):263–277.
108. Hou RCW, Chen YS, Huang JR, Jeng KCG. Cross-linked bromelain inhibits lipopolysaccharide-induced cytokine production involving cellular signaling suppression in rats. *Journal of Agricultural and Food Chemistry*. 2006;54(6):2193–2198.
108. Hou RCW, Chen YS, Huang JR, Jeng KCG. Cross-linked bromelain inhibits lipopolysaccharide-induced cytokine production involving cellular signaling suppression in rats. *Journal of Agricultural and Food Chemistry*. 2006;54(6):2193–2198.
109. Tassman GC, Zafran JN, Zayon GM. Evaluation of a plate proteolytic enzyme for the control of inflammation and pain. *Journal of Dental Medicine*. 1964;19:73–77.
110. Tassman GC, Zafran JN, Zayon GM. A double-blind crossover study of a plant proteolytic enzyme in oral surgery. *The Journal of Dental Medicine*. 1965;20:51–54.
111. Houck JC, Chang CM, Klein G. Isolation of an effective debriding agent from the stems of pineapple plants. *International Journal of Tissue Reactions*. 1983;5(2):125–134.
112. Rosenberg L, Kriehner Y, Silverstein E, et al. Selectivity of a Bromelain Based Enzymatic Debridement Agent: A Porcine Study. Elsevier; 2012.
113. Singer AJ, McClain SA, Taira BR, Rooney J, Steinhaff N, Rosenberg L. Rapid and selective enzymatic debridement of porcine comb burns with bromelain-derived Debrase: acute-phase preservation of noninjured tissue and zone of stasis. *Journal of Burn Care and Research*. 2010;31(2):304–309.
114. Wu SY, Hu W, Zhang B, Liu S, Wang JM, Wang AM. Bromelain ameliorates the wound microenvironment and improves the healing of firearm wounds. *Journal of Surgical Research*. 2012;176:503–509.
115. Hu W, Wang AM, Wu SY, et al. Debriding effect of bromelain on firearm wounds in pigs. *The Journal of Trauma*. 2011;71(4):966–972.
116. Miller JG, Carruthers HR, Burd DAR. An algorithmic approach to the management of cutaneous burns. *Burns*. 1992;18(3):200–211.
117. Sheridan RL, Tompkins RG, Burke JF. Management of burn wounds with prompt excision and immediate closure. *Journal of Intensive Care Medicine*. 1994;237:68–75.
118. Salisbury RE. In-thermal burns. In: McCarthy JC, editor. *Plastic Surgery*. Vol. 1. 1990. pp.787–830.
119. Moss IN, Frazier CV, Martin GJ. Bromelain -the pharmacology of the enzyme. *Archives of International Pharmacody*. 1963;145:166–189.
120. Chaturvedi R, Asim M, Lewis ND, Algood HM, Cover TL, Kim PY, Wilson KT. L-arginine availability regulates inducible nitric oxide synthase-dependent host defense against *Helicobacter pylori*. *Infect Immun* 75: 4305–4315, 2007
121. Lucotti P, Setola E, Monti LD, Galluccio E, Costa S, Sandoli EP, Fermo I, Rabaiotti G, Gatti R, Piatti P. Beneficial effects of a long-term oral L-arginine treatment added to a hypocaloric diet and exercise training program in obese, insulin-resistant type 2 diabetic patients. *Am J Physiol Endocrinol Metab* 291: E906–E912, 2006
122. J.C.L. Neyens, E. Cereda, E.P. Meijer, C. Lindholm, J.M.G.A. Schols. (2017) Arginine-enriched oral nutritional supplementation in the treatment of pressure ulcers: A literature review. *Wound Medicine* 16, 46-51
123. Emanuele Cereda, J. C. L. Neyens, R. Caccialanza, M. Rondanelli, J. M. G. A. Schols. (2016) Efficacy of a disease-specific nutritional support for pressure ulcer healing: A systematic review and meta-analysis. *The journal of Nutrition, Health & Aging*
124. J. Wesley Alexander, Dorothy M. Supp. (2014) Role of Arginine and Omega-3 Fatty Acids in Wound Healing and Infection. *Advances in Wound Care* 3:11, 682-690



125. Sabine Ellinger. (2014) Micronutrients, Arginine, and Glutamine: Does Supplementation Provide an Efficient Tool for Prevention and Treatment of Different Kinds of Wounds? *Advances in Wound Care* 3:11, 691-707
126. Desneves, K.J., Todorovic, B.E., Cassar, A., Crowe, T.C. Treatment with supplementary arginine, vitamin C and zinc in patients with pressure ulcers: a randomised controlled trial. *Clin Nutr.* 2005;24:979–987.
127. Popovic PJ, Zeh HJ III, Ochoa JB. Arginine and immunity. *J Nutr.* 2007;137(6 suppl 2):1681S-1686S. Google Scholar Medline
128. Daly JM, Lieberman MD, Goldfine J et al. Enteral nutrition with supplemental arginine, RNA, and ω 3 fatty acids in patients after operation: immunologic, metabolic and clinical outcome. *Surgery.* 1992;112:56- 67
129. Kemen M, Senkal M, Homann HH et al. Early postoperative enteral nutrition with arginine, ω -3 fatty acids and ribonucleic acid-supplemented diet versus placebo in cancer patients: an immunologic evaluation of Impact®. *Crit Care Med.* 1995;23:652- 659
130. Senkal M, Kemen M, Homann HH et al. Modulation of postoperative immune response by enteral nutrition with a diet enriched with arginine, RNA, and ω -3 fatty acids in patients with upper gastrointestinal cancer. *Eur J Surg.* 1995;161:115- 122
131. Cerra FB, Lehmann S, Konstantinides N et al. Improvement in immune function in ICU patients by enteral nutrition supplemented with arginine, RNA, and menhaden oil is independent of nitrogen balance. *Nutrition.* 1991;7:193- 199
132. Daly JM, Reynolds J, Thom A, . Immune and metabolic effects of arginine in the surgical patient. *Ann Surg.* 1988;208:512-523. Google Scholar Medline
133. Boger RH, Bode-Boger SM. The clinical pharmacology of L-arginine. *Ann Rev Pharmacol Toxicol.* 2001;41:79-99. Google Scholar Medline
134. Gianotti L, Braga M, Fortis C, . A prospective, randomized clinical trial on perioperative feeding with an arginine-, omega-3 fatty acid-, and RNA-enriched enteral diet: effect on host response and nutritional status. *JPEN J Parenter Enteral Nutr.* 1999;23:314-320. Google Scholar Abstract
135. Albina JE, Abate JA, Mastrofrancesco B. Role of ornithine as a proline precursor in healing wounds. *J Surg Res* 55: 97–102, 1993
136. Berger MM, Baines M, Raffoul W, Benathan M, Chioloro RL, Reeves C, Revely JP, Cayeux MC, Sénéchaud I, and Shenkin A: Trace element supplementation after major burns modulates antioxidant status and clinical course by way of increased tissue trace element concentrations. *Am J Clin Nutr* 2007; 85:1293.
137. Berger MM., Baines M., Raffoul W., Benathan M., Chioloro RL., Reeves C., Revely JP., Cayeux MC., Sénéchaud I., and Shenkin A.: Trace element supplementation after major burns modulates antioxidant status and clinical course by way of increased tissue trace element concentrations. *Am J Clin Nutr* 2007; 85:1293.
138. Blass SC., Goose H., Tolba RH., Stoffel-Wagner B., Kabir K., Burger C., Stehle P., and Ellinger S.: Time to wound closure in trauma patients with disorders in wound healing is shortened by supplements containing antioxidant micronutrients and glutamine: a PRCT. *Clin Nutr* 2012; 31:469.
139. Beale RJ., Sherry T., Lei K., Campbell-Stephen L., McCook J., Smith J., Venetz W., Alteheld B., Stehle P., and Schneider H.: Early enteral supplementation with key pharmacutrients improves sequential organ failure assessment score in critically ill patients with sepsis: outcome of a randomized, controlled, double-blind trial. *Crit Care Med* 2008; 36:131.
140. Blass, S.C., Ellinger, S., Goost, H., Engels, J., Wirtz, D.C., Burger, C. et al, Deficiencies in zinc and selenium status are common in elderly trauma patients. *Clin Nutr.* 2007;2:157–158.
141. Allison Paige Kornblatt, Vincenzo Giuseppe Nicoletti, Alessio Travaglia, The neglected role of copper ions in wound healing, *Journal of Inorganic Biochemistry*, Volume 161, 2016, Pages 1-8, ISSN 0162-0134
- 142 T. Ashino, V. Sudhakar, N. Urao, J. Oshikawa, G.F. Chen, H. Wang, Y. Huo, L. Finney, S. Vogt, R.D. McKinney, E.B. Maryon, J.H. Kaplan, M. Ushio-Fukai, T. Fukai, *Circ. Res.* 107 (2010) 787–799.
143. W. Feng, F. Ye, W. Xue, Z. Zhou, Y.J. Kang, *Mol. Pharmacol.* 75 (2009) 174–182.
144. G.F. Hu, *J. Cell. Biochem.* 69 (1998) 326–335.
145. B.E. Kim, T. Nevitt, D.J. Thiele, *Nat. Chem. Biol.* 4 (2008) 176–185.
146. D. La Mendola, D. Farkas, F. Bellia, A. Magri, A. Travaglia, O. Hansson, E. Rizzarelli, *Inorg. Chem.* 51 (2012) 128–141.
147. L. Mandinov, A. Mandinova, S. Kyurkchiev, D. Kyurkchiev, I. Kehayov, V. Kolev, R. Soldi, C. Bagala, E.D. de Muinck, V. Lindner, M.J. Post, M. Simons, S. Bellum, I. Prudovsky, T. Maciag, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 6700–6705.
148. F.X. Maquart, L. Pickart, M. Laurent, P. Gillery, J.C. Monboisse, J.P. Borel, *FEBS Lett.* 238 (1988) 343–346.
149. N. Philips, H. Hwang, S. Chauhan, D. Leonardi, S. Gonzalez, *Connect. Tissue Res.* 51 (2010) 224–229.
150. G.M. Ross, I.L. Shamovsky, S.B. Woo, J.I. Post, P.N. Vrkljan, G. Lawrance, M. Solc, S.M. Dostaler, K.E. Neet, R.J. Riopelle, *J. Neurochem.* 78 (2001) 515–523.
151. C.K. Sen, S. Khanna, M. Venojarvi, P. Trikha, E.C. Ellison, T.K. Hunt, S. Roy, *American journal of physiology, Heart Circ. Physiol.* 282 (2002) H1821–H1827.
152. A. Simeon, F. Monier, H. Emonard, P. Gillery, P. Birembaut, W. Hornebeck, F.X. Maquart, *J. Investig. Dermatol.* 112 (1999) 957–964.
153. F. Soncin, J.D. Guitton, T. Cartwright, J. Badet, *Biochem. Biophys. Res. Commun.* 236 (1997) 604–610.
154. I. Tenaud, I. Sainte-Marie, O. Jumbou, P. Litoux, B. Dreno, *Br. J. Dermatol.* 140 (1999) 26–34.
155. C.Y. Tsai, J.C. Finley, S.S. Ali, H.H. Patel, S.B. Howell, *Biochem. Pharmacol.* 84 (2012) 1007–1013.
156. H. Xie, Y.J. Kang, *Curr. Med. Chem.* 16 (2009) 1304–1314
157. Fawcett WJ, Haxby EJ, Male DA. Magnesium: physiology and pharmacology. *Br J Anaesth.* 1999;83:302–320.
158. The effects of magnesium supplementation on exercise performance. Newhouse IJ, Finstad EW. *Clin J Sport Med* [2000]
159. Clarkson PM , Haymes EM, *Medicine and Science in Sports and Exercise* [01 Jun 1995, 27(6):831-843]
160. Gildasio S. De Oliveira, Jane Bialek, Paul Fitzgerald, John Y. S. Kim, Robert J. McCarthy. Systemic magnesium to improve quality of post-surgical recovery in outpatient segmental mastectomy: a randomized, double-blind, placebo-controlled trial. *Magnesium Research.* 2013;26(4):156-164. doi:10.1684/mrh.2014.0349
161. Shechter M, Sharir M, Paul-Labrador M, Forrester J, Bairey Merz CN. Oral magnesium therapy improves endothelial function in patients with coronary artery disease. *Circulation* 2000;102:2353–2358.
162. U.O. Gustafsson, H. Opielstrup, A. Thorell, J. Nygren, O. Ljungqvist Adherence to the ERAS protocol is associated with 5-year survival after colorectal cancer surgery: a retrospective cohort study *World J Surg*, 40 (2016), pp. 1741-1747
163. C. Gillis, F. Carli Promoting perioperative metabolic and nutritional care *Anesthesiology*, 123 (2015), pp. 1455-1472
164. Chen, Suzan MD, Roffey, Darren M. PhD, Dion, Charles-Antoine BSc, Arab, Abdullah MD, Wai, Eugene K. MD, Effect of Perioperative Vitamin C Supplementation on Postoperative Pain and the Incidence of Chronic Regional Pain Syndrome: A Systematic Review and Meta-Analysis
165. Vitamin C supplementation in the critically ill patient Berger, Mette M.a; Oudemans-van Straaten, Heleen M. *Current Opinion in Clinical Nutrition & Metabolic Care: March 2015 - Volume 18 - Issue 2 - p 193–201*



166. Fattal-Valevski, A (2011). "Thiamin (vitamin B1)". *Journal of Evidence-Based Complementary & Alternative Medicine*. 16 (1): 12–20. doi:10.1177/1533210110392941.

167 "Riboflavin". *Alternative Medicine Review*. 13 (4): 334–340. 2008. PMID 19152481

168. Whitney, N; Rolfes, S; Crowe, T; Cameron-Smith D; Walsh, A (2011). *Understanding Nutrition*. Melbourne: Cengage Learning

169. National Academy of Sciences. Institute of Medicine. Food and Nutrition Board, ed. (1998). "Chapter 6 - Niacin". *Dietary Reference Intakes for Thiamine, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin and Choline*. Washington, D.C.: National Academy Press.

170. University of Bristol (2002). "Pantothenic Acid". Retrieved 16 September 2012.

171. Gropper, S; Smith, J (2009). *Advanced nutrition and human metabolism*. Belmont, CA: Cengage Learning.

172. University of Bristol (2012). "Biotin". Retrieved 17 September 2012.

173. National Academy of Sciences. Institute of Medicine. Food and Nutrition Board, ed. (1998). "Chapter 8 - Folate". *Dietary Reference Intakes for Thiamine, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin and Choline*. Washington, D.C.: National Academy Press.

174. University of Bristol (2002). "Vitamin B12". Retrieved 16 September 2012.

175. DSM (2012). "Vitamin B12". Retrieved 16 September 2012.

176. Daly JM, Lieberman MD, Goldfine J, Shou J, Weintraub F, Rosato EF, et al. Enteral nutrition with supplemental arginine, RNA, and omega-3 fatty acids in patients after operation: immunologic, metabolic, and clinical outcome. *Surgery* 1992;112:56e67.

177. Gianotti L, Braga M, Vignali A, Balzano G, Zerbi A, Bisagni P, et al. Effect of route of delivery and formulation of postoperative nutritional support in patients undergoing major operations for malignant neoplasms. *Arch Surg* 1997;132:1222e9.