

QUICK TIPS FOR SUCCESS

(and if you already know what you're doing, just enjoy the meme!)

The early life cycle of mushrooms are delicate and prone to contamination. This guide offers some quick tips to maximize the performance of your [All-In-One Bag\(s\)](#).


Sanitization and Preparation

The contents of your AIOB may have shifted during shipping. Gently massage rye grain layer to redistribute and even out the moisture content. For best results, allow bag(s) to rest for 24-48 hours before inoculating and use within 30 days. Perishable, use as soon as possible!

- * Wipe down your workspace, AIOB, syringes, and other tools with isopropyl alcohol.
- * Keep the air as still as possible by turning off fans, heaters, A/C, etc.
- * Wash your hands and face. Wear a clean set of clothes, and gloves and mask if you'd like.
- * You will have higher success rates if transfers are done in front of laminar flow hood; we manufacture a hobbist size flow hood (SKU: MSFFU-1). Alternatively, a still air box (SAB) is an economical and effective alternative.
- * Move slowly and deliberately to minimize the movement of contaminants, especially if working in a SAB.
- * If you're using a spore or culture syringe, shake gently for even distribution. Flame sterilize the needle until red hot. Let cool for a few seconds before injecting at least 1-3 mL solution per bag. The more you inject, the faster colonization times will be!

How to Use All-in-One Grow Bags

1. Inoculate your AIOB by injecting your syringe into the injection port. Allow the bags 2-4 weeks to colonize at a comfortable temperature - around 75F. If you're comfortable, your mushrooms are too!
2. Once the grains are roughly 50% colonized, gently massage the grains, spreading the mycelium throughout the bottom layer.
3. Allow another 2-4 weeks for the mycelium to colonize the entire bag.
4. When the bag is fully colonized, fruit it straight from the bag by cutting the top of the bag open. Loosely fold the bag over to allow for fresh air exchange (FAE), and mist if necessary. Alternatively, this would be the time to transfer the bag contents into a fruiting chamber or monotub.



Congrats! Now for the toughest part... having patience!

