

2 results

1 article found by citation matching

Consumption of water containing over 3.5 mg of dissolved hydrogen could improve vascular endothelial function.

PMID: 25378931 [Free PMC article](#). Clinical Trial.

1 **Consumption of water containing over 3.5 mg of dissolved hydrogen could improve vascular endothelial function.**

Sakai T, Sato B, Hara K, Hara Y, Naritomi Y, Koyanagi S, Hara H, Nagao T, Ishibashi T.

Vasc Health Risk Manag. 2014 Oct 17;10:591-7. doi: 10.2147/VHRM.S68844. eCollection 2014.

PMID: 25378931 [Free PMC article](#). Clinical Trial.

2 **Peripheral endothelial function can be improved by daily consumption of water containing over 7 ppm of dissolved hydrogen: A randomized controlled trial.**

Ishibashi T, Kawamoto K, Matsuno K, Ishihara G, Baba T, Komori N.

PLoS One. 2020 May 29;15(5):e0233484. doi: 10.1371/journal.pone.0233484. eCollection 2020.

PMID: 32470022 [Free PMC article](#). Clinical Trial.

« First

< Prev

Page 1 of 1 Next >

Last »

FOLLOW NCBI



Connect with NLM

National Library of Medicine
8600 Rockville Pike
Bethesda, MD 20894

- Web Policies
- FOIA
- HHS Vulnerability Disclosure
- Help
- Accessibility
- Careers

NLM NIH HHS USA.gov