

FULL TEXT LINKS



Biol Pharm Bull. 2008 Jan;31(1):19-26. doi: 10.1248/bpb.31.19.

Inhibitory effect of electrolyzed reduced water on tumor angiogenesis

Jun Ye ¹, Yuping Li, Takeki Hamasaki, Noboru Nakamichi, Takaaki Komatsu, Taichi Kashiwagi, Kiichiro Teruya, Ryuhei Nishikawa, Takeshi Kawahara, Kazuhiro Osada, Kazuko Toh, Masumi Abe, Huaize Tian, Shigeru Kabayama, Kazumichi Otsubo, Shinkatsu Morisawa, Yoshinori Katakura, Sanetaka Shirahata

Affiliations

PMID: 18175936 DOI: [10.1248/bpb.31.19](#)

Free article

Abstract

Vascular endothelial growth factor (VEGF) is a key mediator of tumor angiogenesis. Tumor cells are exposed to higher oxidative stress compared to normal cells. Numerous reports have demonstrated that the intracellular redox (oxidation/reduction) state is closely associated with the pattern of VEGF expression. Electrolyzed reduced water (ERW) produced near the cathode during the electrolysis of water scavenged intracellular H₂O₂ and decreased the release of H₂O₂ from a human lung adenocarcinoma cell line, A549, and down-regulated both VEGF transcription and protein secretion in a time-dependent manner. To investigate the signal transduction pathway involved in regulating VEGF expression, mitogen-activated kinase (MAPK) specific inhibitors, SB203580 (p38 MAPK inhibitor), PD98059 (ERK1/2 inhibitor) and JNKi (c-Jun N-terminal protein kinase inhibitor) were applied. The results showed that only PD98059 blocks VEGF expression, suggesting an important role for ERK1/2 in regulating VEGF expression in A549 cells. As well, ERW inhibited the activation of extracellular signal-regulated kinase (ERK) in a time-dependent manner. Co-culture experiments to analyze in vitro tubule formation assay revealed that A549 cell-derived conditioned medium significantly stimulated the formation of vascular tubules in all analyzed parameters; tubule total area, tubule junction, number of tubules, and total tubule length. ERW counteracted the effect of A549 cell-conditioned medium and decreased total tube length (p<0.01). The present study demonstrated that ERW down-regulated VEGF gene transcription and protein secretion through inactivation of ERK.

Related information

[MedGen](#)

[PubChem Compound \(MeSH Keyword\)](#)

LinkOut – more resources

Full Text Sources

[J-STAGE](#), [Japan Science and Technology Information Aggregator](#), [Electronic Medical Online](#), [Meteo Inc](#)

Research Materials

[NCI CPTC Antibody Characterization Program](#)

Miscellaneous

