

FULL TEXT LINKS



Randomized Controlled Trial [Nutr Res.](#) 2008 Mar;28(3):137-43.

doi: 10.1016/j.nutres.2008.01.008.

Supplementation of hydrogen-rich water improves lipid and glucose metabolism in patients with type 2 diabetes or impaired glucose tolerance

Sizuo Kajiyama ¹, Goji Hasegawa, Mai Asano, Hiroko Hosoda, Michiaki Fukui, Naoto Nakamura, Jo Kitawaki, Saeko Imai, Koji Nakano, Mitsuhiro Ohta, Tetsuo Adachi, Hiroshi Obayashi, Toshikazu Yoshikawa

Affiliations

PMID: 19083400 DOI: [10.1016/j.nutres.2008.01.008](#)

Abstract

Oxidative stress is recognized widely as being associated with various disorders including diabetes, hypertension, and atherosclerosis. It is well established that hydrogen has a reducing action. We therefore investigated the effects of hydrogen-rich water intake on lipid and glucose metabolism in patients with either type 2 diabetes mellitus (T2DM) or impaired glucose tolerance (IGT). We performed a randomized, double-blind, placebo-controlled, crossover study in 30 patients with T2DM controlled by diet and exercise therapy and 6 patients with IGT. The patients consumed either 900 mL/d of hydrogen-rich pure water or 900 mL of placebo pure water for 8 weeks, with a 12-week washout period. Several biomarkers of oxidative stress, insulin resistance, and glucose metabolism, assessed by an oral glucose tolerance test, were evaluated at baseline and at 8 weeks. Intake of hydrogen-rich water was associated with significant decreases in the levels of modified low-density lipoprotein (LDL) cholesterol (ie, modifications that increase the net negative charge of LDL), small dense LDL, and urinary 8-isoprostanes by 15.5% ($P < .01$), 5.7% ($P < .05$), and 6.6% ($P < .05$), respectively. Hydrogen-rich water intake was also associated with a trend of decreased serum concentrations of oxidized LDL and free fatty acids, and increased plasma levels of adiponectin and extracellular-superoxide dismutase. In 4 of 6 patients with IGT, intake of hydrogen-rich water normalized the oral glucose tolerance test. In conclusion, these results suggest that supplementation with hydrogen-rich water may have a beneficial role in prevention of T2DM and insulin resistance.

Related information

[MedGen](#)

[PubChem Compound](#)

[PubChem Compound \(MeSH Keyword\)](#)

[PubChem Substance](#)

LinkOut - more resources

Full Text Sources

[ClinicalKey](#)

[Elsevier Science](#)

Other Literature Sources

[The Lens - Patent Citations](#)

Medical

[ClinicalTrials.gov](#)

[Genetic Alliance](#)

[MedlinePlus Health Information](#)