

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

CAVALRY® 9

Intervet/Merck Animal Health

Clostridium Chauvoei-Septicum-Haemolyticum-Novyi-Sordellii-Tetani-Perfringens Types C & D Bacterin-Toxoid

For Animal Use Only

INDICATIONS: Cavalry® 9 is recommended for the vaccination of healthy cattle 3 months of age or older as an aid in the prevention of disease caused by *Clostridium tetani*, *C. chauvoei*, *C. septicum*, *C. novyi* Type B, *C. haemolyticum*, *C. sordellii*, and *C. perfringens* Types C & D.

Immunity is also provided as an aid in the prevention of diseases caused by the beta and epsilon toxins of an additional organism, *C. perfringens* Type B. Although *C. perfringens* Type B is not a significant problem in North America (US), immunity is derived from a combination of Type C (beta) and Type D (epsilon) *C. perfringens* fractions.

ADMINISTRATION AND DOSAGE: Shake well. Inject 2 mL **subcutaneously** followed by a 2 mL dose in 3 weeks. Calves vaccinated at 3-6 months of age should be given a 3rd vaccination with either this vaccine, or another vaccine containing *C. perfringens* Type D epsilon toxoid, to aid in the prevention of disease caused by *C. perfringens* Type D. Revaccinate annually with 2 mL prior to periods of extreme risk or parturition.

PRESERVATIVES: Gentamicin and thimerosal.

PRECAUTIONS: This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy. This level of performance may be affected by conditions of use such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions. Transient swelling may occur post-vaccination.

CAUTION: Store at 35°-45°F (2°-7°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactic reactions may occur following use. **Antidote:** Epinephrine.

Manufactured by SCHERING-PLOUGH ANIMAL HEALTH LIMITED, UPPER HUTT, NEW ZEALAND

Distributed by INTERVET, INC., OMAHA, NE 68103 USA

U.S. Veterinary Permit No. 311

1 800 521-5767 (USA)

		Code	
10 Doses	20 mL	067691	129542
50 Doses	100 mL	058246	125205
125 Doses	250 mL	067694	131179

CPN: 1047284.2