

HUVEPHARMA, INC.
525 WESTPARK DR., SUITE 230, PEACHTREE CITY, GA, 30269
 Telephone: 770-486-7212
 Toll-Free: 877-994-4883
 Fax: 770-486-7217
 Website: www.huvepharma.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

CORID® 20% SOLUBLE POWDER

Huvepharma
(amprolium)
Coccidiostat

An aid in the treatment and prevention of bovine coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves

Active Ingredient: amprolium	20%
-------------------------------------	-----

CORID® (amprolium) 20% Soluble Powder - FOR ANIMAL USE ONLY

INDICATION: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii* in calves.

CAUTION: For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

USE DIRECTIONS: There are **220 mg** of amprolium in every **1 gram** of CORID 20% Soluble Powder.

5-Day Treatment Protocol	21-Day Prevention Protocol
Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

1 oz = 3 1/2 measuring tablespoonfuls

1 oz = 28.35 grams

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

Stock solution for water proportioners may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/50 gal. One 10 oz pack to 125 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1 bag (10 oz) in 1 gallon of water to make stock solution. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/100 gal. One 10 oz pack to 250 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1/2 bag (5 oz) in 1 gallon of water to make stock solution. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 oz CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 oz of CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

NUMBER OF CALVES ONE PACKET WILL TREAT	5-Day Treatment	21-Day Prevention
100 lb calves	25	12

500 lb calves	5	2.5
---------------	---	-----

RESIDUE WARNING: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

NOT FOR HUMAN USE. Restricted Drug (California) - Use only as directed. Keep this and all drugs out of the reach of children.

STORAGE: Store between 5° - 25°C (41° - 77°F) with brief excursions to 40°C (104°F). Avoid high humidity storage conditions.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

® **CORID is a registered trademark of Huvepharma EOOD.**

Distributed by:

Huvepharma, Inc., Peachtree City, GA 30269

For questions, please call 1-877-994-4883

Approved by FDA under NADA # 033-165

Net Contents:		
10 oz. (238.5 grams)	KCP-S00015	Version:02 Rev:03-2022
12x10 oz Packets Display Tray	KCP-S00016	Version: 02 Rev. 02-2022
24 Packets x 10 oz. each (238.5 grams) Net Wt.: 15 lbs. (6.8 kg)	KCP-S00017	Version: 02 Rev. 02-2022

CPN: 1309059.1