

ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: www.zoetis.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

FluSure XP®/RespiSure-ONE®

Zoetis

Swine Influenza Vaccine, H1N1 & H1N2 & H3N2

Killed Virus

Mycoplasma Hyopneumoniae Bacterin

For use in swine only

For veterinary use only

H1N1 GAMMA

H1N2 DELTA-1

NOW WITH H3N2 CLUSTER-IV-A

NOW WITH H3N2 CLUSTER-IV-B

This product has been shown to be effective for the vaccination of healthy swine, including pregnant sows and gilts, 3 weeks of age or older against respiratory disease caused by swine influenza A virus (SIV) types H1N1, H1N2, and H3N2, and *Mycoplasma hyopneumoniae*. A duration of immunity of at least 10 weeks has been demonstrated against SIV type H3N2 and at least 25 weeks against *M. hyopneumoniae*. Duration of immunity of the remaining fractions has not been established. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

This product contains the following influenza A virus isolates: A/Swine/Iowa/110600/2000(H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2); A/Swine/North Carolina/ 394/2012(H3N2); and A/Swine/Minnesota/872/2012 (H3N2).

Directions: Shake diluent before use. Aseptically rehydrate the freeze-dried vaccine with the liquid bacterin provided, shake well, and administer 2 mL intramuscularly. Administer a single 2-mL dose to healthy swine, followed by a single dose of FluSure XP approximately 3 weeks later. In young pigs, vaccinate after maternally derived antibodies to SIV have declined. For advice on revaccination, contact your veterinarian or the manufacturer.

Precautions: Store at 2°-8°C. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Contains gentamicin and thimerosal as preservatives. In case of allergic reaction, administer epinephrine. Do not mix with other products, except as specified above. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

VLN 190/PCN

Zoetis Inc.

Kalamazoo, MI 49007, USA

50-dose vial of vaccine, rehydrate to 100 mL	4995.2D	40038072
100-mL vial of liquid bacterin for use as diluent		

CPN: 3690251.7