

ZOETIS INC.**333 PORTAGE STREET, KALAMAZOO, MI, 49007**

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: www.zoetis.com

THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

FluSure XP®**Zoetis****Swine Influenza Vaccine, H1N1 & H1N2 & H3N2****Killed Virus**

For use in swine only

For veterinary use only

H1N1 GAMMA**H1N2 DELTA-1****NOW WITH H3N2 CLUSTER-IV-A****NOW WITH H3N2 CLUSTER-IV-B**

This product has been shown to be effective for the vaccination of healthy swine, including pregnant sows and gilts, 3 weeks of age or older against swine influenza A H1N1, H1N2, and H3N2 virus infection. Duration of immunity of at least 10 weeks has been established against swine influenza A virus H3N2; duration of immunity against H1N1 and H1N2 has not been established. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

This product contains the following Influenza A Virus isolates: A/Swine/Iowa/110600/2000 (H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2); A/Swine/North Carolina/A01270394/2012 (H3N2); and A/Swine/Minnesota/A01270872/2012 (H3N2).

A sterile diluent containing Amphigen® is used to rehydrate the freeze-dried vaccine.

Directions: Shake diluent before use. Aseptically rehydrate the freeze-dried vaccine with the accompanying adjuvant-containing sterile diluent, shake well, and administer 2 mL intramuscularly. Healthy swine 3 weeks of age or older should receive two 2-mL doses administered approximately 3 weeks apart. In young pigs, vaccinate after maternally derived antibodies to swine influenza A viruses have declined. For advice on revaccination, consult a veterinarian or the manufacturer.

Precautions: Store at 2°-8°C. Do not freeze. Use entire contents when first opened or rehydrated. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Contains gentamicin as preservative. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

VLN 190/PCN 19A5.2C

Zoetis Inc., Kalamazoo, MI 49007, USA

50-dose vial of vaccine Rehydrate to 100 mL 100 mL vial of sterile diluent	40036618
250-dose vial of vaccine Rehydrate to 500 mL 500 mL vial of sterile diluent	40036622

CPN: 3690246.7