

BOEHRINGER INGELHEIM ANIMAL HEALTH USA INC.

3239 SATELLITE BLVD., BLDG 500, DULUTH, GA, 30096

Telephone: 800-325-9167

Customer Service: 888-637-4251

Technical Service: 888-637-4251

Fax: 816-236-2717

Website: www.boehringer-ingelheim.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

Ingelvac CircoFLEX®

Boehringer Ingelheim Animal Health

Porcine Circovirus Vaccine

Type 2, Killed Baculovirus Vector

This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against porcine circovirus type 2 (PCV2). The onset of protection is 2 weeks post-vaccination. The duration of immunity is at least 4 months. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

The product contains PCV2a, and has been shown to be effective against lymphoid depletion, inflammation and colonization of lymphoid tissue, and the magnitude of viremia.

Directions and dosage: Shake well before use. Using aseptic technique, administer a single **1 mL** dose by intramuscular injection.

Precautions: Store out of direct sunlight at 35 - 46°F (2 - 8°C). Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Do not mix with other products. In case of anaphylactoid reaction, administer epinephrine. This product has not been tested in pregnant animals. In case of human exposure, contact a physician.

Preservative: Neomycin.

Boehringer Ingelheim Animal Health USA Inc., St. Joseph, MO 64506

Phone: 1 (888) 637-4251

VLN/PCN 124/19K5.R0

50 Doses/50 mL	130902-07
100 Doses/100 mL	130903-07
250 Doses/250 mL	130904-07
500 Doses/500 mL	1309169-00

CPN: 1028162.6