

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

Prestige® WNV

Intervet/Merck Animal Health

WEST NILE VIRUS VACCINE, KILLED FLAVIVIRUS CHIMERA

A vaccine containing inactivated, concentrated, adjuvanted, tissue culture origin West Nile virus - flavivirus chimera.

INDICATIONS: This product has been shown to be effective for the vaccination of healthy horses 6 months of age or older against West Nile Virus. Duration of Immunity has not been established. For more information regarding safety and efficacy data, go to productdata.aphis.usda.gov.

This product has been shown to be effective against encephalitis and viremia caused by West Nile Virus.

DOSE: For primary vaccination aseptically administer 1 mL intramuscularly, repeat with a single dose in 3 - 4 weeks. Historically, annual revaccination with this product has been recommended. The need for this booster has not been established. The presence of maternal antibody is known to interfere with the development of active immunity in foals and additional boosters will be required in most young animals. For more information on revaccination frequency, consult your veterinarian.

CAUTIONS: Store at 2°-8°C (35°-46°F). Shake well before using. Use entire contents when first opened. Do not mix with other products. This product has not been tested in pregnant animals. Do not vaccinate within 21 days before slaughter. Inject deep into the muscle only. Local reactions may occur at the injection sites in vaccinates, and most of these reactions resolve in 3 to 7 days following vaccination. Systemic reactions may also occur in animals following vaccination. If allergic reaction occurs, treat with epinephrine. Care should be used to avoid injecting your own fingers, hands, or other body parts. Accidental injection can cause serious local reactions. In case of human exposure, contact a physician.

ANTIBIOTIC: Gentamicin.

PRESERVATIVE: Thimerosal.

FOR ANIMAL USE ONLY

Intervet Inc. d/b/a Merck Animal Health, Omaha, NE 68103, USA

VLN 165A/PCN 1995.R2

1 800 521-5767 (USA)

1 866 683-7838 (CAN)

For patent information:

<http://www.merck.com/product/patent/home.html>

		Code	
10 Doses	10 x 1 mL Syringes with Separate Sterile Needles	182544	195933-02
10 Doses	10 mL	193718	182505-02

CPN: 1047544.1