

ELANCO US, INC.

Farm Animal Business

Distributed by ELANCO US, INC.

1447 140TH ST., LARCHWOOD, IA, 51241

Customer Service: 800-782-8977

Main Switchboard: 317-433-4800

Technical Service: 800-843-4441

Website: www.elanco.us

Email: elanco@elanco.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

RHINI SHIELD™ TX4

Elanco (Farm Animal)

Bordetella Bronchiseptica-Erysipelothrix Rhusiopathiae-Pasteurella Multocida Bacterin-Toxoid

For Use In Animals Only

Nursery

Grow

Gilt

Sow

INDICATIONS: This product has been shown to be effective for vaccination of healthy swine, 7 days of age or older, against atrophic rhinitis caused by *Bordetella bronchiseptica* or the toxin of *Pasteurella multocida* Types A and D, erysipelas caused by *Erysipelothrix rhusiopathiae*, and pneumonia caused by *Pasteurella multocida* Type A. The duration of immunity for this product has not been established. For more information regarding efficacy and safety data, see *productdata.aphis.usda*.

ADJUVANT: Aluminum hydroxide.

DIRECTIONS: Shake well before using. Administer intramuscularly or subcutaneously. Four doses must be given (two gilt/sow doses during pregnancy and two piglet doses). Vaccinate gilts and sows with a 5 mL dose at 5 and 2 weeks prior to farrowing. Vaccinate piglets from vaccinated dams with a 1 mL dose at 7 to 10 days of age and a 2 mL dose 2 weeks later. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended. For advice on revaccination frequency, in general or in the face of maternal antibody, stress, or a disease exposure, consultation with a veterinarian is recommended.

PRECAUTIONS: Store out of direct sunlight at 2 to 8°C (35° to 46°F). DO NOT FREEZE. Do not mix with other products, except as specified on the label. Use entire contents when first opened. Do not vaccinate within 21 days prior to slaughter. In case of human exposure, contact a physician. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

VLN/PCN 196/7054.01

Elanco US Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

Rhini Shield, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

20 Gilt/Sow Doses	100 mL	YL102735X
50/100 Piglet Doses		

CPN: 1114037.9