MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

 Customer Service:
 800-521-5767

 Order Desk:
 800-648-2118

 Technical Service (Companion Animal):
 800-224-5318

 Technical Service (Livestock):
 800-211-3573

 Fax:
 973-937-5557

Website: www.merck-animal-health-usa.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

SABER™

Intervet/Merck Animal Health

POUR-ON INSECTICIDE FOR BEEF CATTLE & CALVES

For Control of Horn Flies and Lice

Season-Long Lice Control† with One Application

May be used with Automatic Dosing System

ACTIVE INGREDIENT:	% [W/W]
Lambdacyhalothrin: [1α-(S), 3α(Z)]-(±)-cyano (3-phenoxyphenyl)-methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate	1.0%
INERT INGREDIENTS	99.0%
TOTAL	100.0%

Keep Out of Reach of Children

CAUTION

EPA Reg. No. 773-74 EPA Est. No. 618-LA-1 U.S. Patent No. 5,045,536

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE - No dilution necessary. Do not apply to lactating or dry dairy cows. Do not apply this product to face of beef cattle or calves.

Apply To	Target Insects	Dosage	
Beef Cattle and Calves	Lice, Horn Flies	Apply product down the backline at the rate of	
		Less than 600 lbs	Greater than 600 lbs
		10 mL (1/3 fl oz)	15 mL (1/2 fl oz)

Repeat treatment as needed; however, do not apply more than once every 2 weeks and do not apply more often than four times within any 6-month period.

Special Note: SABER™ Pour-On is not recommended for use on veal calves. SABER Pour-On is not effective in controlling cattle grubs. Therefore, this product can be used on cattle at any time of the year without fear of host-parasite reactions commonly associated with grub treatment products. SABER Pour-On should be used in an integrated pest management system, which may involve repeat treatments and the use of other pest control practices. Continual exposure of horn flies to a single class of insecticide, (eg, pyrethroids or organophosphates) may lead to the development of resistance to that class of insecticide. In order to reduce the possibility of horn flies developing resistance, it is important to rotate, on a seasonal basis, the class of insecticide used and/or the method of horn fly control. For advice concerning current control practices with relation to specific local conditions, consult resources in resistance management programs and your Cooperative Agricultural Extension Service.

PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed, absorbed through the skin, or inhaled. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, or clothing. Remove contaminated clothing and wash

before reuse. Wash thoroughly with soap and water after handling. Avoid contact with eyes. Wear rubber or nonpermeable protective gloves when applying this product. Data indicate this product to be a contact allergenic. Do not apply product to face of cattle.

FIRST AID

If inhaled

- · Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

If On Skin Or Clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If In Eves

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If Swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the Rocky Mountain Poison Center 303-595-4869 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS - This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of rinse water. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS - Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

†Season-long lice control from January-April.

NOTICE OF WARRANTY

MERCK ANIMAL HEALTH MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THIS PRODUCT UNDER NORMAL CONDITIONS AND IN ACCORDANCE WITH THE STATEMENTS ON THE LABEL.

Copyright © 1997, 1999, Intervet Inc, a subsidiary of Merck & Co., Inc., 2 Giralda Farms, Madison, NJ 07940 All rights reserved.

Net Contents:				
30 fl oz (900 mL) Squeeze 'N' Measure™ Bottle	0061-5062-01	130555 R2 10/12 183079 R1 Rev. 08/18		
Treats 90 550 lb Cattle				
One US Gallon (3.785 L)	0061-5062-04	127312 R2 Rev. 10/12 155084 R1 Rev. 08/15		
Treats 378 550 lb Cattle				

CPN: 1047174.7