

- [Ingredients](#)
- [Charts](#)
- [Information](#)



[Home](#) » [ELANCO US INC.](#) » truFel™ HCP+FeLV (ELANCO US INC.)

[Label](#)

[Use/Dose](#)

ELANCO US INC.

Distributed by ELANCO US, INC.

FORT DODGE, IA, 50501

Customer Service: 317-276-1262

Technical Service: 800-428-4441

Website: www.elanco.us

Email: elanco@elanco.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

TRUFEL™ HCP+FeLV

Elanco US

Feline Leukemia-Rhinotracheitis-Calici-Panleukopenia Vaccine


Modified Live & Killed Virus

For use in animals only

This product has been shown to be effective for the vaccination of healthy cats, 8 weeks of age or older, against feline leukemia, rhinotracheitis, calici and panleukopenia viruses. The duration of immunity for this product has not been determined. For more information regarding efficacy and safety data, go to productdata.aphis.usda.gov.

Directions for Use: Aseptically rehydrate with accompanying diluent. Inject (one) 1 mL dose subcutaneously using aseptic technique. Healthy cats should receive two doses, 3 to 4 weeks apart, except that if the animal is less than 16 weeks of age, the final dose should be given no earlier than 16 weeks of age. The presence of maternal antibody is known to interfere with the development of active immunity in cats and additional boosters will be required in most young animals. Historically, annual revaccination with a single dose has been recommended for this product. The need for this booster has not been established.

For advice on revaccination frequency and annual booster vaccinations, consultation with a veterinarian is recommended. **Do not vaccinate pregnant queens.**



Caution: Store in the dark at 2° to 8°C (35° to 46°F). AVOID FREEZING. SHAKE WELL. Do not mix with other products, except as specified on the label. In case of human exposure, contact a physician. In case of anaphylactoid reaction, administer epinephrine. Inactivate all unused contents before disposal.

Thimerosal, neomycin, polymyxin B and amphotericin B added as preservatives.

Caution: In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package.

Elanco US Inc., Fort Dodge, IA 50501

Phone: 888-545-5973

VLN/PCN 196/15B9.21

TruFel, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2021 Elanco or its affiliates.

25 Doses	25 - 1 mL Vials of Vaccine plus 25 - 1 mL Vials of Diluent	YL103546A
----------	--	-----------

CPN: 1943032.4

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Copyright 2024 - DVMetrics & Services LLC