

Prepare fresh solutions daily.
Mixing Guide for low usage
(high level supplementation):

Vi-tal / Water
¼ tsp. / 1 gallon
1¼ tsp. / 5 gallons
2½ tsp. / 10 gallons

Consumption Guide:

A 6 oz. packet per 128 gallons of
water will treat the following:

CHICKENS (Age)
1 week..... 24,000
3 weeks..... 6,900
5 weeks..... 4,200
8 weeks..... 2,580

REPLACEMENT PULLETS

9 weeks..... 6,000
12+ weeks..... 5,330

LAYING HENS..... 1,800


DUCKS (Age)
1 to 3 weeks..... 2,500

TURKEYS (Age)
1 week..... 14,730
3 weeks..... 4,890
6 weeks..... 2,190
10 weeks..... 1,240
15+ weeks..... 1,040

HOGS
25 lbs..... 350
50 lbs..... 200
100 lbs..... 100

NOTE: In hot weather with
higher water consumption,
divide these numbers in half.

Product No. 148178



Manufactured for: Neogen Corporation
944 Nandino Blvd, Lexington, KY 40511
859-254-1221 or 800-621-8829
animalsafety@neogen.com AN065-0818



Net Contents:
6 ounces (170.1 grams)



Vi-tal

A Water Vitamin, Electrolyte and Mineral Supplement

To provide supplemental vitamins, electrolytes and trace minerals to
aid in growth and productivity; to stimulate water consumption; and to
maintain balance of body fluids in livestock and poultry.

GUARANTEED ANALYSIS:

Salt (NaCl), not less than.....	45.0%
Salt (NaCl), not more than.....	50.0%
Sodium (Na), not less than.....	22.0%
Sodium (Na), not more than.....	26.0%
Potassium (K), not less than.....	10.0%
Magnesium (Mg), not less than.....	1.0%
Vitamin A, not less than.....	6,900,000 IU/lb
Vitamin D-3, not less than.....	1,600,000 IU/lb
Vitamin E, not less than.....	2,100 IU/lb

INGREDIENTS:

Salt, Sodium Bicarbonate, Potassium Chloride, Magnesium Sulfate, Vitamin A Supplement, Niacin Supplement, Vitamin E Supplement, d-Calcium Pantothenate, Vitamin D₃ Supplement, Menadione Sodium Bisulfite Complex (source of Vitamin K Activity), Riboflavin, Vitamin B₁₂ Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Manganese Sulfate, Folic Acid, Cobalt Sulfate, Ethylenediamine Dihydriodide, Copper Sulfate, Zinc Sulfate, Calcium Gluconate.

MIXING DIRECTIONS:

High level supplementation: Mix 1 packet per 128 gallons of water.

Moderate level supplementation: Mix 1 packet per 256 gallons of water.

Low level supplementation: Mix 1 packet per 384 gallons of water.

NOTE: For water proportioners delivering 1 ounce of stock solution per gallon of drinking water, the high, moderate and low levels outlined above are for 1 packet per 1 gallon, 2 gallons and 3 gallons of stock solution, respectively.

Offer for 3 to 5 days —

When receiving and placing poultry and livestock; during stress due to scours, vaccination, castration, and debeaking; during temperature extremes; to stimulate appetite through improved nutrition (use the high or moderate level as outlined in the mixing directions).

Offer for 1 to 2 days —

Before and after transportation (use the high level as outlined in the mixing directions).

Supplementation for 10 days or more —

For aid in nutritional build-up, especially when feed intake is below normal (use low level as outlined in the mixing directions).

Feed Supplementation —

Thoroughly mix 2 packets per ton of complete feed. Use for 5 or more days.

⚠ WARNING: This product can expose you to Cobalt Sulfate, which is known to the State of California to cause cancer. For more information go to www.P65Warnings.ca.gov.