

## ELANCO US, INC.

Farm Animal Business

Distributed by ELANCO US, INC.

1447 140TH ST., LARCHWOOD, IA, 51241

Customer Service: 800-782-8977

Main Switchboard: 317-433-4800

Technical Service: 800-843-4441

Website: [www.elanco.us](http://www.elanco.us)

Email: [elanco@elanco.com](mailto:elanco@elanco.com)



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

---

### VIRA SHIELD™ 6+VL5

*Elanco US (Farm Animal)*

#### **Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza 3-Respiratory Syncytial Virus Vaccine**

Killed Virus

#### **Campylobacter Fetus-Leptospira Canicola-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin For Use In Animals Only**

**INDICATIONS:** This product has been shown to be effective for the vaccination of healthy cattle against disease caused by infectious bovine rhinotracheitis (IBR), bovine virus diarrhea (BVD Type 1 and BVD Type 2), parainfluenza Type 3 (PI3), and bovine respiratory syncytial (BRSV) viruses and *Campylobacter fetus*, *Leptospira canicola*, *grippotyphosa*, *hardjo*, *icterohaemorrhagiae*, and *pomona*. The duration of immunity against BVD Type 2 is as at least 11 months and *C. fetus* is at least one year. The duration of immunity for the other antigens has not been established. For more information regarding efficacy and safety data, see [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

Produced from non-cytopathic (BVD Type 1 and BVD Type 2) and cytopathic (BVD Type 1) isolates. This product is safe for use in pregnant cows and heifers.

**ADJUVANT:** Xtend™ SP.

**DIRECTIONS:** Shake well before using. Administer 5 mL subcutaneously 2 to 4 weeks prior to breeding. In accordance with Beef Quality Assurance guidelines, this product should be administered subcutaneously (under the skin) in the neck. Revaccinate with Vira Shield 6 (Code 1185.20) in 4 to 5 weeks. Historically, annual revaccination has been recommended for this product. The need for this booster has not been established. For advice on revaccination, contact a veterinarian.

**PRECAUTIONS:** Store out of direct sunlight at 2° to 8°C (35° to 46°F). DO NOT FREEZE. Use entire contents when first opened. Do not mix with other products, except as specified on the label. In case of human exposure, contact a physician. Do not vaccinate within 60 days prior to slaughter. Transient swelling may occur at the site of injection. Milk reduction and transient depression may be observed in lactating dairy cows for 3 to 6 days following vaccination. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

Contains amphotericin B, gentamicin, and thimerosal as preservatives.

**VLN/PCN** 196/44B6.20

Elanco US, Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

Vira Shield, Xtend, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

<b>10 Doses</b>	<b>50 mL</b>	YL102481A
<b>50 Doses</b>	<b>250 mL</b>	YL102481B

**CPN:** 1114096.5