

MERCK ANIMAL HEALTH

Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

Vista® 5 VL5 SQ

Intervet/Merck Animal Health

BOVINE RHINOTRACHEITIS-VIRUS DIARRHEA-PARAINFLUENZA 3-RESPIRATORY SYNCYTIAL VIRUS VACCINE, Modified Live Virus, CAMPYLOBACTER FETUS-LEPTOSPIRA CANICOLA-GRIPPOTYPHOSA-HARDJO-ICTEROHAEMORRHAGIAE-POMONA BACTERIN

Cattle Vaccine

Product Description: The reconstituted vaccine-bacterin product contains modified-live cultures of bovine rhinotracheitis (IBR) virus; bovine virus diarrhea (BVD) virus (Types 1 and 2); parainfluenza₃ virus (PI₃) and bovine respiratory syncytial virus (BRSV) and inactivated cultures of *Campylobacter fetus* and *Leptospira canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona* with a proprietary adjuvant.


Indications: This product is for the vaccination of healthy cows and heifers prior to breeding as an aid in the reduction of abortion due to infectious bovine rhinotracheitis (IBR); as an aid in the prevention of fetal infection, including persistently infected calves caused by bovine virus diarrhea virus (BVD) (Types 1 & 2). Reproductive Duration of Immunity (DOI) has been demonstrated to be at least 217 days for IBR and at least 206 days for BVD (Types 1 & 2). In addition, it can be used as an aid in the prevention of disease caused by IBR, BVD (Type 2); and bovine respiratory syncytial virus (BRSV), as an aid in the control of disease caused by BVD (Type 1), and parainfluenza₃ virus (PI₃), as an aid in reducing infertility (reproductive disease caused by *Campylobacter fetus*), and as an aid in preventing leptospirosis (caused by *Leptospira canicola*, *L. grippotyphosa*, *L. hardjo* - including the *L. borgpetersenii* serovar *hardjo bovis*, *L. icterohaemorrhagiae*, and *L. pomona*) and as an aid in prevention of urinary shedding of hardjo organisms. Respiratory Duration of Immunity (DOI) has been demonstrated to be at least 182 days for IBR, at least 206 days for BVD (Type 1), and at least 200 days for BVD (Type 2). Safe for use in pregnant heifers and cows or calves nursing pregnant cows provided the cows and heifers in the herd are vaccinated prior to breeding, within the previous 12 months, with any of the modified live IBR and BVD containing vaccine(s) in this product line.

Mixing Directions: 10 Doses - Rehydrate freeze dried vial of Vista® 5 SQ with accompanying vial of VL5 SQ. Mix reconstituted vial well.

50 Doses - Rehydrate freeze dried vial of Vista® 5 SQ with part of the accompanying vial of VL5 SQ using the transfer needle provided (see pictorial directions). Mix reconstituted vial well and transfer rehydrated vaccine into VL5 SQ vial using transfer needle. Remove transfer needle from VL5 SQ vial and mix reconstituted vial well. Peel label from bottle of Vista® 5 SQ and place on VL5 SQ vial containing all vaccine.

Use Directions: Inject 2.0 mL subcutaneously 14-60 days prior to breeding. Annual revaccination is recommended. A revaccination dose can be administered at more frequent intervals based upon individual farm disease risk assessment or any time epidemic conditions exist or are reported. Consult your veterinarian.

Mixing Instructions



CAUTION: THE TRANSFER NEEDLE, INCLUDED IN THE CARTON PACKAGING, IS SHARP AND MAY CAUSE INJURY TO SELF OR ANIMALS IF NOT HANDLED OR DISPOSED OF PROPERLY.

1. Insert needle beyond breather valve into the vial of lyophilized vaccine to remove vacuum.
2. Turn the vial of lyophilized vaccine upside down and fully insert needle into the bacterin diluent bottle. Be sure to clear cap with breather valve.
3. Turn attached vial and bottle so the vial of lyophilized vaccine is underneath the bacterin diluent bottle and squeeze enough bacterin diluent into the vial of lyophilized vaccine to rehydrate the vaccine. Mix well.
4. Turn attached bottles so rehydrated vaccine is above the bacterin diluent bottle. Squeeze air into rehydrated vaccine vial.
5. Squeeze and release the bacterin diluent bottle until all the solution in the rehydrated vaccine vial drains into the bacterin diluent bottle.
6. Separate the vial of lyophilized vaccine and needle from the bacterin diluent bottle. Using the tab, remove the back of the vaccine vial label. Place the separated label over the bacterin diluent bottle label to accurately identify the newly created solution.

Cautions: Store at 2°-7°C (35°-45°F). Do not freeze. Use immediately after reconstitution; do not save partial contents. Burn containers and all unused product. Use only in healthy cattle. Do not vaccinate within 21 days before slaughter. Fetal health risks associated with vaccination of pregnant animals with modified live vaccines cannot be unequivocally determined by clinical trials conducted for licensure. Management strategies based on vaccination of pregnant animals with modified live vaccines should be discussed with a veterinarian. If allergic reaction occurs, treat with epinephrine. Contains penicillin, streptomycin and thimerosal as preservatives.

FOR ANIMAL USE ONLY

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103

U.S. Vet Lic. No 165A

1 800 521-5767 (USA)

For patent information:

<http://www.merck.com/product/patent/home.html>

		Code	
--	--	------	--

10 Doses	20 mL	006341	142036-07
50 Doses	100 mL	006342	144899-07

CPN: 1047456.4