

Label Use/Dose

BOEHRINGER INGELHEIM ANIMAL HEALTH USA INC.

3239 SATELLITE BLVD., BLDG 500, DULUTH, GA, 30096

Telephone: 800-325-9167

Customer Service: 888-637-4251

Technical Service: 888-637-4251

Fax: 816-236-2717

Website: www.boehringer-ingelheim.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

BOVIKALC® DRY

Boehringer Ingelheim Animal Health

Anionic Mineral Feed Supplement

For dairy cows at drying off

CONTAINS AMMONIUM CHLORIDE

Product Characteristics: Bovikalc® Dry is an oral mineral supplement to be given to dairy cows at drying off.

Actions: Following oral administration, the bolus dissolves into the rumen within 30 minutes and the anionic salts are released and then absorbed, which impacts the cow's acid-base balance.

Ingredients: Calcium chloride, calcium sulfate, water, ammonium chloride, vegetable fat, a-tocopherol acetate, fruit juice color, Vitamin D3, sodium selenite.

Guaranteed Analysis:

Equivalent Crude Protein from Non Protein Nitrogen, max	20%
Calcium, min	18.3%
Calcium, max	20%
Selenium, min	9.6 ppm

How to use Bovikalc Dry?

To be used in dairy cows at drying off.

Give 2 boluses at last milking or 8-12 hours before the last milking.

Do not administer more than 4 boluses per 24 hours.

Please discuss protocol with Herd Nutritionist and/or Herd Veterinarian before implementation.

Place the bolus in the bolus applicator with the rounded end pointing outward.

If right-handed, place yourself at the cow's right side.

Grab the cow's upper jaw with your left hand. Keep the cow's head at a natural position and a little towards yourself. Gently guide the loaded BOVIKALC applicator into the cow's mouth.

When you feel resistance, gently guide the applicator over the tongue. When the applicator is behind the tongue, squeeze the handle to release the bolus.

Wait a moment for the cow to swallow and then remove the applicator.



Precautions:

CAUTION: USE ONLY AS DIRECTED. FOR CATTLE ONLY.

Follow label instructions carefully. Incorrect use or handling may cause serious side effects.

User assumes risk of use or handling contrary to instructions, or under abnormal conditions or conditions not reasonably foreseeable.

Only administer using the Bovikalc branded applicator.

Do not give Bovikalc Dry to cows lying down.

Do not administer Bovikalc Dry if the coating on the bolus becomes damaged or the bolus is broken.

Keep out of the reach of children.

Bovikalc Dry is not for human use.

Storage:

Store at or below 30°C (86°F).

The bolus should be stored in the original plastic tube until immediately before use, as it is susceptible to humidity.

Do not use broken or damaged boluses.

Please note the caution on the packaging.

Bovikalc is a registered trademark of Boehringer Ingelheim GmbH, used under license.

Manufactured in Denmark by:

Boehringer Ingelheim Animal Health Denmark A/S

Marketed by:

Boehringer Ingelheim Animal Health USA Inc., Duluth, GA 30096

Toll Free 1-888-637-4251

MADE IN DENMARK

GTIN: (01)05703655003185

Net Contents:	
4 Boluses	151592-001
48 Boluses	151596-002
FRAGILE: HANDLE WITH CARE	

CPN: 1028337.0

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).