

Label Use/Dose

## ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: [www.zoetis.com](http://www.zoetis.com)



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

---

## FluSure XP®/FarrowSure® GOLD B

### Zoetis

Parvovirus-Swine Influenza Vaccine, H1N1 & H1N2 & H3N2

Killed Virus

Erysipelothrix Rhusiopathiae-Leptospira Bratislava-Canicola-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin

For use in swine only

For veterinary use only

**H1N1 GAMMA**

**H1N2 DELTA-1**

**NOW WITH H3N2 CLUSTER-IV-A**

**NOW WITH H3N2 CLUSTER-IV-B**

This product has been shown to be effective for the vaccination of healthy, breeding swine, including pregnant sows and gilts, against reproductive failure caused by porcine parvovirus; respiratory disease caused by swine influenza A virus (SIV) types H1N1, H1N2, and H3N2; and disease caused by *Erysipelothrix rhusiopathiae*, *Leptospira bratislava*, *L. canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae*, and *L. pomona*. A duration of immunity of at least 18 weeks has been demonstrated against *E. rhusiopathiae*. Duration of immunity against the remaining fractions has not been established. For more information regarding efficacy and safety data, go to [productdata.aphis.usda.gov](http://productdata.aphis.usda.gov).

This product contains the following Influenza A Virus isolates:

A/Swine/Iowa/110600/2000 (H1N1); A/Swine/Oklahoma/0726H/2008 (H1N2);

A/Swine/North Carolina/394/2012 (H3N2); and

A/Swine/Minnesota/872/2012 (H3N2).

**Directions:** Aseptically rehydrate the freeze-dried vaccine with the liquid component provided, shake well, and administer 2 mL intramuscularly. Healthy swine should receive 2 doses 3-5 weeks apart with the second dose administered 2-4 weeks prior to breeding. Healthy gilts, however, should receive the second dose as near as possible to 14 days prior to breeding. For advice on revaccination, consult a veterinarian or the manufacturer.

**Precautions:** Store at 2°-8°C. Do not freeze. Use entire contents when first opened or rehydrated. Sterilized syringes and needles should be used to administer this vaccine. Do not vaccinate within 21 days before slaughter. Contains gentamicin and thimerosal as preservatives. Temporary swelling at the injection site may be observed. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy. Do not mix with other products, except as specified above. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

VLN 190/PCN

Zoetis Inc.

Kalamazoo, MI 49007, USA

50-dose vial of vaccine, rehydrate to 100 mL	49F6.2B	40036883
100-mL vial of liquid component for use as diluent		

**CPN:** 3690249.6

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

Copyright 2024 - DVMetrics & Services LLC