



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

REPROSTAR™ VL5 HB

Elanco US (Farm Animal)

Campylobacter Fetus-Leptospira Canicola-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin

For Use In Animals Only

INDICATIONS: This product has been shown to be effective for the vaccination of healthy cattle against disease caused by *Campylobacter fetus*, *Leptospira canicola*, *grippotyphosa*, *hardjo-bovis*, *icterohaemorrhagiae* and *pomona*. The duration of immunity for *C. fetus* and *L. hardjo-bovis* is at least 12 months. The duration of immunity for the other antigens in this product has not been established. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

The product has been shown to be effective against kidney colonization, urinary shedding, and reproductive tract infection caused by *L. hardjo-bovis*. Vaccinated animals subsequently exposed to *L. borgpetersenii* serovar *hardjo* (type *hardjo-bovis*) have been shown to clear renal infections within 8 weeks of exposure.

ADJUVANT: Xtend™ SP.

DIRECTIONS: Shake well before using. Administer 2 mL intramuscularly 2 to 4 weeks prior to breeding. Revaccinate in 4 to 6 weeks with ReproSTAR L5 HB. (code 2665.07) Historically, annual revaccination has been recommended for this product. The need for this booster has not been established. For advice on revaccination frequency consultation with a veterinarian is recommended.

PRECAUTIONS: Store out of direct sunlight at 2° to 8°C (35° to 46°F). DO NOT FREEZE. This product has not been tested in pregnant animals. Use entire contents when first opened. Do not mix with other products, except as specified on the label. In case of human exposure, contact a physician. Do not vaccinate within 60 days prior to slaughter. Transient swelling may occur at the site of injection. Milk reduction and transient depression may be observed in lactating dairy cows for 3 to 6 days following vaccination. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine. Contains thimerosal as a preservative.

VLN/PCN: 196/2863.05

Elanco US Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

ReproSTAR, Xtend, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

10 Doses	20 mL	YL102762A
50 Doses	100 mL	YL102762B

CPN: 1114107.4

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Copyright 2024 - DVMetrics & Services LLC