



ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: www.zoetis.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

UltraChoice® 8

Zoetis

Clostridium Chauvoei-Septicum-Haemolyticum-Novyi-Sordellii-Perfringens Types C & D Bacterin-Toxoid

For veterinary use only

This product has been shown to be effective for the vaccination of healthy cattle and sheep against disease caused by *Clostridium chauvoei*, *Cl. septicum*, *Cl. haemolyticum*, *Cl. novyi*, *Cl. sordellii*, and *Cl. perfringens* types C and D. Duration of immunity has not been established. For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Although *Cl. perfringens* type B is not a significant problem in North America, immunity is provided by the beta toxoid of type C and the epsilon toxoid of type D.

Directions: Mix well. *Cattle:* Aseptically administer 2 mL subcutaneously. In accordance with Beef Quality Assurance guidelines, this product should be administered SC under the skin. Healthy cattle should receive 2 doses administered 4-6 weeks apart. *Sheep:* Aseptically administer 1 mL subcutaneously in the neck, followed by a second dose 4-6 weeks later. For more information on revaccination and in the face of *Cl. haemolyticum* exposure, contact your veterinarian.

Precautions: Store at 2°-8°C. Do not freeze. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Contains formalin as preservative. Temporary local swelling at injection site may occur after administration. Field reports indicate that a transient reduction in milk production may occur following vaccination. This product has not been tested in pregnant animals. As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy. Do not mix with other products. In case of human exposure, contact a physician.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471.

VLN 190/PCN 7160.02

Zoetis Inc., Kalamazoo, MI 49007, USA

10 cattle doses 20 sheep doses	20 mL	
50 cattle doses 100 sheep doses	100 mL	
250 cattle doses 500 sheep doses	500 mL	40033096

CPN: 3690077.3

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party

data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

[Contact Us](#) - [Logout](#)

Copyright 2024 - DVMetrics & Services LLC