

Label Use/Dose

BOEHRINGER INGELHEIM ANIMAL HEALTH USA INC.

3239 SATELLITE BLVD., BLDG 500, DULUTH, GA, 30096

Telephone: 800-325-9167

Customer Service: 800-243-0127

Fax: 816-236-2717

Website: www.boehringer-ingelheim.com



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

VETERA® Gold^{XP}

Boehringer Ingelheim Animal Health

Encephalomyelitis-Rhinopneumonitis-Influenza-West Nile Virus Vaccine

Eastern & Western, Killed Virus

Tetanus Toxoid

Indications: This product has been shown to be effective for the vaccination of healthy horses 4 months of age or older, including pregnant mares, against eastern and western encephalomyelitis (EEE & WEE), west nile virus (WNV), and tetanus, and against respiratory disease due to equine herpesvirus types 1 and 4 (EHV-1 & EHV-4) and A2 equine influenza virus (EIV). The duration of immunity against EIV OH/03 is at least 6 months and against WNV is at least 12 months. The duration for the remaining organisms has not been determined. For more information regarding efficacy and safety data go to productdata.aphis.usda.gov.

This product has also been shown to be effective against shedding due to A2 EIV and against viremia, mortality, and neurologic clinical disease due to WNV.

Composition: Contains Kentucky Lineage (KY/95), Florida sublineage clade 1 (OH/03), and Florida sublineage clade 2 (RI/07) EIV strains. Amphotericin B and gentamicin added as preservatives.

Directions and dosage: Shake well before use. Using aseptic technique, inoculate horses intramuscularly with a 1 mL dose. Administer a second 1 mL dose intramuscularly in 3 to 4 weeks using a different injection site. Historically, annual revaccination with this product has been recommended. The need for annual booster vaccination has not been established for this product. For advice on revaccination frequency, contact your veterinarian or the manufacturer.

Precautions: Store at 35-46°F (2-8°C). Do not freeze. Do not mix with other products. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine. In case of human exposure, contact a physician.

For use in animals only

Vetera® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH. Used under license.

© 2019 Boehringer Ingelheim Animal Health USA Inc. All rights reserved.

US Patent Nos. 7572620, 8535685, 8697089, 8784838, 6803041, 7226604, and 8821889.

Boehringer Ingelheim Animal Health USA Inc., St. Joseph, MO 64506

Phone: 1 (888) 637-4251

VLN/PCN 124/4855.A0

1 Dose/1 mL	137110-01
10 Doses/10 mL	137109-01

CPN: 1028268.3

THIS SERVICE AND INCLUDED DATA ARE PROVIDED "AS IS" WITHOUT ANY EXPRESS OR IMPLIED REPRESENTATION OR WARRANTY. Without limiting the foregoing, the service and data are based on third party data, and DVMetrics is not responsible or liable for such third-party data. Each user assumes full risk, responsibility, and liability related to use of the service and data. The service and data are further subject to the Terms of Use.

By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

Copyright 2023 - DVMetrics & Services LLC