

**MERCK ANIMAL HEALTH**

Intervet Inc.

**2 GIRALDA FARMS, MADISON, NJ, 07940**

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: [www.merck-animal-health-usa.com](http://www.merck-animal-health-usa.com)



THIS SERVICE AND DATA ARE PROVIDED "AS IS". DVMetrics assumes no liability, and each user assumes full risk, responsibility, and liability, related to its use of the DVMetrics service and data. See the Terms of Use for further details.

**Vision® 8 with SPUR®\***

**Intervet/Merck Animal Health**

**Cattle Vaccine**

**CLOSTRIDIUM CHAUVOEI-SEPTICUM-HAEMOLYTICUM-NOVYI-SORDELLII-PERFRINGENS TYPES C & D BACTERIN-TOXOID**

For use in healthy cattle and sheep as an aid in preventing disease caused by *Clostridium chauvoei* (Blackleg), *septicum* (Malignant edema), *haemolyticum* (Bacillary Hemoglobinuria/Red Water), *novyi* (Black disease), *sordellii* and *perfringens* Types C & D (Enterotoxemia).

**DOSE:** Cattle-2mL, Sheep-2mL. Inject subcutaneously. Repeat in 3 to 4 weeks. In animals subject to reexposure to *Cl. haemolyticum* revaccinate every 5 to 6 months. Annual revaccination is recommended. Shake well before using. Store at 2°-7°C (35°-45°F). Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur.

**ANTIDOTE:** Epinephrine. Contains formaldehyde as preservative.

\*Adjuvant-Intervet's Proprietary Technology.

**FOR ANIMAL USE ONLY**

Intervet Inc., Merck Animal Health, division of Intervet Inc., Omaha, NE 68103 USA

U.S. Vet Lic. No. 165A

1-800-521-5767

For patent information:

<http://www.merck.com/product/patent/home.html>

		Code	
10 Doses	20 mL	069136	151167-05
50 Doses	100 mL	069137	156458-05

**CPN:** 1047409.4