



# Mitochondrial Cofactors

## Cellular Energy Formula\*

The body is a complex organism that requires substantial energy for everyday functioning. The energy that drives biological activity comes from adenosine triphosphate (ATP), which contains high-energy bonds for use in biochemical reactions. Because ATP cannot be stored, it must constantly be replenished. ATP is produced in a series of enzymatic steps within the mitochondria. If this process is disturbed by nutrient deficiencies, stress, or a myriad of other conditions, fatigue can occur.\*

Mitochondrial Cofactors contains coenzyme Q10, Panmol® NADH, acteyl-L-carnitine, magnesium, and vitamin B6 as pyridoxal-5-phosphate. These bioactive cofactors have been shown to enhance energy production, with particular benefits for skeletal muscles, heart, and brain.\*

### Key Features

- Supports ATP production by activating key mitochondrial enzymes\*
- Facilitates the transport of fatty acids into the mitochondria for use as fuel\*



SKU #78270  
90 vegetarian capsules

# Mitochondrial Cofactors

**CoQ10 Ubiquinone (Ubidecarenone)** coenzyme Q10, a cofactor that has both bioenergetic and antioxidant properties.\* The main function of CoQ10 is to transfer electrons along the complexes of the mitochondrial electron transport chain, culminating in the formation of ATP.\* Low levels of CoQ10 are consistently associated with fatigue.\* In human studies, supplementation with CoQ10 increased mitochondrial efficiency in the brain.\* Healthy adults consuming CoQ10 experienced significantly less fatigue and greater exercise capacity compared to placebo groups.\*

**NADH (PANMOL®)** is a microencapsulated form of nicotinamide adenine dinucleotide (NADH), a bioactive derivative of niacin (vitamin B3).\* NADH serves an electron carrier for the ultimate production of ATP.\* An analysis of human mitochondria revealed that NADH metabolism was one of the most important biological determinants of fatigue.\* In a clinical study, individuals were asked to rate the extent to which fatigue had interfered with certain aspects of their day-to-day functioning.\*

**Acetyl-L-Carnitine** is a short-chain ester of L-carnitine that is naturally produced in the body.\* However, the biosynthesis of carnitine only accounts for 25% of daily needs.\* Carnitine plays a key role in energy metabolism, since it enables fatty acids to enter the mitochondria, where they are broken down to form ATP.\* Low blood and tissue carnitine levels often correlate with fatigue.\* Supplemental carnitine was shown to increase energy production in the brain and to reduce muscle fatigue under various conditions.\*

**Pyridoxal-5-phosphate (PLP, also known as P5P)** is a form of vitamin B6 that serves as a cofactor for numerous biosynthetic enzymes.\* PLP is needed for the synthesis of heme, the oxygen-binding molecule within hemoglobin, which supplies oxygen to every tissue in the body.\* A supply of PLP is important for normal heart and skeletal muscle contractions.\* The intake of vitamin B6 also upregulates pathways that promote the growth and repair of skeletal muscle.\* Low PLP levels are associated with decreased mitochondrial oxidative capacity and fatigue.\*

**Magnesium Bisglycinate Chelate** is a highly soluble form of magnesium that is chelated with the neutral amino acid, glycine.\* Magnesium influences the rate of ATP production by stimulating the activity of several enzymes in the trichloroacetic acid cycle (TCA, also known as the Krebs' cycle).\* Additionally, magnesium plays a pivotal role in the activity of the mitochondrial ATP synthase, the enzyme that produces the bulk of cellular ATP.\* Human and animal studies have shown that magnesium supports the normal daily functioning of the heart, brain, and skeletal muscle.\*

## References:

- An JH, et al. *Endocr J*. 2016 Oct 29;63(10):885-95.  
Andreani C, et al. *Oxid Med Cell Longev*. 2018 Oct 24;2018:8936251.  
Atamna H, et al. *Proc Natl Acad Sci USA*. 2002 Nov 12;99(23):14807-12.  
Badrasawi M, et al. *Clin Interv Aging*. 2016;11:1675.  
Barbrioli B, et al. *J Cereb Blood Flow Metab*. 1999 May;19(5):528-32.  
Bonora M, et al. *Purinergic Signal*. 2012 Sep;8(3):343-57.  
Bremer J. *Physiol Rev*. 1983 Oct 1;63(4):1420-80.  
Capaldi RA, Aggeler R. *Trends Biochem Sci*. 2002;27(3):154-60.  
Castro-Marrero J, et al. *Antioxid Redox Signal*. 2015 Mar 10;22(8):679-85.  
Castro-Marrero J, et al. *Clin Nutr*. 2016 Aug;35(4):826-34.  
Chee C, et al. *Aging Cell*. 2021 Feb;20(2):e13303.  
Chen HC, et al. *Nutrients*. 2019 Oct 23;11(10):2550.  
Chiechio S, et al. *Int J Mol Sci*. 2018 Jan;19(1):11.  
Ciacci C, et al. *Dig Liver Dis*. 2007 Oct 1;39(10):922-8.  
Ciregia F, et al. *Transl Psychiatry*. 2016 Sep 27;6(9):e904.  
Cojocaru IM, et al. *Rom J Intern Med*. 2009 Jan 1;47(2):169-71.  
Cox IM, et al. *Lancet*. 1991 Mar 30;337(8744):757-60.  
Dhalla NS, et al. *Clin Chem Lab Med*. 2013 Mar 15;51(3):535-43.  
Diaz-Castro J, et al. *Nutrients*. 2020 Feb 6;12(2):424.  
Dutta A, et al. *Exp Physiol*. 2008 Oct 19;93(10):1139-46.  
El Alaoui-Talibi ZA, et al. *Am J Physiol*. 1997 Apr 1;272(4):H1615-24.  
Fielding R, et al. *Nutrients*. 2018 Mar 13;10(3):349.  
Filler K, et al. *BBA Clin*. 2014 Jun 1;112-23.  
Fiorentini D, et al. *Nutrients*. 2021 Mar 30;13(3):1136.  
Fluge O, et al. *JCI Insight*. 2016 Dec 22;1(21):e89376.  
Forsyth LM, et al. *Ann Allergy Asthma Immunol*. 1999 Feb;82(2):185-91.  
Frey U, et al. *Front Neurosci*. 2021;15:322.  
Fukuda M, et al. *Biofactors*. 2016 Jul 8;42(4):431-40.  
Galkin MA, Syroeshkin AV. *Biochemistry*. 1999;64:1176-85.  
Garrido-Maraver J, et al. *Mol Syndromol*. 2014 Jul;5(3-4):187-97.  
Golf SW, et al. *Cardiovasc Drugs Ther*. 1998 Sep;12 Suppl 2:197-202.  
Golshani-Hebroni S, et al. *Gene*. 2016 Apr 25;581(1):1-13.  
He L, et al. *Ind Health*. 2008;46(5):506-12.  
Hiatt WR, et al. *J Clin Invest*. 1989;84:1167-73.  
Kalicki B, et al. *Cent Eur J Immunol*. 2019;44(1):23-32.  
Kepka A, et al. *Postepy Hig Med Dosw (Online)*. 2016 Jan 1;70.  
Kumrunsee T, et al. *Eur J Nutr*. 2021 Aug 26. PMID: 34436643.  
Heap AC, et al. *J R Soc Med*. 1999 Apr;92(4):183-5.  
Hernández-Camacho JD, et al. *Front Physiol*. 2018 Feb 5;9:44.  
Horowitz R, Freeman PR. *Healthcare (Basel)*. 2018 Nov 5;6(4):129.  
Huang A, Owen K. *Med Sport Sci*. 2012;59:135-42.  
Hunter GA, Ferreira GC. 2009 Feb 16;55(1):102-10.  
Johnson S. *Med Hypotheses*. 2001 Feb 1;56(2):163-70.  
Kennedy DO, et al. *Nutr Metab (Lond)*. 2016 Dec;13(1):1-6.  
Kerner J, Hoppel C. *Biochim Biophys Acta*. 2000;1486:1-17.  
Ko YH, et al. *J Biol Chem*. 1999 Oct 8;274(41):28853-6.  
Kolte D, et al. *Cardiol Rev*. 2014 Jul 1;22(4):182-92.  
Kumar A, et al. *Sci Rep*. 2020 Jul 16;10(1):11824.  
Kuratsune H, et al. *Neuroimage*. 2002 Nov 1;17(3):1256-65.  
Kwon OS, Chung YB. *Arch Pharm Res*. 2004 Jun;27(6):676-81.  
Langsjoen PH, Langsjoen AM. *Clin Pharmacol Drug Dev*. 2014 Jan;3(1):13-7.  
Littaru GP, Tiano L. *Nutrition*. 2010;26(3):250-4.  
Lohninger A, et al. *Monatsh. Chem*. 2005 Aug;136(8):1425-42.  
Madsen KL, et al. *J Clin Endocrinol Metab*. 2018 Dec 1;103(12):4580-8.  
Malaguerneira M, et al. *Arch Gerontol Geriatr*. 2008 Mar 1;46(2):181-90.  
Malaguerneira M, et al. *J Interferon Cytokine Res*. 2011 Sep;31(9):653-9.  
Marcovina SM, et al. *Transl Res*. 2013;161(2):73-84.  
Mendelsohn BA, et al. *PLoS Biol*. 2018 Aug 27;16(8):e2004624.  
Modanloo M, Shokrzadeh M. *Iran J Kidney Dis*. 2019 Mar;13(2):74-86.  
Montesano A, et al. *Oxid Med Cell Longev*. 2015 Mar 8;2015.  
Nicolson GL, et al. *Funct Foods Health Dis*. 2012 Mar 28;2(3):35-47.  
Nicolson GL. *Altern Ther Health Med*. 2014 Aug;13(4):35.  
Odai T, et al. *Nutrients*. 2020 Nov 9;12(11):3437.  
Orsucci D, et al. *Curr Med Chem*. 2011;18(26):4053-64.  
Pilchova I, et al. *Oxid Med Cell Longev*. 2017;2017:6797460.  
Pizzorno J. *Integr Med (Encinitas)*. 2014 Apr;13(2):8-15.  
Plioplys AV, Plioplys S. *Neuropsychobiology*. 1995;32(3):132-8.  
Poljsak B, et al. *Oxid Med Cell Longev*. 2020 Dec 12;2020:8819627.  
Rebouche CJ. *FASEB J*. 1992;6:3379-86.  
Rebouche CJ. *Ann N Y Acad Sci*. 2004 Nov;1033(1):30-41.  
Reinhardt RA. *Arch Intern Med*. 1988 Nov;148(11):2415-20.  
Reuter SE, Evans AM. *Clin Pharmacokinet*. 2012;51(9):553-72.  
Ringseis R, et al. *Eur J Nutr*. 2013 Aug;52(5):1421-42.  
Rosenfeldt F, et al. *Biofactors*. 2003;18(1-4):91-100.  
Santaella ML, et al. *P R Health Sci J*. 2004;23(2).  
Santos DA, et al. *Magnes Res*. 2011 Dec;24(4):215-9.  
Sarmiento et al. *Curr Drug Metab*. 2016;17(4):345-58.  
Schuette SA, et al. *JPEN J Parenter Enteral Nutr*. Sep-Oct 1994;18(5):430-5.  
Smeland OB, et al. *Neurochem Int*. 2012 Jul 1;61(1):100-7.  
Stein LR, Imai SI. *Trends Endocrinol Metab*. 2012 Sep;23(9):420-8.  
Suidasari S, et al. *Front Nutr*. 2016 Jan 19;2:39.  
Syroeshkin AV, et al. *Biochemistry (Mosc)*. 1999 Oct;64(10):1128-37.  
Tanabe K, et al. *Jpn Circ J*. 1998 May;62(5):341-6.  
Tardy AL, et al. *Nutrients*. 2020 Jan;12(1):228.  
Tomas C, et al. *PLoS One*. 2017 Oct 24;12(10):e0186802.  
Uberti F, et al. *Nutrients*. 2020 Feb 22;12(2):573.  
Uziel G, et al. *Muscle Nerve*. 1988 Jul;11(7):720-4.  
van Dijk M, et al. *J Cachexia Sarcopenia Muscle*. 2018 Feb;9(1):146-59.  
Varanoske AN, et al. *Nutrients*. 2017 Sep 7;9(9):988.  
Vermeulen RC, Scholte HR. *Psychosom Med*. 2004 Mar 1;66(2):276-82.  
Veronesi N, et al. *Am J Clin Nutr*. 2014 Sep 1;100(3):974-81.  
Wall BT, et al. *J Physiol*. 2011;589:963-73.  
Yamanaka R, et al. *Sci Rep*. 2016 Jul 26;6(1):1-2.  
Zhang Y, et al. *Nutrients*. 2017 Sep;9(9):946.

## Supplement Facts

Serving Size	3 Capsules	
Servings Per Container	30	
Amount Per Serving	% Daily Value**	
Vitamin B6 (as Pyridoxal-5-Phosphate)	25 mg	1470%
Magnesium (as Magnesium Bisglycinate)	100 mg	23%
Acetyl-L-Carnitine	300 mg	†
Coenzyme Q10 (Ubiquinone)	100 mg	†
NADH (PANMOL® NADH) (Reduced B-Nicotinamide Adenine Dinucleotide)	5 mg	†
† Daily value not established. **Percent Daily Value are based on a 2,000 calorie diet.		

Other ingredients: Hydroxypropyl methylcellulose, microcrystalline cellulose, silicon dioxide, calcium palmitate, stearic acid.

**Suggested Use:** As a dietary supplement, 3 capsules, one or two times daily with meals, or as directed by a healthcare practitioner.

PANMOL®  
NADH