

PMRA Approved Label

2022-1680

2023-03-30

LALGUARD M52 OD

Bioinsecticide

For Use on Outdoor Turf, Greenhouse Crops and Cannabis (Marihuana) Produced
Commercially Indoors

COMMERCIAL
EMULSIFIABLE CONCENTRATE
READ THE LABEL BEFORE USING
SHAKE VIGOROUSLY BEFORE USING

ACTIVE INGREDIENT: *Metarhizium brunneum* Strain F52
Contains a minimum of 2×10^9 CFU/mL

REGISTRATION NO: 30829 PEST CONTROL PRODUCTS ACT

WARNING - EYE IRRITANT
POTENTIAL SENSITIZER

Net Contents _____

Danstar Ferment AG
Poststrasse 30
Zug, Switzerland CH-6300

Canadian agent:
Lallemand Inc./LALLEMAND PLANT CARE
59 Industrial Park Crescent, Unit 1
Sault Ste. Marie, ON P6B 5P3
Phone: 1-888-236-7378

Batch Code/Expiry Date: XXXX/dd-mm-year

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May cause sensitization. Keep away from food, drink and animal feed.

Causes eye irritation. DO NOT get in eyes. Avoid contact with skin or clothing. Avoid breathing mists.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes and socks, eye goggles, and a dust-mist filtering respirator/mask (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator/mask with any N-95, R-95, P-95 or HE filter. Wash thoroughly with soap and water after handling. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Do not re-enter or allow re-entry into treated areas until the spray is dried unless wearing appropriate personal protective equipment, including a long-sleeved shirt, long pants, shoes plus socks, and waterproof gloves. In addition, a dust-mist filtering respirator/mask (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator/mask with any N-95, R-95, P-95 or HE filter is required until spray mist has settled.

Can be applied up to and including the day of harvest. The pre-harvest interval is 0 days.

DO NOT APPLY BY AIR.

ENVIRONMENTAL PRECAUTIONS: As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters. For guidance contact the Provincial Regulatory Agency.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

FIRST AID:

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

DIRECTIONS FOR USE

GENERAL: LALGUARD M52 OD is composed of spores of the insect pathogenic fungus *Metarhizium brunneum* strain F52. Insects that come into contact with the fungus will become infected. The spores of *Metarhizium brunneum* attach to the surface of the insect, germinate and begin to grow. It will penetrate the exoskeleton of the insect and grow inside the insect, causing the insect to die. Under moderate temperatures it usually takes 3 to 7 days, once the insect is exposed for death to occur. At very cold or hot temperatures death may occur more slowly.

Note: LALGUARD M52 OD contains live spores that may be harmed by storage at high temperatures or held as a water suspension for more than 24 hours. See storage instructions on this label.

MODE OF ACTION AND APPLICATION TIMING: Application frequency will depend on the environment to which LALGUARD M52 OD is applied and the target insects' life cycle.

LALGUARD M52 OD may be used for foliar and drench applications against labelled pests. It is best to begin applying LALGUARD M52 OD at early stages of population development.

Applications of drenches to growing media may be done as appropriate to standard pest management practices. When possible, application to growing media should be done in advance of a target insect's life stage that will be passing through the first 15 cm of growing media. Although the penetration of LALGUARD M52 OD into growing media will vary greatly with growing media type and level of irrigation it is best to target populations as close to the surface as possible or time application of LALGUARD M52 OD with a growing media incorporation event.

APPLICATION INSTRUCTIONS

Re-entry interval: 0 hours

Mixing instructions: SHAKE VIGOROUSLY BEFORE USING. Apply LALGUARD M52 OD with conventional ground application equipment. Do not mix LALGUARD M52 OD with fungicides. LALGUARD M52 OD contains emulsifiers and mixes readily in water. Fill spray tank with half of the desired amount of water and start agitation. Slowly add desired amount of LALGUARD M52 OD to spray tank. Add remaining desired amount of water. Triple rinse empty

LALGUARD M52 OD container with water and add rinse water to spray tank. Continue agitation throughout spraying. Do not mix more LALGUARD M52 OD than can be applied within six hours of mixing. Do not apply fungicides near the time of application of LALGUARD M52 OD.

Plant safety: LALGUARD M52 OD has not been tested on all varieties of listed plant species. Test LALGUARD M52 OD on a small number of plants to check for potential damage before applying to a large number of plants.

Foliar application, Greenhouse use - Apply at a rate of 0.5-5.0 L per 1000 L. Use the higher application concentration when pest pressure is high. Re-apply as required. The need for and timing of re-application should be determined by monitoring. An application interval of 5-10 days is recommended. Mix well by external mixing, in-tank mixing, or pump circulation to form emulsion. Follow spray equipment manufacturer's instructions for final spray volume to obtain adequate coverage. Do not apply at pressures above 1380 KPa (200 PSI). **Spray to wet all foliage, but avoid runoff. Do not apply through a thermal pulse fogger.**

CROP	PEST	APPLICATION	CONCENTRATION
Zucchini	Reduces numbers of thrips	Foliar	0.5 – 5.0 L/1000 L
Fruiting vegetables (tomato, pepper, eggplant), Cucumber, Lettuce	Reduces number of mites, thrips and whiteflies	Foliar	0.5 – 5.0 L/1000 L
Strawberry	Reduces number of spider mites, thrips and whiteflies	Foliar	0.5 – 5.0 L/1000 L
Ornamentals excluding conifers	Reduces number of spider mites, thrips, whiteflies and aphids	Foliar	0.5 – 5.0 L/1000 L

Foliar application, Cannabis (Marihuana) Produced Commercially Indoors - Apply at a rate of 0.5-5.0 L per 1000 L. Use the higher application concentration when pest pressure is high. Re-apply as required. The need for and timing of re-application should be determined by monitoring. An application interval of 5-10 days is recommended. Mix well by external mixing, in-tank mixing, or pump circulation to form emulsion. Follow spray equipment manufacturer's instructions for final spray volume to obtain adequate coverage. Do not apply at pressures above 1380 KPa (200 PSI). **Spray to wet all foliage, but avoid runoff. Do not apply through a thermal pulse fogger.**

Restricted Entry Interval: DO NOT enter or allow worker entry into treated areas for 4 hours or until sprays have dried.

PEST	APPLICATION	CONCENTRATION
Reduces number of mites, thrips and whiteflies	Foliar	0.5 – 5.0 L/1000 L

Soil drench application Greenhouse use – Add 108 mL of LALGUARD M52 OD to 10 L of drench solution. Drench application should be thoroughly watered-in without causing water to come out of the bottom of the pots/grow bags – depending on the growing media type and moisture this will be around 250 ml per 4 L pot or grow bag. Re-apply as required. The need for and timing of re-application should be determined by monitoring. **Do not apply via drip irrigation.**

CROP	PEST	APPLICATION	CONCENTRATION
Fruiting vegetables (tomato, pepper, eggplant), Cucumber, Lettuce	May reduce number of thrips	Drench	108 mL/10 L

Soil drench application, Cannabis (Marihuana) Produced Commercially Indoors – Add 108 mL of LALGUARD M52 OD to 10 L of drench solution. Drench application should be thoroughly watered-in without causing water to come out of the bottom of the pots/grow bags – depending on the growing media type and moisture this will be around 250 ml per 4 L pot or grow bag. Re-apply as required. The need for and timing of re-application should be determined by monitoring. **Do not apply via drip irrigation.**

PEST	APPLICATION	CONCENTRATION
May reduce number of thrips	Drench	108 mL/10 L

Foliar application, Turf use - Apply at a rate of 64-96 mL/100 m² against hairy chinch bugs and 137-155 mL/100 m² against ticks. Use the higher application rate when pest pressure is high. Re-apply as required. The need for and timing of re-application should be determined by monitoring. An application interval of 4 weeks for hairy chinch bug or 3 weeks for ticks is recommended. Mix well by external mixing, in-tank mixing, or pump circulation to form emulsion. Follow spray equipment manufacturer's instructions for final spray volume to obtain adequate coverage. Do not apply at pressures above 1380 KPa (200 PSI). **Spray to wet all foliage, but avoid runoff. Do not apply through a thermal pulse fogger.**

CROP	PEST	APPLICATION	RATE
Turf	Hairy chinch bug (suppression)	foliar	64 - 96 mL/100 m ²
	Ticks (suppression)	foliar	137 – 155 mL/100 m ²

SITES FOR USE:

This product is for outdoor turf and greenhouse agricultural use.

STORAGE: Keep in original container during storage. To prevent contamination, store this product away from food or feed. Keep away from direct sunlight, fire or open flame, or other source of heat.

LALGUARD M52 OD consists of living microbes. Store closed product in a cool dry place at 20°C. Use product by expiry date noted on the label. Once open, use entire contents of bottle. Exposing product to temperatures in excess of the recommended storage temperature will reduce the shelf-life. Do not freeze.

DISPOSAL:

1. Triple- or pressure rinse the empty container. Add the rinsings to the treatment site.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.

4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.