



37878 PS

USN PHEDRACUT LIPO X BLACK LIQUID

(liquid)

SCHEDULING STATUS: S0**COMPLEMENTARY MEDICINE: COMBINATION PRODUCT (WESTERN HERBAL MEDICINE / HEALTH SUPPLEMENT)**

This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use.

1. NAME OF THE MEDICINE

USN PHEDRACUT LIPO X BLACK LIQUID

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 25 ml contains:	*	%NRV
Ascorbic acid (Vitamin C)	500 mg	500 %
L-Carnitine L-Tartrate	500 mg	
N-Acetyl L-Tyrosine	200 mg	
Caffeine Anhydrous	200 mg	
Fucus vesiculosus L. (Kelp) [whole herb; extract providing 10 % fucoxanthin]	120 mg	
Raspberry Ketone [fruit; extract providing 98 % p-hydroxyphenyl butanone]	100 mg	
Thermozine™ Aframomum melegueta K. Schum (Grains of Paradise) [seed; 10:1 extract providing 400 mg of dried herb equivalent]	40 mg	
Nicotinic acid (Vitamin B3)	17.5 mg	109 %
Pyridoxine HCl providing Pyridoxine (Vitamin B6)	0.33 mg	
Chromium Picolinate providing Chromium (elemental)	0.27 mg	16 %
	402 µg	
	50 µg	143 %

*%Nutrient Reference Value (NRVs) for individuals 4 years and older (2010)

CONTAINS CAFFEINE (200 mg per 25 ml).

A cup of instant coffee contains approximately 80 mg of caffeine.

Excipients with known effects:

Contains sugar (1 000 mg Fructose per 25 ml) and sugar alcohol (2 500 mg Sorbitol per 25 ml).

Contains sweetener (104.5 mg Sucralose per 25 ml).

Preservatives:

Potassium sorbate 0.08 % m/v and Sodium benzoate 0.15 % m/v.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Liquid.

Black coloured liquid with a fireball cinnamon flavour.

4. CLINICAL PARTICULARS**4.1 Therapeutic indications**

USN PHEDRACUT LIPO X BLACK LIQUID is indicated as a thermogenic aid to assist weight loss as part of a weight management programme. Only effective when combined with a balanced, energy-restricted eating plan and regular physical activity.

4.2 Posology and method of administration**Adults:**

- Take 25 ml (2 tablespoons) 2 times daily.
- For best results, take 1 dosage (25 ml) on an empty stomach, 20 minutes prior to morning workout/breakfast.
- Take another dosage 20 minutes before lunch.
- Do not use USN PHEDRACUT LIPO X BLACK LIQUID for more than five consecutive days per week. May be alternated with PHEDRACUT capsules in the same day.
- Shake well before use.
- Do not exceed the recommended daily dosage.

Children:

Not suitable for children under the age of 18 years (see section 4.4).

4.3 Contraindications

Hypersensitivity to any of the active ingredients or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use**Bleeding disorders:**

USN PHEDRACUT LIPO X BLACK LIQUID may have antiplatelet effects and may increase the risk of bruising and bleeding when used in patients with bleeding disorders. Patients should be advised to discontinue USN PHEDRACUT LIPO X BLACK LIQUID at least 2 weeks prior to surgical procedures (see section 4.5).

Cardiac conditions:

USN PHEDRACUT LIPO X BLACK LIQUID may reduce blood flow to the heart muscle during exercise which might lead to cardiovascular complications such as chest pain and dysrhythmia, even in healthy individuals. USN PHEDRACUT LIPO X BLACK LIQUID may also temporarily increase blood pressure when used short-term.

Diabetes mellitus:

USN PHEDRACUT LIPO X BLACK LIQUID may increase insulin levels and/or decrease blood glucose levels and cause hypoglycaemia. Dose adjustment of antidiabetic medicine might be necessary (see section 4.5).

Glaucoma:

USN PHEDRACUT LIPO X BLACK LIQUID may temporarily increase intraocular pressure in patients with narrow-angle glaucoma.

Renal disease:

USN PHEDRACUT LIPO X BLACK LIQUID may exacerbate renal dysfunction in patients with renal disease. USN PHEDRACUT LIPO X BLACK LIQUID may also increase the risk of incontinence or worsen symptoms of urinary incontinence, such as urinary frequency and urgency (overactive bladder syndrome).

Liver disease:

USN PHEDRACUT LIPO X BLACK LIQUID may exacerbate liver dysfunction in patients with liver disease.

Hypothyroidism:

USN PHEDRACUT LIPO X BLACK LIQUID may inhibit the activity of thyroid hormone in target tissues.

Epilepsy:

USN PHEDRACUT LIPO X BLACK LIQUID may increase seizure frequency or severity in people with a history of seizures.

Sleep deprivation:

Use of USN PHEDRACUT LIPO X BLACK LIQUID is not recommended before bedtime as it may result in sleep deprivation. USN PHEDRACUT LIPO X BLACK LIQUID is not intended as a substitute for sleep.

Patients following a low protein diet should consult a healthcare provider before they take USN PHEDRACUT LIPO X BLACK LIQUID.

Contains sugar:

USN PHEDRACUT LIPO X BLACK LIQUID contains sugar (fructose) and sugar alcohol (sorbitol). The additive effect of concomitantly administered products containing fructose or sorbitol and dietary intake of fructose or sorbitol should be taken into account.

Patients with hereditary fructose intolerance (HFI) should not take USN PHEDRACUT LIPO X BLACK LIQUID.

The content of sorbitol in USN PHEDRACUT LIPO X BLACK LIQUID may affect the bioavailability of other medicinal products for oral use administered concomitantly.

USN PHEDRACUT LIPO X BLACK LIQUID also contains maltodextrin. Maltodextrin has a high glycemic index value and should be used sparingly by diabetic and/or hypoglycemic individuals, as it may increase blood sugar levels.

Paediatric population:

The safety and efficacy of USN PHEDRACUT LIPO X BLACK LIQUID in children under the age of 18 years have not been established. USN PHEDRACUT LIPO X BLACK LIQUID should not be used in children under 18 years (see section 4.2).

4.5 Interaction with other medicines and other forms of interaction**Anticoagulant/antiplatelet medicines:**

USN PHEDRACUT LIPO X BLACK LIQUID may enhance the effects of anticoagulant/antiplatelet medicines or herbal supplements. Concomitant use may increase the risk of bruising and bleeding.

Antidiabetic medicines:

The use of USN PHEDRACUT LIPO X BLACK LIQUID with antidiabetic medicines or herbal supplements with hypoglycaemic effects may have an additive effect when used concomitantly (see section 4.4). Caution is advised.

Ephedrine and other stimulants:

Concomitant use of USN PHEDRACUT LIPO X BLACK LIQUID with ephedrine or other stimulants such as bitter orange extract, synephrine, octopamine and ephedra may increase blood pressure and heart rate and potentially increase the risk of serious cardiovascular adverse effects. Caution is advised.

Caffeine-containing products or foods:

The use of caffeine-containing products should be limited when taking USN PHEDRACUT LIPO X BLACK LIQUID. Use with other caffeine-containing products or foods may increase the risk of caffeine-related adverse effects. Other natural products which contain caffeine include coffee, tea, cola nut, cocoa, guarana and maté.

Monoamine oxidase inhibitors (MAOIs):

Concomitant use of USN PHEDRACUT LIPO X BLACK LIQUID with MAOIs may increase blood pressure raising effects. Caution is advised.

Levothyroxine:

The use of USN PHEDRACUT LIPO X BLACK LIQUID with levothyroxine may decrease serum levels of levothyroxine. Levothyroxine should be taken at least 30 minutes before or 3 – 4 hours after taking USN PHEDRACUT LIPO X BLACK LIQUID.

Lithium:

The concomitant use of USN PHEDRACUT LIPO X BLACK LIQUID with lithium may affect serum lithium levels. Caution is advised.

4.6 Fertility, pregnancy and lactation:

Safety in pregnancy and lactation has not been established. Total caffeine intake more than 200 mg per day is not recommended during pregnancy or breastfeeding.

4.7 Effects on ability to drive and use machines:

USN PHEDRACUT LIPO X BLACK LIQUID may cause side effects such as dizziness which can affect the ability to drive a vehicle and use machines (see section 4.8).

4.8 Undesirable effects

USN PHEDRACUT LIPO X BLACK LIQUID is generally well tolerated.

Immune system disorders:

Frequency unknown: hypersensitivity and/or allergic reactions.

Psychiatric disorders:

Frequency unknown: anxiety, nervousness, insomnia.

Nervous system disorders:

Frequency unknown: restlessness, dizziness, tremors, agitation, headache.

Cardiac disorders:

Frequency unknown: dysrhythmia, hypertension.

Vascular disorders:

Frequency unknown: flushing.

Gastrointestinal disorders:

Frequency unknown: gastric irritation, nausea, vomiting, abdominal cramps, dyspepsia, diarrhoea.

Renal and urinary disorders:

Frequency unknown: diuresis.

Reporting of suspected adverse reactionsReporting suspected adverse reactions after authorisation of USN PHEDRACUT LIPO X BLACK LIQUID is important. It allows continued monitoring of the benefit/risk balance of USN PHEDRACUT LIPO X BLACK LIQUID. Healthcare providers are asked to report any suspected adverse reactions to the South African Health Products Regulatory Authority (SAHPRA) via the 6.04 Adverse Drug Reaction Reporting Form, found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>.**4.9 Overdose**

In overdose, side effects can be precipitated and/or be of increased severity (see section 4.8).

In the event of overdose, treatment should be symptomatic and supportive.

5. PHARMACOLOGICAL PROPERTIES**5.1 Pharmacodynamic properties****Category and class:**

D 33.7 Combination Product.

USN PHEDRACUT LIPO X BLACK LIQUID contains a variety of herbal ingredients, vitamins, minerals, amino acids and caffeine to assist with weight loss as part of a weight management programme.

5.2 Pharmacokinetic properties

Ascorbic acid (Vitamin C) is readily absorbed from the gastrointestinal tract and is widely distributed in the body. The main route of elimination is through urine.

L-Carnitine (from L-Carnitine L-Tartrate) is mainly distributed to the muscle. The half-life of L-Carnitine is about 15 hours and is excreted in the urine.

Tyrosine (from N-Acetyl L-Tyrosine) is an amino acid that is converted in the brain by tyrosine hydroxylase to L-dopa, dopamine, norepinephrine and then epinephrine, and are excreted in the urine.

Caffeine is rapidly and almost completely (99 %) absorbed after oral administration. Cytochrome P450 1A2 plays a role in the metabolism of caffeine in the liver. The half-life of caffeine in healthy adults is 3 – 6 hours and is excreted in the urine. Pregnancy and oral contraceptives appear to delay caffeine clearance and smoking increases it.

Raspberry ketone and metabolites (conjugated with glucuronic acid or sulfate) are excreted in the urine within 24 hours.

Nicotinic acid (Vitamin B3) is rapidly absorbed from the gastrointestinal tract and is excreted mainly via urine.

Pyridoxine (Vitamin B6) is passively absorbed from the upper gastrointestinal tract, converted in the liver to coenzyme pyridoxal phosphate and excreted in the urine.

Approximately 0.5 % to 2 % of chromium is absorbed. Chromium as chromium picolinate have increased absorption, retention and accumulation of chromium. Absorbed chromium is rapidly excreted in the urine and unabsorbed chromium in the faeces.

There is insufficient information available about the pharmacokinetics of *Fucus vesiculosus* L. (Kelp) and *Aframomum melegueta* K. Schum (Grains of Paradise).**6. PHARMACEUTICAL PARTICULARS****6.1 List of excipients**

Carbon black (colourant)

Citric acid anhydrous (E330)

Fructose

Glycerine

Hot cinnamon flavour

Maltodextrin

Natural flavours

Potassium sorbate (E202)

Purified water

Silicon dioxide (E551)

Sodium benzoate (E211)

Sorbitol (E420)

Sucralose.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

14 months.

Store at or below 25 °C.

Use within 1 month after first opening.

6.4 Special precautions for storage

Refrigerate after opening.

6.5 Nature and contents of container



USN PHEDRACUT LIPO X BLACK LIQUID

37878 PS

SCHEDULING STATUS: **S0**

USN PHEDRACUT LIPO X BLACK LIQUID

Contains sugar (1 000 mg Fructose per 25 ml) and sugar alcohol (2 500 mg Sorbitol per 25 ml).

Contains sweetener (104,5 mg Sucralose per 25 ml).

COMPLEMENTARY MEDICINE: COMBINATION PRODUCT (WESTERN HERBAL MEDICINE / HEALTH SUPPLEMENT)

This unregistered medicine has not been evaluated by SAHPRA for its quality, safety or intended use.

Read all of this leaflet carefully because it contains important information for you.

USN PHEDRACUT LIPO X BLACK LIQUID is available without a doctor's prescription, for you to maintain your health.

Nevertheless, you still need to use USN PHEDRACUT LIPO X BLACK LIQUID carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share USN PHEDRACUT LIPO X BLACK LIQUID with any other person.
- Ask your healthcare provider or pharmacist if you need more information or advice.

What is this leaflet?

1. What USN PHEDRACUT LIPO X BLACK LIQUID is and what it is used for.
2. What you need to know before you take USN PHEDRACUT LIPO X BLACK LIQUID.
3. How to take USN PHEDRACUT LIPO X BLACK LIQUID.
4. Possible side effects.
5. How to store USN PHEDRACUT LIPO X BLACK LIQUID.
6. Contents of the pack and other information.

1. What USN PHEDRACUT LIPO X BLACK LIQUID is and what it is used for

USN PHEDRACUT LIPO X BLACK LIQUID is indicated as a thermogenic aid to assist you with weight loss as part of a weight management programme. Only effective when combined with a balanced, energy-restricted eating plan and regular physical activity.

2. What you need to know before you take USN PHEDRACUT LIPO X BLACK LIQUID

Do not take USN PHEDRACUT LIPO X BLACK LIQUID:

If you are hypersensitive (allergic) to any of the ingredients (listed in section 6).

Warnings and precautions

Take special care with USN PHEDRACUT LIPO X BLACK LIQUID:

- CONTAINS CAFFEINE. Each 25 ml contains 200 mg caffeine. A cup of instant coffee contains approximately 80 mg of caffeine.
- If you have a bleeding disorder or must go for surgery, USN PHEDRACUT LIPO X BLACK LIQUID should be discontinued at least 2 weeks before any surgical procedures.
- If you have heart problems or high blood pressure.
- If you have diabetes mellitus (high blood sugar) and you are taking medicines to control your blood sugar levels.
- If you have glaucoma (a condition causing increased pressure behind your eye).
- If you have any kidney problems or overactive bladder syndrome.
- If you have a liver disease.
- If you have an underactive thyroid.
- If you have epilepsy/seizure disorder.
- If you are following a low protein diet.
- Do not take USN PHEDRACUT LIPO X BLACK LIQUID before bedtime, as it may result in sleep deprivation.

Children and adolescents

Do not give USN PHEDRACUT LIPO X BLACK LIQUID to children younger than 18 years, as the safety and efficacy in children have not yet been established.

Other medicines and USN PHEDRACUT LIPO X BLACK LIQUID

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor or pharmacist if you are currently taking:

- Anticoagulant/antiplatelet medicines (used to thin your blood), or herbal supplements with blood thinning effects.
- Antidiabetic medicines (used to treat high blood sugar levels), or herbal supplements which can influence blood sugar levels.
- Ephedrine and other stimulants, such as bitter orange extract, synephrine, octopamine and ephedra.
- Other caffeine-containing products.
- Monoamine oxidase inhibitors (MAOIs) (used to treat depression).
- Levotyroxine (used to treat an underactive thyroid).
- Lithium (used to stabilise your mood).

PASIËNTINLIGTINGSBLAD



USN PHEDRACUT LIPO X BLACK LIQUID

(vloeistof)

SKEDULERINGSTATUS: **S0**

USN PHEDRACUT LIPO X BLACK LIQUID

Bevat suiker (1 000 mg Fructose per 25 ml) en suikeralkohol (2 500 mg Sorbitol per 25 ml).

Bevat versroter (104,5 mg Sukralose per 25 ml).

KOMPLEMENTÈRE MEDISYNE: KOMBINASIEPRODUK (WESTERSE KRUIEMEDISYNE / GESONDHEIDAANVULMING)

Hierdie ongeregistreerde medisyne is nie vir gehalte, veiligheid of beoogde gebruik deur SAHPRA geëvalueer nie.

Lees hierdie hele inligtingsblad aandagdig deur want dit bevat inligting wat belangrik is vir jou.

USN PHEDRACUT LIPO X BLACK LIQUID is beskikbaar sonder 'n dokter se voorskrif, sodat jy jou gesondheid in stand kan hou. Jy moet USN PHEDRACUT LIPO X BLACK LIQUID nochtans met sorg neem om die beste resultate te verkry.

- Hou hierdie inligtingsblad. Dit mag nodig wees dat jy dit weer lees.
- Moenie USN PHEDRACUT LIPO X BLACK LIQUID met enigemand anders deel nie.
- Vra gerus jou apteker indien jy verdere inligting of advies nodig het.

Wat is in hierdie inligtingsblad

1. Wat is USN PHEDRACUT LIPO X BLACK LIQUID en waaroor word dit gebruik.
2. Wat jy moet weet vooroor dat jy USN PHEDRACUT LIPO X BLACK LIQUID neem.
3. Hoe om USN PHEDRACUT LIPO X BLACK LIQUID te neem.
4. Montlike newe-effekte.
5. Hoe om USN PHEDRACUT LIPO X BLACK LIQUID te bêre.
6. Inhou van die pak en ander inligting.

1. Wat is USN PHEDRACUT LIPO X BLACK LIQUID en waaroor word dit gebruik

USN PHEDRACUT LIPO X BLACK LIQUID is aangedui as 'n termogeniese hulpmiddel om gewigverlies as deel van 'n verlankingsprogram, te ondersteun. Slegs effekief indien met 'n gebalanseerde, energie-beperkte eetplan en gereeld oefening gekombineer word.

2. Wat jy moet weet vooroor dat jy USN PHEDRACUT LIPO X BLACK LIQUID neem

Moenie USN PHEDRACUT LIPO X BLACK LIQUID neem nie

Indien jy hypersensitief (allergies) is vir enige van die ander bestanddele (geleys by afdeling 6).

Waarskuwings en voorsorgmaatreëls

Wees vereit versigtig met USN PHEDRACUT LIPO X BLACK LIQUID:

- BEVAT KAFEÏEN. Elké 25 ml bevat 200 mg kafeïen. 'n Kopjie koffie bevat ongeveer 80 mg kafeïen.
- Indien jy 'n bloedingsversteuring het of vir 'n operasie moet gaan. Hou twee weke voor enige chirurgiese procedure (operasie) op om USN PHEDRACUT LIPO X BLACK LIQUID te neem.
- Indien jy hartprobleme of hoë bloeddruk het.
- Indien jy diabetes mellitus (hoë bloedsuiker) het en jy medisyne gebruik om jou bloedsuikervlakke te beheer.
- Indien jy glokom het ('n toestand wat verhoogde druk in jou oog veroorsaak).
- Indien jy enige nierprobleme of ooraktiewe blaasindroom het.
- Indien jy 'n leverziekte het.
- Indien jy 'n onderaktiewe skildklier het.
- Indien jy epilepsie/'n aanvalversteuring het.
- Indien jy 'n lae protein-dieet volg.
- Moenie USN PHEDRACUT LIPO X BLACK LIQUID voor slaaptyd neem nie, want dit kan slaaptekort tot gevolg hê.

Kinders en tiener

Moenie nie USN PHEDRACUT LIPO X BLACK LIQUID aan kinders jonger as 18 jaar gee nie, aangesien die veiligheid en doeltreffendheid by kinders nog nie vastgestel is nie.

Ander medisyne en USN PHEDRACUT LIPO X BLACK LIQUID

Sé altyd vir jou gesondheidsorgverskaffer indien jy enige ander medisyne neem. (Dit sluit komplementêre of tradisionele medisyne in.)

Sé vir jou dokter of apteker indien jy tans die volgende gebruik:

- Antikoagulantmedisyne (word gebruik om jou bloed dun te hou) of kruie-aanvullings met bloedverdunnende effekte.
- Antidiabetiese medisyne (gebruik om hoë bloedsuikervlakke te behandel) of kruie-aanvullings met bloedsuikerverlagende effekte.
- Efedrien of ander stimulante, soos bitterlemoen-ekstrak, synephrine, oktopamien en efedra.
- Ander kafeïen-bevattende produkte.
- Monoaminoëksidase inhibeerders (MAOIs) (gebruik om depressie te behandel).
- Levotyroxine (gebruik vir die behandeling van 'n onderaktiewe skildklier).
- Litium (gebruik om jou gemoed te stabiliseer).

USN PHEDRACUT LIPO X BLACK LIQUID with food and drink

Do not take USN PHEDRACUT LIPO X BLACK LIQUID with other caffeine-containing food and drinks, such coffee, tea, cola nut, cocoa, guarana and maté.

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking USN PHEDRACUT LIPO X BLACK LIQUID. Safety in pregnancy and lactation has not been established. If you are of childbearing age, pregnant or breastfeeding and have concerns that your daily intake of caffeine from all sources may exceed 200 mg per day, please consult a healthcare provider prior to use.

Driving and using machines

USN PHEDRACUT LIPO X BLACK LIQUID may cause side effects, such as dizziness and can affect your ability to drive a vehicle and use machines (see section 4). Do not drive or operate machinery until you know how USN PHEDRACUT LIPO X BLACK LIQUID affects you.

USN PHEDRACUT LIPO X BLACK LIQUID contains sugar

USN PHEDRACUT LIPO X BLACK LIQUID contains 1 000 mg fructose and 2 500 mg sorbitol per 25 ml. If your doctor has told you that you have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to your doctor before you take USN PHEDRACUT LIPO X BLACK LIQUID.

USN PHEDRACUT LIPO X BLACK LIQUID also contains maltodextrin. This should be taken into account if you have diabetes mellitus, as maltodextrin can increase blood sugar levels.

3. How to take USN PHEDRACUT LIPO X BLACK LIQUID

Always take USN PHEDRACUT LIPO X BLACK LIQUID exactly as described in this leaflet or as your pharmacist has told you. Check with your pharmacist if you are not sure.

Adults:

- The usual dose is 25 ml (2 tablespoons) 2 times daily.
- For best results, take 1 dosage (25 ml) on an empty stomach, 20 minutes prior to morning workout/breakfast.
- Take another dosage 20 minutes before lunch.
- Do not use USN PHEDRACUT LIPO X BLACK LIQUID for more than five consecutive days per week. May be alternated with PHEDRACUT capsules in the same day.
- Shake well before use.
- Do not exceed the recommended daily dosage.

If you take more USN PHEDRACUT LIPO X BLACK LIQUID than you should

In the event of an overdose, consult your doctor or pharmacist.

If neither is available, contact the nearest hospital or poison centre.

If you forget to take USN PHEDRACUT LIPO X BLACK LIQUID

Take your next dose at the normal time.

Do not take a double dose to make up for individual forgotten doses.

4. Possible side effects

USN PHEDRACUT LIPO X BLACK LIQUID can have side effects.

Not all side effects reported for USN PHEDRACUT LIPO X BLACK LIQUID are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking USN PHEDRACUT LIPO X BLACK LIQUID, please consult your healthcare provider for advice.

If any of the following happens, stop taking USN PHEDRACUT LIPO X BLACK LIQUID and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Swelling of your hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing.
- Rash or itching.
- Fainting.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to USN PHEDRACUT LIPO X BLACK LIQUID. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

Frequency unknown:

- Changes in the way your heart beats, for example, if you notice it beating faster than normal.
- If you pass more urine than is normal for you.

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequency unknown:

- Nausea (feeling sick), vomiting (being sick), stomach pain, stomach cramps, heartburn, diarrhoea.
- Anxiousness, nervousness, restlessness, agitation, headache, dizziness.
- Insomnia (difficulty falling or staying asleep).
- Tremors (uncontrolled shaking).
- Increased blood pressure (headache, dizziness).
- Flushing (reddening of the skin).

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the 6.04 Adverse Drug Reactions Reporting Form, found online under SAHPRA's publications: <http://www.sahpra.org.za/Publications/Index/8>.

By reporting side effects, you can help provide more information on the safety of USN PHEDRACUT LIPO X BLACK LIQUID.

5. How to store USN PHEDRACUT LIPO X BLACK LIQUID

- Store at or below 25 °C.
- Refrigerate after opening.
- Use within 1 month after first opening of the container.
- Store all medicines out of reach of children.
- Do not use after the expiry date printed on the container.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What USN PHEDRACUT LIPO X BLACK LIQUID contains

Each 25 ml contains:

Ascorbic acid (Vitamin C)	500 mg	*%NRV
L-Carnitine L-Tartrate	500 mg	500 %
N-Acetyl L-Tyrosine	200 mg	
Caffeine Anhydrous	200 mg	
<i>Fucus vesiculosus</i> L. (Kelp) [whole herb; extract providing 10 % fucoxanthin]	120 mg	
Raspberry Ketone [fruit; extract providing 98 % p-hydroxyphenyl butanone]	100 mg	
Thermozine™ <i>Aframomum melegueta</i> K. Schum (Grains of Paradise) [seed; 10:1 extract providing 400 mg of dried herb equivalent]	40 mg	
Nicotinic acid (Vitamin B3)	17,5 mg	109 %
Pyridoxine HCl providing Pyridoxine (Vitamin B6)	0,33 mg	
Chromium Picolinate providing Chromium (elemental)	0,27 mg	16 %
	402 µg	
	50 µg	143 %</td