


Compound Data Sheet

Structure:



Name	17α-Methyltestosterone
Synonym(s)	17α-Methyl-17β-hydroxyandrost-4-en-3-one
Epichem Code #	EPL-DV4 Batch 1
CAS #	58-18-4
Molecular Formula	C ₂₀ H ₃₀ O ₂
Molecular Weight	302.46 g/mol
Purity	99.4% at 242nm by HPLC (assuming all components detected with the same response factor). No impurities detected by ¹ H NMR analysis. LC/MS: [M+H] ⁺ = 303.3
Appearance	White solid
Melting point	160.2-166.7°C
Date of Manufacture	3 April 2019
Storage Requirements	Avoid exposure to heat, light and moisture.
Special Precautions	This compound is for laboratory use only. Its toxicological properties may not have been fully established. It should be handled only by suitably qualified personnel.
Date of Shipment	TBA
Retest Date	TBA (Proper Storage and Handling Required)

Document prepared by:

Checked by:

James Rixson, PhD
Head of Production
20 October 2020

Boon Tan
Quality Manager
20 October 2020

I. Identity

The identity of this product was established using the following analyses:


Ia. ¹H NMR Spectrum

Conditions: 400 MHz, DMSO-d₆

¹H NMR spectrum consistent with chemical structure.

Acquisition Time (sec)	3.7547	Comment	LBC266-193@23H 1H DMSO (E:\data\external\epichem) cyqoh 6		
Date	21 Jun 2019 17:08:16	Date Stamp	21 Jun 2019 17:08:16		
File Name	\naphthalene\company\NMR files\LBC266-193@23H\1\data\111r		Frequency (MHz)	400.13	
Nucleus	1H	Number of Transients	8	Origin	spect
Owner	nmr	Points Count	32768	Original Points Count	24038
SW(cyclical) (Hz)	6402.05	Solvent	DMSO-d6	Pulse Sequence	zg
Sweep Width (Hz)	6401.85	Temperature (degree C)	24.996	Receiver Gain	144.00
				Spectrum Offset (Hz)	2800.9097
				Spectrum Type	STANDARD


LBC266-193@23H.001.001.1r.esp



Ib. Mass Spectrum

The mass spectrum of this material was analysed by Liquid Chromatography Mass Spectroscopy (LCMS) using in-house EM005.WI08.

Retention Time (MS)	MS Area	Mol. Weight or Ion
3.164	2394756	383.20 I
		304.20 I
		303.25 I




Theoretical value: 303.3 [M+H]⁺.

The signal of the Mass Spectrum is consistent with the theoretical value and its interpretation is consistent with the structural formula.

Ic. IR Spectrum

The infra-red spectrum of this material was analysed by Fourier-Transform Infrared Spectroscopy (FTIR) using in-house EM005.WI09.




The interpretation of the signals of the Fourier-Transform Infrared Spectrum is consistent with the structural formula.

II. Purity

The purity of this material was analysed by high performance liquid chromatography (HPLC) using in-house EM005.WI07.

HPLC Conditions:

Column	Conditions				Detector	Injector
Agilent Zorbax SB-C18 4.6 x 250mm 5.0 micron	25°C				DAD 242nm	Auto 1.0 µL 1.0 mg/mL 100% acetonitrile
	Time (min)	% Line A (Water + 0.1% (v/v) TFA)	% Line B (Acetonitrile + 0.1% (v/v) TFA)	Flow rate (mL/min)		
	0.00	50	50	1.5		
	9.00	5	95	1.5		
	14.00	5	95	1.5		
	15.00	50	50	1.5		
20.70	50	50	1.5			



Area Percent Report – Sorted by Signal

Peak Number	Retention Time (rounded)	Area	Area % (rounded)
1	2.07	0.02	0.00
2	2.17	2.17	0.11
3	2.49	4.01	0.20
4	2.80	0.81	0.04
5	2.97	0.10	0.01
6	3.32	0.86	0.04
7	5.04	0.65	0.03
8	5.33	0.53	0.03
9	5.62	0.37	0.02
10	5.82	1960.61	99.37
11	6.76	2.93	0.15
Totals			100 (rounded)

For the calculation the system peaks were ignored. The content of the analyte was determined as a ratio of the peak area of the analyte and the cumulative areas of the purities, added up to 100%.

Results:

Average 99.4% (average duplicate analyses)