



URINE DRUG SCREEN

Multi-Drug Urine Test Dip Card

MRK220006

Single Test or Multi-Drug Panel

A comprehensive approach to screening for the presence of multiple drugs within urine. Based on the principle of lateral flow immunoassay technology, urine drug screens present qualitative results in a matter of minutes. A simple test procedure with easy to interpret and quick results make this an ideal drug screen within various settings.

FEATURES

- Qualitative detection of various drugs and/or their respective metabolites in urine
- Optional urine specimen validity tests provide a semi-quantitative screen for Oxidants, Specific Gravity, pH, Nitrile, Glutaraldehyde, and Creatine
- 25 Tests/Kit, 20 Kits/Case
- Customizable test configurations
- OEM and Private-Label options are available

BENEFITS

- FDA-Cleared and CLIA-Waived (See test menu on back page)
- 24-Month shelf life from date of manufacture
- On-site screening
- Internal, procedural control included
- Qualitative results in 5 minutes





TEST MENU

Multi-Drug Urine Dip Card

MRK220006

All-in-one urine-based test that can be utilized in practically any drug screening scenario. Our drug test dip cards can offer panel sizes up to 14 analytes with an additional 6 specimen validity tests.

FDA 510k Cleared for Over-the-Counter (OTC)	
Test	Cut-off
Amphetamine (AMP)	500 ng/mL
	1000 ng/mL
Barbiturates (BAR)	300 ng/mL
Buprenorphine (BUP)	10 ng/mL
Benzodiazepines (BZO)	300 ng/mL
Cocaine (COC)	150 ng/mL
	300 ng/mL
Ecstasy (MDMA)	500 ng/mL
Marijuana (THC)	50 ng/mL
Methadone Metabolite (EDDP)	300 ng/mL
Methadone (MTD)	300 ng/mL
Methamphetamine (MET)	500 ng/mL
	1000 ng/mL
Morphine (MOP)	300 ng/mL
Opiates (OPI)	2000 ng/mL
Oxycodone (OXY)	100 ng/mL
Phencyclidine (PCP)	25 ng/mL
Propoxyphene (PPX)	300 ng/mL
Tricyclic Antidepressants (TCA)	1000 ng/mL


For Forensic Use Only*	
Test	Cut-off
Alcohol	0.04%
Clonazepam (CLO)	100 ng/mL
Cotinine (COT)	200 ng/mL
Ethyl Glucuronide (EtG)	300 ng/mL
	500 ng/mL
Fentanyl (FEN)	10 ng/mL
	20 ng/mL
Gabapentin (GAB)	1000 ng/mL
Hydrocodone (HCD)	300 ng/mL
Hydromorphone (HMO)	300 ng/mL
Ketamine (KET)	1000 ng/mL
Kratom (KRA)	250 ng/mL
LSD	20 ng/mL

For Forensic Use Only*	
Test	Cut-off
Methaqualone (MQL)	300 ng/mL
Methcathinone (MTC)	300 ng/mL
Bath Salts (MDPV)	1000 ng/mL
6-monoacetylmorphine (6-MAM)	10 ng/mL
AB-Pinaca (K3)	10 ng/mL
Pregabalin (PGB)	1000 ng/mL
Synthetic Marijuana (K2)	25 ng/mL
	50 ng/mL
Tramadol (TRA)	100 ng/mL
	200 ng/mL
UR-144	50 ng/mL
Zolpidem (ZOL)	300 ng/mL
Zopiclone (ZOP)	50 ng/mL

TEST PROCEDURE

Allow the test dip card to come to room temperature prior to test.

- 1) Remove the test dip card from the foil pouch.
- 2) Remove the cap from the test dip card. Label the dip card with patient or control identifications.
- 3) Immerse the absorbent tip into the urine sample for 10-15 seconds. Urine sample should not touch the plastic dip card.
- 4) Replace the cap over the absorbent tip and lay the dip card flatly on a non-absorptive clean surface.
- 5) Read results at 5 minutes.



*Forensic Use Only (FUO) devices are intended for use in drugs of abuse testing for law enforcement purposes within the United States. FUO devices are not designed, tested, developed, or labeled for use in other settings, such as clinical diagnostics.