



Bio BasicsTM
www.biobasics.biz

Crop Specific Instructions

APPLES 1 pint/acre each application

1st application: At full pink

3rd application: 3 weeks after 2nd spraying

2nd application: At calyx (petal fall)

4th application: 4 weeks after 3rd spraying

CARROTS 1 pint/acre each application

1st application: At tuber initiation

2nd application: 2 to 3 weeks after 1st spraying

CELERY

1st application: Use 2 pints/acre applied to the seed bed at time of seeding or up to 20 days thereafter. 2nd application: Use 2 pints/acre at the time seedlings are transplanted. 3rd application: Use 1 pint/acre 2 to 3 weeks after transplanting.

CORN 1 pint/acre each application

1st application: At the 1 to 1 ½ foot stage

2nd application: At tassel time

COTTON

Pinhead square: Apply 2 to 4 ounces/acre weekly for 4 weeks beginning at pinhead square. First bloom: Apply ½ pint/acre at first white flower and again two weeks later. Stripper cotton: Apply single application of ½ pint/acre during first two weeks of bloom.

GRAPES

1 pint/acre each application

1st application: Between leaf out and prebloom

2nd application: At petal fall

3rd application: After harvest

ORANGES 1 pint/acre each application

1st application: At prebloom

3rd application: 3 weeks after 2nd spraying

2nd application: At calyx (petal fall)

4th application: 4 weeks after 3rd spraying

PEACHES 1 pint/acre each application

1st application: At prebloom

2nd application: At calyx (petal fall)

3rd application: 3 weeks after 2nd spraying

4th application: 4 weeks after 3rd spraying

PEANUTS 1 pint/acre each application

1st application: At pegging

2nd application: 2 to 3 weeks after 1st application

PEPPERS 1 pint/acre each application

1st application: Just prior to first bloom

2nd application: 10 days after 1st spraying

3rd application: 10 days after 2nd spraying

4th application: 10 days after 3rd spraying

POTATOES 1 pint/acre each application

1st application: At tuber set

2nd application: At full blossom

Russet Burbanks which do not show full blossom should be sprayed 2 to 3 weeks after 1st spraying.

SOYBEANS 1 pint/acre each application

Application: At first bud formation

STRAWBERRIES

1st application: Use 2 pints/acre as a transplant solution

2nd application: Use 1 pint/acre at prebloom

3rd application: Use 1 pint/acre at petal fall

4th application: Use 1 pint/acre after harvest

SUGAR BEETS 1 pint/acre each application

1st application: At tuber initiation

2nd application: 2 to 3 weeks after 1st spraying

TOMATOES

1st application: Use 2 pints/acre applied to the seedbed at time of seeding or up to 20 days thereafter. 2nd application: Use 2 pints/acre at the time seedlings are transplanted. 3rd application: Use 1 pint/acre 2 to 3 weeks after 1st bloom.

WHEAT 1 pint/acre each application

Application: 1 to 2 weeks before boot stage