

Fillers

Lip Filler

- Mini Lip Lift: \$400
- Full Lip Lift: \$695

Dermal Fillers & Injectables

- **Juvederm:** cheek, chin, jawline, tear through, temples: \$600-\$800.
- **Volite** an anti aging hyaluronic acid treatment that hydrates, plumps skin and smooths fine lines: \$695/syringe.

Before & After



Fillers

Dermal Filler Instructions

Pre-Treatment

- Schedule your Dermal Filler appointment at least 2 weeks prior to a special event which you may be attending, such as a wedding or a vacation. Bruising and swelling may be apparent and take up to 2 weeks to heal.
- Avoid alcoholic beverages at least 24 hours prior to treatment (alcohol may thin the blood and increase the risk of bruising).
- Avoid anti-inflammatory/blood thinning medications (unless prescribed by your physician) if possible, for a period of 5 days before treatment. Medications and supplements such as acetylsalicylic acid (ASA), vitamin E, ginkgo biloba, ginseng, St. John's Wort, Omega 3/Fish Oil supplements, ibuprofen, naproxen sodium and other NSAIDS have a blood thinning effect and can increase the risk of bruising and swelling after injections.
- Reschedule your appointment at least 24 hours in advance (to avoid cancellation fees) if you have a rash, cold sore or blemish on the area to be treated.
- If you have a history of cold sores and you are having your lips or mouth area treated, please pretreat with anti-viral medication prior to treatment (as prescribed by your physician)
- We recommend taking Arnica Montana (herb) 2 days prior to injections to reduce bruising and swelling.
- If you choose, please take 5 pellets 3 times per day 2 days prior to treatment, the day of your treatment, and 2 days after your treatment. You may purchase at most natural food stores and/or Amazon.
- You should avoid having dental work 2 weeks before or after receiving filler.
- Be sure to have a good meal, including food and drink before your procedure. This will decrease the chance of lightheadedness or low blood sugar during your treatment.
- If possible avoid drinking caffeine the day of your procedure.