


Pocketbook

Radiology

Romée Snijders, MD & Veerle Smit, MD
Gwendolyn Vuurberg, MD, PhD

QUALITY
GUARANTEED
BY MEDICAL
SPECIALISTS



Compendium
medicine

A completely new pocket on diagnostic imaging and radiographic findings in the most important acute diagnoses.

The Compendium Method®

Manual

In *Compendium Medicine* we use the same concise, visual and schematic description of the various medical specialties as much as possible. Everything is geared towards overview and structure, facilitating study and practice. We call this the Compendium Method.

Fixed layout

All our medical specialties are presented in the same, recognisable way and each has its own colour and icon. The pocketbooks have a fixed chapter structure. The table of contents of each pocketbook tells you exactly which topics are covered. The symbols in the corner of the page indicate the specialty or chapter.

- ATLS
- Anatomy
- Physiology
- Patient history
- Physical examination
- Diagnostics
- Treatment
- Differential diagnosis
- Conditions
- Clinical reasoning
- Appendixes
- References
- Abbreviations
- Index

Illustrations

The images provide at-a-glance insight into topics like anatomy or the typical patient. They are also intended for study and practice, for example by checking whether you can identify the letters in a picture without looking at the caption.




Figure 1 // Anatomical planes
A: Coronal/frontal **B:** Median **C:** Sagittal **D:** Axial/transverse




Conditions

Each condition in this pocketbook starts with a full-sentence definition, followed by a telegram-style explanation. For each condition the following icons (as applicable) are discussed. The icons are also useful when studying: you can cover the text and quiz yourself.

- D** Definition
- E** Epidemiology
- Ae** Aetiology
- R** Risk factors
- Hx** Patient history
- PE** Physical examination
- DDx** Differential diagnosis
- Dx** Diagnostics
- Tx** Treatment
 - General
 - Paramedical care
 - Drug treatment
 - Invasive, non-medicated treatment
- P** Prognosis
- !** Watch out/don't forget

Tables

We make as much use as possible of tables, for example to compare conditions. In this way, the differences are immediately clear. Features that match are centred over the columns to which they apply. This allows you to see the similarities and differences right away.

| | D | Hx PE | DDx | Dx | ! |
|--|---|---|-----|----|---|
| | |    | | | |
| | | | | | |








Diagrams

→ = positive/yes/+ → = negative/no/-

Diagrams help you reason clinically starting from a particular symptom, using the green and red arrows as signposts. Always remember that the full differential diagnosis may consist of multiple diagnoses.










Icons & frames

Throughout this pocketbook you will find highlighted frames.

| | | |
|--|--|--|
|  QR code |  Note |  Reference to another chapter |
|  Alarm! |  Description of the typical patient |  Formula |
|  Mnemonic | | |

Punctuation marks

The punctuation in our books also focuses on overview and ensures that the subject matters are covered concisely and effectively.

| | | |
|---|--|--|
|  Rare |  Most common |  Decrease |
|  Uncommon |  Consequence |  Female sex |
|  Very common |  Increase/improvement |  Male sex |

Abbreviations

We make extensive use of abbreviations, medical terms and symbols for scientific units and quantities. Below are some examples of the abbreviations used in this pocketbook.

| | | | |
|-----|----------------|------|--------------|
| sec | second/seconds | mo | month/months |
| min | minute/minutes | min. | minimum |
| hr | hour/hours | max. | maximum |
| d | day/days | e.g. | for example |
| wk | week/weeks | L | litres |

Index

The pocketbooks include a comprehensive and easy-to-use index. It contains all the topics covered in the books so you can quickly navigate and find the information you are looking for.

Appendixes

In the pocketbooks you will find space for your notes. In addition, handy appendixes have been added; these contain specific information that you would like to have at hand and are therefore located at the back of the pocketbooks.

His/her

We realise that sex and gender identity are not binary and that there is more variation than just 'woman' or 'man'. For the sake of the readability, however, we have chosen to use the pronouns he/him when referring to anyone, regardless of sex or gender identity.



Want to know more about the Compendium Method? Scan the QR code.



Table of contents

Radiology

ATLS trauma care

- ABCDE approach
- Secondary survey

Anatomical planes

Imaging modalities

- Ultrasound
 - Basics
 - Tissue differentiation
 - FAST ultrasound
 - Abdominal ultrasound
 - Doppler and Duplex ultrasound
 - Conventional X-ray
 - Basics
 - Chest
 - Abdomen
 - Bones
 - Conventional X-ray examination
 - Digital subtraction angiography (DSA)
 - Fluoroscopy beyond the radiology department
 - Computed tomography (CT)
 - Basics
 - Tissue differentiation
 - Head-neck
 - Chest
 - Abdomen
 - Bones
- #### Magnetic resonance imaging (MRI)
- Basics
 - Tissue differentiation
 - Contrast agents
 - Head-neck
 - Abdomen

- Bones
 - Nuclear medicine
 - Basics
 - Planar scintigraphy
 - Pulmonary scintigraphy (V/Q scan)
 - Single photon emission computed tomography (SPECT)
 - Positron emission tomography (PET)
 - Dual-energy X-ray absorptiometry (DEXA) scan
 - Overview of nuclear imaging modalities
 - Comparison of imaging modalities
- #### Contrast
- Contrast agents
 - Contrast phases
 - Imaging modalities using contrast
 - Conventional X-rays and fluoroscopy
 - Ultrasound
 - CT scan
 - MRI scan
 - Angiography and venography
 - Contrast allergy
 - Contrast-induced nephropathy (CIN)
- #### Imaging requests
- RI-RADS
 - Examples of imaging requests
- #### Pointers for imaging during pregnancy
- Pulmonary embolisms during pregnancy
- #### Radiological signs of child abuse

Trauma mimics

- Secondary ossification centre
- Epiphysis
- Apophysis
- Accessory ossification centres
- Accessory ossicles and sesamoids
- Anatomical variations and physiological development
- Haversian canals
- Vertebral variations
- Intercarpal congruence
- Cranial sutures

Invasive diagnostics and treatment

- General
- Elective procedures
 - Biopsies and punctures
 - Peritoneal dialysis catheter (PD catheter)
 - Radiofrequency ablation (RFA) and cryoablation
- Drainages and ascites aspiration
 - Ascites aspiration and drainage
 - Abscess drainage
 - Gallbladder drainage
 - Nephrostomy catheter
- Endovascular procedures
 - Venous access
 - Intra-arterial thrombectomy (IAT)
 - Embolisation
 - Percutaneous transluminal angioplasty (PTA)
 - Thrombolysis

Conditions

Head-neck

- Intracranial haemorrhage
 - Epidural haematoma
 - Subdural haematoma
 - Subarachnoid haemorrhage
 - Parenchymal haemorrhage
 - Thrombus
 - Ischaemic stroke (Cerebrovascular Accident (iCVA))
 - Cerebral venous thrombosis (CVT)
 - Mass effect
 - Herniation
 - Hydrocephalus
 - Infectious conditions
 - Cerebral abscess
 - Retropharyngeal abscess
 - Trauma
 - Skull base fracture
 - Facial fracture
- #### Spine
- Traumatic spinal injury
 - NEXUS criteria
 - Denver criteria
 - Traumatic cervical spine injury
 - Vertebral fracture
 - Epidural haematoma
 - Non-traumatic spinal injury
 - Spondylodiscitis
 - Torticollis
- #### Thorax
- Lines and tubes
 - Pulmonary pathology
 - Pneumonia
 - Empyema
 - Congestive heart failure (CHF)

Table of contents

Radiology

Thoracic trauma
 Pneumothorax
 Haemothorax
 Pulmonary haemorrhage
 Pulmonary contusion
 Rib fracture
 Vascular conditions
 Pulmonary embolism
 Aortic dissection

Abdomen

Lines and tubes
 Acute pathology
 Appendicitis (acute)
 Cholecystitis
 Pancreatitis
 Diverticulitis
 Mesenteric lymphadenitis
 Necrotising enterocolitis (NEC)
 Vascular conditions
 Intestinal ischaemia
 Infarction of parenchymal organs
Renal infarction
Splenic infarction
 Organ haemorrhage
Liver laceration
Spleen laceration
Kidney laceration
 Postoperative haemorrhage
 Ruptured acute abdominal
 aortic aneurysm (AAA)
 Gastrointestinal conditions
 Intussusception
 Intestinal volvulus
Children

Adults
 Internal herniation
 Adynamic ileus
Paralytic ileus
Mechanical ileus
 Gastrointestinal perforation
 Urogenital conditions
 Nephrolithiasis and urolithiasis
 Hydronephrosis
 Epididymitis
 Testicular torsion
 Ovarian torsion
 Ectopic pregnancy
 Pelvic inflammatory disease (PID)
 Miscellaneous
 Ascites
 Choledocholithiasis

Extremities

AO classification
 Shoulder
 AC dislocation
 Clavicular fracture
 Shoulder dislocation
 Proximal humerus fracture
 Elbow/underarm
 Elbow dislocation
 Radial head subluxation
 Supracondylar humeral fracture
 Olecranon fracture
 Proximal radial fracture
 Forearm fractures
 Galeazzi fracture
 Monteggia fracture
 Essex-Lopresti fracture

Hand/wrist
 Distal radius fracture
Colles fracture
Smith fracture
 Scaphoid fracture
 Volar plate avulsion injury
 Mallet finger
 Skier's thumb
 Metacarpal fractures (2-5)
 Boxer's fracture
 Phalangeal fractures
 Pelvis/hip
 Pelvic fracture
 Collum fracture
 Hip dislocation
 Knee/lower leg
 Tibial plateau fracture
 Patellar fracture
 Patellar dislocation
 Ankle/foot
 Ankle fracture
 Lisfranc (dislocation) injury
 Miscellaneous
 Muscle/tendon rupture
 Ligamentous injury
 Avulsion fracture
 Pathological fracture
 Deep vein thrombosis (DVT)
 Critical limb ischaemia (CLI)

Non-radiological diagnoses

Head-neck
 Epileptic seizure
 Meningitis
 Chest

Acute respiratory distress syndrome (ARDS)
 Myocardial infarction
 Pericarditis
 Tension pneumothorax
 Abdomen
 Cholangitis
 Hepatitis
 Pancreatitis
 Extremities
 Pulled elbow

Clinical reasoning

Diagnostic evaluation of dyspnoea
 Diagnostic evaluation of abdominal pain

Reference list

Illustrations

Epilogue

About us

Abbreviations

Index

Contrast agents

During imaging studies, patients may be administered a contrast agent. Contrast agents are chemical agents that provide better contrast and make it easier to differentiate tissues. Contrast agents are usually administered intravenously, as this enables assessment of e.g. abdominal organs or blood vessels and high-light pathology susceptible to contrast enhancement, such as tumours or vascular malformations. Contrast agents may also be administered orally or rectally to assess post-operative anastomotic leakage or assess the course of and passage through the digestive tract, for example. Contrast agents can also be injected intra-articularly to better assess intra-articular structures like the labrum. See table 1 for the most commonly used types of contrast agents and the corresponding imaging techniques and routes of administration.

When using contrast, patients may be scanned in several phases, following the contrast agent through the body as it passes various organs. The appropriate scanning phase therefore depends on the clinical question.

| CONTRAST AGENT | IMAGING TECHNIQUES | SPECIFIC TARGET ORGAN | ROUTE OF ADMINISTRATION |
|------------------------|--|---|--|
| Barium sulphate | Conventional (swallow test, dynamic rectal exam - DRE), CT | Digestive tract | Oral, rectal |
| Iodinated | CT, conventional (e.g. for choking or anastomotic leakage) | Varying, scan phase depends on contrast agent's route of administration | Intravenous, oral, urethral, intra-articular |
| Gadolinium | MRI | - | Intravenous, oral, intra-articular |
| Primovist | | Liver/bile ducts | Intravenous |

Table 1 // Various contrast agents with their corresponding imaging techniques and routes of administration

Guidelines for imaging during pregnancy

During pregnancy, the fetus is highly susceptible to the adverse effects of radiation and drugs. This is because rapid and frequent cell division takes place in an embryo/fetus, which makes DNA extra susceptible to iatrogenic damage from X-rays and other sources (see table 2). In addition to the fetus, any other rapidly dividing tissue, such as mammary tissue, has an increased risk of iatrogenic damage from X-rays. It is important to keep health risks for both mother and child in mind when requesting certain types of imaging.

Although the degree of sensitivity of these tissues depends on the stage of pregnancy, the X-ray recommendations are the same at each stage.



During X-ray imaging, avoid using a lead apron or lead screens. The current generation of X-ray cameras automatically adjust the radiation dose based on the amount of radiation reaching the detector plate (Automatic Exposure Control (AEC)). Using a lead apron/lead screens can increase scatter radiation. The number of rays reaching the detector plate will also decrease, prompting the X-ray camera to automatically increase the radiation dose.



Ultrasound devices of the radiology department use different settings compared to ultrasound devices used in obstetrics and by gynaecologists. Even though ultrasounds are considered safe during pregnancy, it is recommended not to directly image the fetus or use Doppler ultrasound to image adjacent structures during early pregnancy.



| IMAGING MODALITY | RISKS DURING PREGNANCY | | RISKS DURING BREASTFEEDING | CONSIDERATIONS |
|--|---|--|---|--|
| Conventional X-ray exam | Low radiation exposure → mildly teratogenic | | None | Only if strictly necessary. Try to limit radiation exposure to a minimum. |
| Ultrasound | None | | | Can be safely used during pregnancy |
| CT scan • Child | High radiation exposure → teratogenic | | None (if contrast is used - see below) | <ul style="list-style-type: none"> Consider whether an ultrasound or MRI is a suitable alternative. If this is not possible, try to minimise radiation exposure. Radiation dose depends on the type of scan and stage of pregnancy (e.g. CT for pulmonary embolisms with indirect foetal radiation before the end of the 3rd trimester vs. CT abdomen with direct radiation). |
| • Mother | During pregnancy and after delivery, mammary tissue is at risk of iatrogenic damage due to its proliferation in preparation for the lactation period | | During pregnancy and after delivery, mammary tissue is at risk of iatrogenic damage due to its proliferation in preparation for the lactation period | |
| MRI scan | Can be safely used during pregnancy >18 wks. Possible risks for the child in the 1 st trimester have not yet been fully investigated. Likely no risks at low magnetic field strength (≤1.5 Tesla). | | None | <ul style="list-style-type: none"> The risks posed by MRI during pregnancy have not yet been fully investigated. MRI is always preferred over a CT scan in pregnant women, if possible |
| Contrast exam | For all types of contrast agent, carefully consider whether the exam must be performed during pregnancy and whether the use of contrast agent is necessary | | For all types of contrast agent, carefully consider whether the exam must be performed during pregnancy and whether the use of contrast agent is necessary | |
| • Iodinated contrast agent (IV) | Probably safe. The risk of affecting foetal thyroid gland function seem small. | | Safe. A small amount of contrast agent enters breast milk (iodine 1%, gadolinium 0.04%), and it is poorly absorbed by the neonate's gastrointestinal tract. | The use of contrast agent is not recommended unless it significantly improves the diagnostic process and therefore the foetal and/or maternal outcome |
| • Gadolinium-based contrast agent (IV) | Unknown. In patients, gadolinium is deposited in the brain. Effect on fetus unknown. | | | |
| Barium sulphate (oral) | Unknown | | Unknown | The risks of barium sulphate are unknown, so it is generally not recommended in pregnant women. An iodinated oral contrast agent diluted with water can serve as an alternative. |

Table 2 // Imaging risks during pregnancy

Radiologic signs of child abuse

Child abuse, also known as 'non-accidental injury' (NAI), is a difficult subject. Doctors rarely expect to find child abuse and do not want to assume that injuries are inflicted deliberately. Confronting the parents can also be challenging and the resulting investigation may have a significant impact on both the parents and the environment of the child. This is why it is important to recognise the signs of NAI on conventional imaging and conduct a thorough and repeated history, preferably including a collateral history with eyewitnesses (e.g. other parents) to the trauma.

On suspicion of child abuse, imaging can play a vital role. Situations in which to suspect child abuse:

- Injury inconsistent with the reported trauma mechanism;
- Injury in an unusual location;
- Injury inconsistent with the child's developmental stage;
- Long delay before seeking medical help.

If a particular fracture is inconsistent with the child's age (e.g. a femoral fracture in an infant), the person assessing the image should be extra alert to the possibility of child abuse (see table 3). In case of strong suspicions of child abuse, a comprehensive skeletal survey may be performed to identify occult or old fractures that support the suspicion.



Injuries raising the suspicion of child abuse are sometimes described as 'non-accidental injury' (NAI) in the radiological report.



It is important to always consider whether an injury is consistent with the trauma mechanism. For example, a transverse humeral fracture supposedly resulting from a fall is highly suspicious, while a direct impact injury (e.g. kick from a horse) is less suspicious.



When in doubt, consultation between the requesting physician and radiologist is important!

| TYPE OF FRACTURE | TRAUMA MECHANISM | SPECIAL ATTENTION IS WARRANTED IN: |
|--|--|--|
| Combination of sternum fractures, scapula fractures, posterior rib fractures/spinous process fractures, skull fractures and/or brain injuries (subdural haematoma (SDH), subarachnoid haemorrhage and cerebral oedema) | Severe shaking/anterior-posterior force (ie. shaken baby syndrome) Slap on the back (fracture spinous process) | <ul style="list-style-type: none"> • All children, but pay special attention to children who are not yet able to walk or children who are regularly presented with injuries. • Impaired consciousness. This may be indicative of an SDH. |
| Metaphyseal corner fractures and avulsion fractures | Push and pull forces, shaking the child back and forth by holding the torso as the head and limbs move back and forth (shaken baby syndrome) | |
| <ul style="list-style-type: none"> • Multiple fractures in different stages of healing • Signs of fractures sustained at ages of the child | Repeated trauma | |
| Vertebral fractures, vertebral compression fractures | Compression (axial impact) | |
| Femur fracture, humeral fracture and radius/ulna fracture | High impact force, acceleration-deceleration forces | |
| Transverse or spiral fractures of the long bones | Rotational forces | |

Table 3 // Radiologic signs of child abuse.

Red: very specific **Orange:** moderately specific **Yellow:** little specific.

In children with fractures, always consider whether the trauma mechanism is consistent with the injury (see figure 1). Blaming a brother/sister or the absence of parents are warning signs in the patient history. Always double-check the patient history, ask about what happened several times, preferably speak to both parents separately and, if possible, have another adult testify in case of strong suspicions of child abuse.

Proper screening is also important, as some underlying conditions can closely resemble non-accidental injury on diagnostic imaging, such as haemorrhages in coagulopathy, skeletal abnormalities in connective tissue disorders (including osteogenesis imperfecta), metabolic diseases or skeletal dysplasia. Even normal anatomical variation may resemble (abusive) injuries (see section Trauma mimics).



For more information, see the section on Trauma mimics.




Figure 1 // Probability of child abuse based on fracture findings



Suspect **H**arm from **M**other **OR** **F**ather:
S: sternum, scapula, spine/vertebrae
H: humerus, head, hands*
M: multiple fractures, metaphyseal corner or other avulsion fractures
O: old fractures
R: ribs
F: femur*, feet*
 * Especially suspicious in children who haven't started walking.




Figure 2 // Signs of shaken baby syndrome



Endovascular procedures

| | CVL | PICC LINE | MIDLINE CATHETER |
|---------------------------------------|--|--|------------------|
| Placement • Insertion point | <ul style="list-style-type: none"> Internal jugular vein Axillary/subclavian vein Superficial femoral vein | <ul style="list-style-type: none"> Brachial vein Brachiocephalic vein | |
| • Catheter tip | Superior/inferior vena cava | | Axillary vein |
| Indications | Long-term antibiotic (AB) use | | |
| | <ul style="list-style-type: none"> Central access Parenteral nutrition Chemotherapy Central AB therapy | Venous access for at least ± 2 wks | |
| | <ul style="list-style-type: none"> Haemodialysis Central venous pressure measurement (CVP) | Venous access for at least ±4 wks | |
| Contraindications | Home treatment | Relative: PICC line via left arm in case of pacemaker/ICD due to conflict between catheter and leads | |
| | | Acute situation (CVL preferred) | |
| Complications | Thrombus risk ↑ (CVL preferred) | | |
| | <ul style="list-style-type: none"> Arrhythmias Haemo-/pneumothorax | | |
| | | <ul style="list-style-type: none"> Arterial puncture Sepsis (Line tip) thrombosis | |

Table 4 // Types of venous access



A midline catheter is very similar to a PICC line, but is not centrally located because the tip of the catheter does not pass the axillary vein.



After the central line has been inserted, a chest X-ray should be made to assess its position and potential complication.



Not all forms of venous access are suitable for administering contrast during imaging, mainly due to the size of the lumen and the corresponding flow rate. Check this beforehand to avoid the risk of the line breaking in the patient.

Venous access

The peripheral venous catheter is a common form of venous access. For longer-term venous access or for certain medications, a more centrally placed line may be required (see figure 2). Depending on the indication, you can choose between a central venous line (CVL), peripherally inserted central catheter (PICC) line or midline catheter (see table 4). Depending on which medication is to be given through the line, one, two or three lumens can be chosen.




Figure 2 // A: CVL via jugular vein B: CVL via subclavian vein D: Femoral catheter via femoral vein C: PICC line via brachiocephalic vein



Acute pathology

| CONDITION | D | Hx PE | DDx | Dx | ! |
|-----------------------------|--|--|---|--|--|
| Appendicitis (acute) | Acute inflammation of the wall of the vermiform appendix located in the extension of the caecum near Bauhin's valve. | <ul style="list-style-type: none"> PE Point tenderness over McBurney's point, psoas sign + PE CRP ↑, leukocytes ↑ | <p>Adults:</p> <ul style="list-style-type: none"> Gastroenteritis Cholecystitis Right-sided diverticulitis IBD (Crohn's disease/colitis ulcerosa) Tubo-ovarian abscess (TOA)/pelvic inflammatory disease (PID) <p>Children:</p> <ul style="list-style-type: none"> Gastroenteritis Mesenteric lymphadenitis Intussusception (inflamed) Meckel's diverticulum | <ul style="list-style-type: none"> Abdominal ultrasound: demonstrating appendicitis (pain near appendix, incompressible appendix, transverse diameter appendix >6 mm, wall thickening >2 mm, peri-appendicular inflamed fat, free fluid, appendiceal faecoliths (see figure 3). CT abdomen with IV contrast if ultrasound is inconclusive despite strong suspicion: appendix diameter >6 mm, appendiceal faecoliths, peri-appendicular inflamed fat Abdominal MRI in pregnant women and sometimes in children if ultra-sound is inconclusive despite strong suspicion: diameter of appendix diameter >6 mm, wall thickening, appendiceal faecoliths, restricted diffusion | <ul style="list-style-type: none"> Complicated appendicitis: presence of faecoliths, disappearance of mucosal layer, abscess formation and/or suspected perforation Free air secondary to perforation of the appendix is rare In obese patients, a non-contrast CT abdomen may suffice because the intestinal loops are further apart due to mesenteric fat ↑, increasing the visibility of local inflammations. Symptoms secondary to malignancy, e.g. mucinous cystadenoma |
| Cholecystitis | Acute inflammation of the gallbladder usually due to an obstructing gallstone in the neck of the gallbladder or cystic duct. | <ul style="list-style-type: none"> H Colic pain ☹ PE Murphy sign + PE CRP ↑, leukocytes ↑ | <ul style="list-style-type: none"> Cholelithiasis Choledocholithiasis Pancreatitis Acute hepatitis Appendicitis (retrocaecal or elevated caecum) | <p>Abdominal ultrasound: demonstrating cholecystitis - sonographic Murphy sign, incompressible gallbladder (hydrops), gallstones, wall thickening (>3 mm) (see figure 4). Depending on the location of the obstruction, dilated bile ducts may also be seen in the liver and pancreas.</p> | <ul style="list-style-type: none"> Possible symptoms with an obstructing/passed stone in the choledochal duct (duct diameter >5 mm) Gangrenous inflammation with abscess formation or perforation of the gallbladder. Cholelithiasis is an important risk factor for acute cholecystitis. In case of recurrent symptoms, elective cholecystectomy may be considered. Acalculous cholecystitis: cholecystitis without obstructive concrement (2-18%) |
| Pancreatitis | Acute inflammation of the pancreas, usually caused by gallstones or alcohol consumption. | <ul style="list-style-type: none"> H Epigastric pain (radiating to back) PE Peritoneal excitation +/- PE Lipase ↑, (amylase ↑), CRP ↑ | <ul style="list-style-type: none"> Ulceration Cholelithiasis Gastrointestinal perforation | <ul style="list-style-type: none"> CT abdomen with IV contrast: oedematous pancreatic parenchyma, peripancreatic fat stranding, locoregional lymphadenopathy Important: pancreatitis is primarily diagnosed based on clinical and lab findings. A CT can help assess severity and necrotising component approx. 3 days after onset of symptoms. | <ul style="list-style-type: none"> Abdominal ultra-sound does not rule out pancreatitis and therefore has little added value. In biliary pancreatitis, ultra-sound can be used to detect gallstones. In pancreatitis due to an obstructing stone near Vater's papilla, an ERCP may be performed by the gastroenterologist Cave necrotising pancreatitis involving extensive necrotising fluid collections due to the lytic properties of the released pancreatic enzymes |

Tabel 5A // Acute pathology of the abdomen



In children, referred pain due to pneumonia can mimic abdominal conditions.



Inflamed fat is also known as *fat stranding*.



| CONDITION | D | Hx PE | DDx | Dx | ! |
|--|---|---|---|---|--|
| Diverticulitis | Inflammation of one or more (false) diverticula (bulging pouches in the intestinal wall) usually of the sigmoid and colon, but possibly also of the duodenum. | <ul style="list-style-type: none"> H Pain (mostly) in lower left abdomen, altered bowel habits, rectal bleeding PE T ↑ CRP ↑, leukocytes ↑ | <ul style="list-style-type: none"> Gastroenteritis Obstipation IBD (Crohn's disease/colitis ulcerosa) Appendicitis | <ul style="list-style-type: none"> Abdominal CT: presence of diverticula, perifocal inflamed fat, free fluid and possible abscess formation CT abdomen with IV contrast: diverticula with fat stranding, possibly (covered) perforation and abscess formation | <ul style="list-style-type: none"> In Caucasian patients, diverticula are mainly (95%) localised in the sigmoid, in patients with an Asian background, they are mainly (75%) found in the ascending colon In cases of suspected complicated diverticulitis (with abscess formation), CT abdomen is the first choice, as deep-seated abscesses are easier to miss on ultrasound |
| Mesenteric lymphadenitis | Inflammation of the lymph nodes of the abdominal cavity, occurs mostly in children <15 years of age. | <ul style="list-style-type: none"> Presentation similar to appendicitis: H Abdominal pain over McBurney's point, N+, V+ CRP ↑ | <ul style="list-style-type: none"> Appendicitis Invagination Obstipation | Abdominal ultrasound: rule out appendicitis, ≥3 enlarged lymph nodes in the lower abdomen (>5 mm) | Mesenteric lymphadenitis is a diagnosis of exclusion, self-limiting and often does not require treatment |
| Necrotising enterocolitis (NEC) | Life-threatening neonatal intestinal infection often associated with intestinal ischaemia. Does not occur outside the neonatal period. | <ul style="list-style-type: none"> H Rectal bleeding, (bilious) vomiting PE Abdominal distension and tenderness, enhanced vein definition, signs of shock | <ul style="list-style-type: none"> Infectious enteritis/colitis Spontaneous intestinal perforation Volvulus Intussusception Meconium ileus Cow's milk protein allergy Hirschsprung or other congenital disorders | Abdominal X-ray:dilated intestinal loops, intestinal pneumatosis, air in portal veins and, in case of perforation, free air in the abdominal cavity | <ul style="list-style-type: none"> Risk factors: preterm birth (< 32 wks), dysmaturity High mortality (15-30%) Granular faeces (stool with soap bubble sign on abdominal X-ray) does not occur in the first weeks after birth, so this could be consistent with intramural gas. |

Table 5B // Acute pathology of the abdomen




Figure 3 // Acute appendicitis with wall thickening (red arrow) up to 7.1 mm, peri-appendicular fat infiltration (blue arrow) and peri-appendicular fluid (white arrow)





Figure 4 // Cholecystitis secondary to cholelithiasis.

Hydropic gallbladder with wall thickening (red arrow) secondary to an obstructing 22 mm stone in the neck of the gallbladder (white arrow).

Enjoyed the Preview? Get the Full Pocket Radiology Now!



- ✓ Imaging modalities and technical details
- ✓ What is needed for a radiology request
- ✓ Invasive diagnostic procedures and treatments performed at the radiology department
- ✓ A range of acute conditions

BUY NOW



"I sincerely believe this is the best radiological textbook for beginners: up-to-date, practical, and clear figures."

Prof. P. Simoni, MD, PhD, MBA - Radiologist