



Dicamba Agricultural Herbicide

For weed control in corn, cotton, sorghum, small grains, pasture, hay, rangeland, general farmstead (non-cropland), rights-of-way, public utility and industrial areas, fallow, sugarcane, asparagus, turf and grass seed crops.

Active Ingredient:

Dimethylamine salt of dicamba (3, 6-dichloro-*o*-anisic acid)* 49.77%

Inert Ingredients: 50.23%

Total 100.00%

*This product contains 41.35% 3, 6-dichloro-*o*-anisic acid (dicamba) or 4 pounds per gallon (480 g/L).

Keep Out of Reach of Children

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See additional Precautionary Statements and Directions for Use inside booklet.

Oracle® is a registered trademark of Gharda Chemicals International, Inc.

EPA Reg. No.: 93182-10

EPA Est. No.: 2217-KS-1^P

EPA Est. No.: 44616-MO-2^H

EPA Est. No.: 19713-TN-8

(Superscript designates first letter of lot number on jug.)

Net Contents: 2.5 Gallons

Distributed By:



Gharda Chemicals International, Inc.

760 Newtown-Yardley Rd., Suite 110

Newtown, PA 18940

TABLE OF CONTENTS

	Page
Precautionary Statements	3
Directions for Use	4
General Information	4 - 8
Mixing and Application	4
Best Stewardship Practices	4
Ground and Surface Waters Protection	5
Sensitive Crop Precautions	5
Band Treatments	6
Compatibility Test	6
Procedure For Cleaning Spray Equipment	6
General Weed List	7 - 8
Field, Seed, Popcorn and Silage Corn	9 - 13
Cotton (Preplant Application)	13
Sorghum (Milo)	13 - 14
Small Grains (Wheat, Barley, and Oats)	14 - 19
Fall and Spring Seeded Wheat	16 - 17
Fall Seeded Barley	18
Spring Seeded Barley	19
Fall and Spring Seeded Oats	19
Sugarcane	20
Pasture, Hay, Rangeland and General Farmstead (non-Cropland), Rights-of-way, and Public Utility and Industrial areas	20 - 22
Cut Surface Tree Treatments	22
Dormant Applications for the Control of Multiflora Rose	22
Conservation Reserve Program (CRP)	22 - 23
Asparagus	23
Turf and Lawns	23 - 24
Grass Seed Crops	25
Preplant Directions (Post-Harvest/Fallow/Crop Stubble/Set-A-Side) for Broadleaf Weed Control Before Wheat, Corn, Sorghum, Soybeans	26 - 28
Control of Perennial Broadleaf Weeds in Cropland (Spot Application Only)	28 - 29
Wiper Application Uses	29
Storage and Disposal	29
Notice of Warranty and Disclaimer	29
Registered Trademarks	30

ORACLE[®] Dicamba Agricultural Herbicide

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed.

FIRST AID (Substituted Benzoic acid)	
If swallowed:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Remove person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER (Substituted Benzoic acid) Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: 1-(866)-359-5660	

Personal Protective Equipment (PPE)

Some material that is chemical-resistant to this product is natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks
- Chemical-resistant gloves (except for applicators using groundboom equipment, pilots and flaggers)
- Protective eyewear

See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as directed on label.

DIRECTIONS FOR USE**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeve shirt and short pants
- chemical-resistant footwear plus socks
- chemical-resistant gloves made of any waterproof material
- chemical-resistant headgear for overhead exposure
- protective eyewear

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

Before applying ORACLE, read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

INFORMATION

The following directions apply to all uses of ORACLE. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

Mixing and Application

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF ORACLE. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

ORACLE is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see COMPATIBILITY TEST on page 6) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of a diluted spray per treated acre when using ground application equipment, or 1 to 10 gallons of a diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, ORACLE should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

Best Stewardship Practices

ORACLE provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Ground and Surface Waters Protection

1. Point source contamination - To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment, or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.

2. Movement by surface runoff or through soil - Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the general information section of this label.
3. Movement by water erosion of treated soil - Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

Sensitive Crop Precautions

ORACLE may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems, or foliage. These plants are most sensitive to ORACLE during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING ORACLE.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of ORACLE with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10 or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gpa, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply ORACLE adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
- To avoid injury to desirable plants, equipment used to apply ORACLE should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT on page 6) before reusing to apply any other chemicals.

All crop uses of ORACLE are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. **Tank mix recommendations are for use only in states where the tank mix product and application site are registered.**

Band Treatments

ORACLE may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per treated acre} = \text{Band RATE per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per treated acre} = \text{Band VOLUME per treated acre}$$

Compatibility Test

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**Amount of Herbicide to Add to One Pint of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)**

Herbicide Formulations	Rate Per Acre	Level Teaspoons
Dry	1 lb.	1 1/2
Liquid	1 pt.	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Re-run the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

Procedure For Cleaning Spray Equipment

The steps listed below are suggested for thorough cleaning of spray equipment following applications of ORACLE or tank mixes of ORACLE or tank mixes of ORACLE plus 2,4-D amine.

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.
The steps listed below are suggested for thorough cleaning of spray equipment used to apply ORACLE as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. ORACLE tank mixes with water-dispersible formulations require the use of a water/detergent rinse.
5. Complete step 1.
6. Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
7. Flush the detergent solution out of the spray tank through the boom.
8. Repeat step 1, and follow with steps 2, 3, and 4.

General Weed List

This is a general list of weeds which may be treated with ORACLE in accordance with this label as specified under the rates and timing sections of the Individual Use Headings. Proper usage of this product will give control or growth suppression of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species including:

Annuals

Amaranth, Spiny
(Spiny Pigweed)
Aster, Slender
Bedstraw
Beggarweed, Florida
Broomweed, Common
Buckwheat, Wild
Buffalobur
Burclover, California
Burdock
Buttercup, Roughseed
Carpetweed
Catchfly, Nightflowering
Chamomile, Corn
Chickweed, Common
Clovers (Annual)
Cockle, Corn
Cockle, Cow
Cocklebur, Common
Croton, Tropic
Croton, Woolly
Daisy, English
Evening primrose, Cutleaf
Fleabane, Annual
Goosefoot, Nettleleaf
Henbit
Jimsonweed
Knotweed
Kochia
Ladysthumb
Lambsquarters, Common
Lambsquarters (triazine resistant)
Lettuce, Prickly
Mallow, Common
Mallow, Venice
Mayweed
Morningglory, Ivyleaf
Morningglory, Tall
Mustard, Tansy
Mustard, Wild
Mustard (Yellowtops)
Nightshade, Black
Pennycress, Field (Fanweed,
Frenchweed, Stinkweed)
Pepperweed, Virginia
(Peppergrass)
Pigweed, Prostrate
Pigweed, Redroot
(Carelessweed)
Pigweed, Rough
Pigweed, Smooth
Pigweed (triazine resistant)
Pigweed, Tumble
Poorjoe
Puncturevine
Purslane, Common
Pusley, Florida
Radish, Wild
Ragweed, Common
Ragweed, Giant (Buffaloweed)
Ragweed, Lance-Leaf
Rubberweed, Bitter
(Bitterweed)
Sesbania, Hemp
Shepherdspurse
Sicklepod
Sida, Prickly (Teaweed)
Smartweed, Green
Smartweed, Pennsylvania
Sneezeweed, Bitter
Sowthistle, Annual
Sowthistle, Spiny

Annuals (continued)

Spikeweed, Common
Spurge, Prostrate
Spurry, Corn
Starbur, Bristly
Sumpweed, Rough
Sunflower, Common (Wild)
Sunflower, volunteer
Thistle, Russian
Velvetleaf
Waterhemp
Waterprimrose, Winged
Wormwood, Annual

Biennials

Burdock, Common
Carrot, Wild (Queen Anne's
Lace)
Cockle, White
Evening primrose, Common
Geranium, Carolina
Gromwell
Knapweed, Diffuse
Knapweed, Spotted
Mallow, Dwarf
Plantain, Bracted
Ragwort, Tansy
Starthistle, Yellow
Sweetclover
Teasel
Thistle, Bull
Thistle, Milk
Thistle, Musk
Thistle, Plumeless

Perennials

*Alfalfa
 Artichoke, Jerusalem
 Aster, Spiny
 Aster, Whitehead
 Beadstraw, Smooth
 Bindweed, Field
 Bindweed, Hedge
 Blueweed, Texas
 *Bursage, (Bur Ragweed, Lakeweed, Povertyweed)
 Bursage, Woollyleaf (Lakeweed)
 Buttercup, Tall
 Campion, Bladder
 Chickweed, Field
 Chickweed (Mouseear, Canada)
 Chicory
 *Clover, Hop
 *Dandelion, Common
 *Dock, Broadleaf (Bitterdock)
 *Dock, Curly
 Dogbane, Hemp
 *Dogfennel (Cypressweed)
 Fern, Bracken
 Garlic, Wild
 Goldenrod, Canada
 Goldenrod, Missouri
 Goldenweed, Common
 Hawkweed
 Horsenettle, Carolina
 Ironweed
 Knapweed, Black
 Knapweed, Russian
 Mare's Tail (Horseweed)

Milkweed, Climbing
 Milkweed, Common
 Milkweed, Honeyvine
 Milkweed, Western Whorled
 Nettle, Stinging
 Nightshade, Silverleaf (White Horsenettle)
 Onion, Wild
 *Plantain, Broadleaf
 Plantain, Buckhorn
 Pokeweed
 Ragweed, Western
 Redvine
 Sericia Lespedeza
 Smartweed, Swamp
 Snakeweed, Broom
 *Sorrel, Red (Sheep Sorrel)
 Sowthistle
 Sowthistle, Perennial
 Spurge, Leafy
 Sundrop, Halfshrub (Evening Primrose)
 Thistle, Canada
 Toadflex, Dalmation
 Tropical Soda Apple
 Trumpetcreeper (Buckvine)
 Vetch
 Waterhemlock
 Waterprimrose, creeping
 *Woodsorrel, Creeping
 Common Yellow
 Wormwood, Common
 Wormwood, Louisiana
 *Yankeeweed
 Yarrow, Common

Woody

Alder
 Ash
 Aspen
 Basswood
 Beech
 Birch
 **Blackberry
 **Blackgum
 **Cedar
 Cherry
 Chinquapin
 Cottonwood
 **Creosotebush
 Cucumbertree
 **Dewberry
 **Dogwood
 Elm
 Grape
 **Hawthorn (Thornapple)
 Hemlock
 Hickory
 Honeylocust
 Honeysuckle
 Hornbeam
 Huckleberry
 Huisache
 Ivy, Poison
 Kudzu
 **Growth Suppression

Locust, Black
 Maple
 Mesquite
 Oak
 Oak, Poison
 Olive, Russian
 Persimmon, Eastern
 Pine
 **Plum, Sand (Wild Plum)
 Poplar
 Rabbitbrush
 **Redcedar, Eastern
 **Rose, McCartney
 **Rose, Multiflora
 Sagebrush, Fringe
 Sassafras
 Serviceberry
 Spicebush
 Spruce
 Sumac
 **Sweetgum
 Sycamore
 Tarbush
 Willow
 Witchhazel
 **Yaupon
 **Yucca

*Noted perennials may be controlled using ORACLE at rates lower than those specified for other listed perennial weeds. (See APPLICATION RATES AND TIMINGS section on page 21.

FIELD, SEED*, POPCORN* AND SILAGE CORN

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4 as well as the following:

* Do not apply ORACLE to seed corn or popcorn without first verifying with your local seed corn company (supplier) the ORACLE selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

ORACLE is not registered for use on sweet corn.

Direct contact of ORACLE with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the surface, delay application until corn has emerged.

Up to 2 applications of ORACLE may be made during a growing season. Do not exceed a total 1 1/2 pints of ORACLE per treated acre per crop year. Allow two weeks or more between applications of ORACLE. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

Applications of ORACLE to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizers (1/2-1 gallon per acre of 28%, 30%, 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate*) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after crop emergence or crop injury may result.

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tank mix applications with ORACLE. Refer to their label for specific recommendations.

* Not for use in California

Weeds Controlled

ORACLE will control many ANNUAL broadleaf weeds or give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn. (Refer to the GENERAL WEED LIST on pages 7 and 8).

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

Preplant/Preemergence In No-Tillage Corn

Applications of ORACLE may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply ORACLE at 1 pint per treated acre on medium or fine textured

soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply ORACLE after 4-6 inches of regrowth has occurred.

Preemergence In Conventional or Reduced Tillage Corn

ORACLE may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of ORACLE does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

Early Postemergence (All Tillage Systems)

Spike through 8 inch tall corn

ORACLE at 1 pint per treated acre may be applied during the period from corn emergence through the five leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 1/2 pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, loamy sand). See Late Postemergence applications given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

Late Postemergence (All Tillage Systems)

8 to 36 inches tall corn

Application of ORACLE at 1/2 pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when: (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D.

DO NOT apply ORACLE when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

Overlay (Sequential) Treatments

ORACLE may be applied to ground previously treated with one or more of the following herbicides:

acetochlor (Surpass[®], Harness[®])

alachlor (Lasso[®])

atrazine

Bicep[®] II Magnum

Bullet[®]

butylate (Sutan+[®])

Clarity[®]

dimethenamid (Frontier[®])

EPTC (Eradicane[®])

Guardsman[®]

glyphosate (Roundup[®])

halosulfuron (Permit[®])

Lariat[®]

Marksman[®]

s-metolachlor (Dual[®] II Magnum)

paraquat (Gramoxone[®])

pendimethalin (Prowl[®])

simazine (Princep[®])

Apply ORACLE at 1/2 pint per treated acre to ground previously treated with full rates of Clarity[®] or Marksman[®] herbicides. Allow at least 2 weeks between applications.

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS.

Tank Mix Treatments

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds in corn. **Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions.**

Rates and Timings

ORACLE Plus	Preplant/Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Post-Emergent (All Tillage Systems)	Late Post-Emergent (All Tillage Systems)	Additional Directions
ACCENT® (nicosulfuron)	—	—	1/2-1 oz. ai/A	1/2-1 oz. ai/A (to improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.)	Application may be made to emerged weeds before corn is greater than 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this tank mixture.
atrazine	1 1/4-2 lbs. ai/A	1 1/4-2 lbs. ai/A	1 1/4-2 lbs. ai/A Crop oil concentrates may be used with this mixture if corn is 5 inches or less in height.	1 1/4-2 lbs. ai/A Do not apply if corn is greater than 12 inches tall.	Application may be made before grasses are 1 1/2" tall. Follow all state and Federal restrictions pertaining to atrazine applications.
BEACON® (primisulfuron)	—	—	0.31-0.62 oz. ai/A	0.31-0.62 oz. ai/A (to improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.)	Application may be made to emerged weeds when corn is 4 to 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this tank mixture.
DUAL® II Magnum/II Magnum SI (metolachlor)	0.8-1.6 lbs. ai/A	0.8-1.6 lbs. ai/A (use only on fine or medium soils with 2.5% or greater organic matter.)	0.8-1.6 lbs. ai/A	—	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall.

Rates and Timings (continued)

ORACLE Plus	Preplant/ Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Post-Emergent (All Tillage Systems)	Late Post-Emergent (All Tillage Systems)	Additional Directions
FRONTIER® (dimethenamid)	13-25 fl. oz/A	13-25 fl. oz/A (use only on fine or medium textured soils with 2.5% or greater organic matter.)	13-25 fl. oz/A	—	Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides postemergence control of grass weeds if they are greater than 1 inch tall at the time of application.
GRAMOXONE® (paraquat)	1/4-1 lb. ai/A	1/4-1 lb. ai/A	—	—	Application may be made to emerged weeds but prior to corn emergence.
HARNESS® or SURPASS® (acetochlor)	1 1/2-3 lbs. ai/A	1 1/2-3 lbs. ai/A (use only on fine or medium textured soils with 2.5% or greater organic matter.)	—	—	Application should be made prior to corn emergence.
LASSO® (alachlor)	1 1/2-4 lbs. ai/A	1 1/2-4 lbs. ai/A (use only on fine textured soils with greater than 2.5% organic matter).	1 1/2-4 lbs. ai/A	—	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall. If microencapsulated forms of alachlor are used (Lasso MT, Partner), applications must be made prior to grass emergence.
PRINCEP® (simazine)	2.0-3.0 lbs. ai/A	2.0-3.0 lbs. ai/A	—	—	Application may be made prior to corn or weed emergence.

Rates and Timings (continued)

ORACLE Plus	Preplant/ Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Post-Emergent (All Tillage Systems)	Late Post-Emergent (All Tillage Systems)	Additional Directions
PROWL® (pendimethalin)	—	3/4-1 1/2 lbs. ai/A (use only on fine or medium textured soils with 2.5% or greater organic matter.)	3/4-1 1/2 lbs. ai/A	—	Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2 leaf stage of growth.
ROUNDUP® (glyphosate)	1.0-3.0 lbs. ai/A	1.0-3.0 lbs. ai/A	—	—	Application may be made to emerged weeds but prior to corn emergence.
2,4-D	1/4-1/2 lbs. ai/A	1/4-1/2 lbs. ai/A	Do not use at this application timing	1/8 lbs. ai/A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of the weed foliage.

COTTON (Except California)

PREPLANT APPLICATION: Apply up to 8 fluid ounces of ORACLE per acre to control emerged broadleaf weeds prior to planting cotton in conventional or conservation tillage systems.

For best performance, apply ORACLE when weeds are in the 2 – 4 leaf stage and rosettes are less than 2" across.

Following application of ORACLE and a minimum accumulation of 1" of rainfall or overhead irrigation, a waiting interval of 21 days is required per 8 fluid ounces per acre or less. These intervals must be observed prior to planting cotton.

Do not apply preplant to cotton west of the Rockies.

Do not make ORACLE preplant applications to geographic areas with average annual rainfall less than 25".

If applying a spring preplant treatment following application of a fall preplant (postharvest) treatment, then the combination of both treatments may not exceed 2 lbs. acid equivalent per acre.

Tank Mix Treatments

For control of grasses or additional broadleaf weeds, ORACLE may be tank mixed with Bladex®, Caparol®, Gramonone® Extra, and Roundup® herbicides

SORGHUM (Milo)

Observe all PRECAUTIONS on pages 3 to 5, including the reference to crops growing under stress. Read and follow MIXING AND APPLICATION instructions on page 4.

Applications of ORACLE to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. Preharvest Interval (PHI) restrictions are as follows: grain sorghum – 30 days; fodder – 30 days; and forage – 20 days.

If sorghum is grown for pasture or hay, refer to the pasture use section of this label. Do not apply ORACLE to sorghum grown for seed production.

Make no more than one application per growing season.

Weeds Controlled

ORACLE, when applied at the specified rate for sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as control their seedlings. (Refer to GENERAL WEED LIST on pages 7 and 8.)

Rates and Timings

ORACLE may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of ORACLE must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3-5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

Broadcast Rate Per Treated Acre:

1/2 pint (1/4 lb. ai)

Tank Mix Treatment**ORACLE plus Atrazine**

For improved control of emerged, actively growing broadleaf weeds including triazine resistant species and added suppression of perennial broadleaf weeds, tank mix 1/2 pint ORACLE with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 1/2 pint ORACLE with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all state and Federal restrictions pertaining to atrazine applications.

ORACLE plus Buctril®

For improved control of broadleaf weeds, tank mix 1/2 pint ORACLE with 1-1 1/2 pint Buctril Herbicide per treated acre. Make application at 4 leaf to 15 inch tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS.

Overlay (Sequential) Treatments

ORACLE may be applied to ground previously treated with one or more of the following herbicides:

Herbicide	Maximum Rate Per Treated Acre (lbs. ai)
atrazine ¹	2.5
s-metolachlor (Dual® II Magnum) (Concep®-treated seed)	1.33

¹ Maximum use rate for atrazine is determined by soil type, tillage practices used, surface residue, and state or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

Preharvest Uses**For Use Only in the States of Texas and Oklahoma**

ORACLE may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications use at least 2 gallons of water-based carrier per treated acre.

Delay harvest until 30 days after treatment.

Broadcast Rate Per Treated Acre:

1/2 pint (1/4 lbs. ai)

**SMALL GRAINS (WHEAT, BARLEY AND OATS)
NOT UNDERSEEDED TO LEGUMES****Important**

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

Preharvest Interval (PHI) restriction for grain is 7 days.

If small grains are used for pasture hay, the following restrictions apply:

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
- Do not harvest hay from treated areas before 37 days after treatment.

NOTE: Observe all precautions and restrictions on the labels of products used in tank mix treatments.

Weeds Controlled

ORACLE, or combinations with listed tank mix partners, will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that ORACLE be applied in a tank mix with other herbicides. Refer to specific crop tank mix options.

Alkanet ¹	Chervil, Bur ¹
Bedstraw, Catchweed ¹	Chickweed, Common ¹
Bindweed, Field ²	Cockle, Corn
Buckwheat, Tartary	Cockle, Cow
Buckwheat, Wild	Cocklebur, Common
Carpetweed ¹	Cornflower (Bachelorbutton) ¹
Chamomile, Corn	Dandelion, Common ²

Dock, Curly ²	Pennycress, Field (Fanweed,
Dragonhead, American ¹	Frenchweed, Stinkweed)
Evening Primrose, Cutleaf ¹	Pepperweed, Peppergrass ¹
Falseflax, Smallseeded ¹	Pigweed, Redroot
Fiddleneck (Tarweed) ¹	(Carelessweed)
Flixweed ¹	Pigweed, Rough
Fumitory ¹	Pigweed, Tumble
Gromwell, Corn ¹	Pineappleweed ¹
Groundsel, Common ¹	Plantain, Broadleaf ²
Hempnettle ¹	Poppy, Red Horned ¹
Henbit	Puncturevine ¹
Jacobs Ladder ¹	Purslane, Common ¹
Knawel (German Moss)	Radish, Wild ¹
Knotweed, Prostrate	Ragweed, Common ¹
Kochia	Ragweed, Giant
Ladysthumb	(Buffaloweed) ¹
Lambsquarters, Common	Rocket, London ¹
Lettuce, Miners ¹	Rocket, Yellow ¹
Lettuce, Prickly	Salsify (Goatsbeard) ¹
Mallow, Common	Shepherdspurse ¹
Mayweed, Chamomile	Smartweed, Green
(Dogfennel) ¹	Smartweed, Pennsylvania
Mustard, Blue (Purple) ¹	Sorrel, Red (Sheep Sorrel) ¹
Mustard, Tansy	Sowthistle, Annual
Mustard, Treacle ¹	Starthistle, Yellow ¹
Mustard, Tumble (Jim Hill) ¹	Sunflower, Common (Wild)
Mustard, Wild ¹	Thistle, Canada ²
Nightshade, Black	Thistle, Russian
Nightshade, Cutleaf ¹	Yarrow, Common ²
Nightshade, Silverleaf ²	Velvetleaf
(White Horsenettle)	Vetch ¹

¹ These weeds will be controlled with ORACLE tank mixtures. Refer to tank mix label for specific weeds controlled.

² ORACLE tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

Rates and Timings

Application of ORACLE may be made before, during or after planting small grains. For best performance, make applications when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Application of ORACLE to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use ORACLE at 2 to 4 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 2 to 3 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, Russian thistle and prickly lettuce or dense vegetative growth.

ORACLE used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for ORACLE rate and application timing.

For applications prior to the emergence of weeds or when sulfonylurea resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of ORACLE with a tank mix herbicide. Non-sulfonylurea herbicides such as 2,4-D or MCPA tank mixed with ORACLE will offer more consistent control of sulfonylurea resistant weeds.

When tank mixing with sulfonylurea herbicides, such as Ally®, Amber®, Express®, Finesse®, Glean® and Harmony® Extra, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

FALL AND SPRING SEEDED WHEAT

ORACLE MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT EXCEEDS THE 5 LEAF STAGE. Early developing wheat varieties such as TAM 107, MADISON, or WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

Tank Mix Treatments

ORACLE may be tank mixed with one or more, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

Broadcast Rate Per Treated Acre:

Apply 2-4 fluid ounces of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A) ¹
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A) ¹
Ally®	metsulfuron-methyl	60% DF	1/10 oz.
Amber®	triasulfuron	75% DF	0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/3 oz.
Buctril®	bromoxynil ²	2 lb/gal	1-1 1/2 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	1-2 pts.
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	2-2 2/3 pts.
Stinger®	clopyralid	3 lb/gal	1/4-1/3 pt.
Karmex® ³	diuron ²	80% DF	1/2-1 1/2 lbs.
Sencor® ³	metribuzin ²	75% DF	1-10 oz.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

² Herbicides with the same active ingredient and/or different formulation may be used.

³ Tank mixtures for fall seeded wheat only.

**Special Use Tank Mixes For Spring and Fall Seeded Wheat
(See Footnotes for Applicable Uses)**

Apply 3-4¹ fluid ounces of ORACLE with:

Product ²	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D or MCPA Amine	2,4-D or MCPA	4 lb/gal	1-2 pts. ³ (0.5-1.0 lb ai/A) ⁴
2,4-D or MCPA Ester	2,4-D or MCPA	4 lb/gal	1-1.5 pts. ³ (0.5-0.75 lb ai/A) ⁴
Ally®	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber®	triasulfuron	75% DF	0.14-0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Ally® + 2,4-D Amine or Ester ⁵	metsulfuron-methyl + 2,4-D	60% DF+ 4 lb/gal	1/20-1/10 oz. + 8 fl oz.
Amber® + 2,4-D Amine or Ester ⁵	triasulfuron + 2,4-D	75% DF + 4 lb/gal	0.14-0.28 oz. + 8 fl oz.
Express® + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/12-1/6 oz + 8 fl oz.
Harmony® Extra + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/6-1/3 oz + 8 fl oz.
Roundup® ⁶	glyphosate	3 lb/gal	12-16 fl oz.

¹ ORACLE may be used at 6 fluid ounces on fall seeded wheat in Western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of ORACLE may be applied on fall seeded wheat after it exceeds the 3 leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. ORACLE may be tank mixed with 2,4-D amine at 8 fluid ounces after wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For Fall applications only, do not use if the potential for crop injury is not acceptable.

² Do not use low rates of sulfonyleurea herbicides (such as Ally®, Amber®, Express®, Finesse®, Glean®, and Harmony® Extra) on more mature weeds and/or on dense vegetative growth.

³ NOTE: For use on Fall Seeded Wheat Only. Do Not Use unless potential crop injury will be acceptable.

⁴ When using formulations other than 4 lb/gal use pounds active/acre listed.

⁵ Use for improved control of Russian thistle, flixweed, gromwell, mayweed and fiddleneck.

⁶ ORACLE may be applied at 2 fluid ounces with Roundup® as a preplant application to small grains with no waiting period prior to planting. Add 0.5% by volume of an agriculturally approved non-ionic surfactant.

FALL SEEDED BARLEY

ORACLE MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

NOTE: For spring barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

Tank Mix Treatments

ORACLE may be tank mixed with one or more, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

Broadcast Rate Per Treated Acre:

Apply 2-4 fluid ounces of ORACLE with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8 fl oz. (0.25 lb ai/A) ²
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (0.25-0.375 lb ai/A)
Ally®	metsulfuron-methyl	60% DF	1/20 - 1/10 oz.
Amber®	triasulfuron	75% DF	0.14 - 0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12 - 1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6 - 1/3 oz.
Sencor®	metribuzin ³	75% DF	1-10 oz.
Buctril®	bromoxynil	2 lb/gal	1-1 1/2 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4-1 1/2 pts.

¹ Do not use low rates of sulfonylureas (Ally®, Amber®, Express®, and Harmony® Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

SPRING SEEDED BARLEY

ORACLE MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4 LEAF STAGE.

Tank Mix Treatments

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

Broadcast Rate Per Treated Acre:

Apply 2-3 fluid ounces of ORACLE with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (0.25-0.375 lb ai/A) ²
Ally®	metsulfuron-methyl	60% DF	1/20 - 1/10 oz.
Amber®	triasulfuron	75% DF	0.14 - 0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12 - 1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6 - 1/3 oz.
Sencor®	metribuzin ³	75% DF	1-10 oz.
Buctril®	bromoxynil	2 lb/gal	1-1 1/2 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4-1 1/2 pts.

¹ Do not use low rates of sulfonylureas (Ally®, Amber®, Express®, and Harmony® Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

FALL AND SPRING SEEDED OATS

ORACLE MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5 LEAF STAGE. APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE.

Tank Mix Treatments

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

Broadcast Rate Per Treated Acre:

Apply 2-4 fluid ounces of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (0.25-0.375 lb ai/A) ¹

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

SUGARCANE

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

Preharvest Interval (PHI) restriction for sugarcane is 87 days.

Weeds Controlled

ORACLE, when applied at specified rates, will control many ANNUAL, BIENNIAL and PERENNIAL broadleaf weeds commonly found in sugarcane. (Refer to GENERAL WEED LIST on pages 7 and 8.)

Rates and Timings

Application of ORACLE may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timings of ORACLE are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage and Type	Amount Product	Broadcast Rate Per Treated Acre lbs. ai
Annual		
Small, actively growing	1/2-1 pt	1/4 - 1/2
Established weed growth	1-1 1/2 pts	1/2 - 3/4
Biennial	1-2 pts	1/2 - 1
Perennial	2 pts	1 ¹

¹ Application made over the top of actively growing sugarcane may result in crop injury.

When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

Retreatment may be made as needed, however, do not exceed a total of 2 applications of ORACLE per year.

Tank Mix Treatments

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

Herbicide	Rates per treated acre (lbs. ai)
Ametryn (Evik®)	2/5 - 8
asulam (Asulox®)	2 - 3 1/3
atrazine	2/5 - 4
2,4-D	1/2 - 3 ¹

¹ Application of ORACLE plus 2,4-D tank mix at the higher listed rate range may result in crop injury.

PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (Non-Cropland)

ORACLE is recommended for use on pasture, hay, rangeland, general farmstead (non-cropland, including fence rows and non-irrigation ditchbanks) for broadleaf weed and brush control. ORACLE may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds must be recognized at the State level but programs may be administered at State, County or other levels.

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

ORACLE uses described in this section also pertain to small grains (such as barley, forage, sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

NEWLY SEEDED AREAS, including small grains grown for pasture may be severely injured if rates of ORACLE greater than 1 pint/acre are applied.

ESTABLISHED GRASS CROPS growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 1 pint ORACLE (1/2 lb. ai) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

ANIMALS CANNOT BE REMOVED FROM TREATED AREA FOR SLAUGHTER PRIOR TO 30 DAYS AFTER LAST APPLICATION.

THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND GRAZING FOR NON-LACTATING ANIMALS.

Timing Restrictions for Lactating Dairy Animals**Following Treatment**

ORACLE Rate per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb. ai)	7 days	37 days
Up to 1 quart (1 lb. ai)	21 days	51 days

NOTE: Observe all precautions and restrictions on labels of products used in tank mixtures.

Mixing and Application

ORACLE can be applied using water, oil in water emulsions (including invert systems), or sprayable fluid fertilizer as a carrier. A COMPATIBILITY TEST (page 6 of this booklet) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

ORACLE may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment apply 1 to 40 gallons of diluted spray per treated acre in a water-based carrier.

ORACLE may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run-off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

Weeds and Brush Controlled

ORACLE, when applied at specified rates, will give control of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species commonly found in pasture, hay, rangeland, and general farmstead (non-cropland) areas. (Refer to GENERAL WEED LIST on pages 7 and 8.)

Noted (*) PERENNIAL weeds may be controlled with lower rates of either ORACLE or ORACLE plus 2,4-D. See RATES AND TIMINGS below.

Rates and Timings

Application rates and timing of ORACLE are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage and Type	Amount Product	Broadcast Rate Per Treated Acre lbs. ai
Annual Small, actively growing Established weed growth	1/2 - 1 pt. 1 - 1 1/2 pts.	1/4 - 1/2 1/2 - 3/4
Biennial ¹ Rosette diameter Less than 3 inches 3 inches or more Bolting	1/2 - 1 pt. 1 - 2 pts. 2 pts.	1/4 - 1/2 1/2 - 1 1
Perennial Suppression or top growth control Control of noted (*) Perennials	1/2 - 1 qt. 1 qt.	1/2 - 1 1
Woody Brush & Vines Top growth suppression Top growth control ²	1/2 - 1 qt. 1 qt.	1/2 - 1 1

¹ For best performance, make application when BIENNIAL weeds are in the rosette stage.

² Species noted in GENERAL WEED LIST, pages 7 and 8, will require tank mixtures for adequate control.

Retreatment may be made as needed, however, do not exceed a total of 2 applications of ORACLE per year.

Tank Mix Treatments

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND OTHER RESTRICTIONS.

ORACLE may be tank mixed with one or more, but not limited to, the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines.

Herbicide	Rates Per Treated Acre (lbs. ai)
Pasture, hay, rangeland and general farmstead (non-cropland) use:	
glyphosate (Roundup®)	3/4 to 3 3/4
metsulfuron methyl (Ally®)	0.0038 to 0.011
paraquat (Gramoxone®)	1/2 to 1
picloram (Tordon®)	1/8 to 3
triclopyr (Garlon®)	3/4 to 9
2,4-D	1/4 to 6

Due to the variations that may occur on formulated products and specific use ingredients (e.g. water supplies), a COMPATIBILITY TEST as described on page 6 is recommended prior to actual tank mixing.

CUT SURFACE TREE TREATMENTS

ORACLE may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix of 1 part ORACLE with 1 to 3 parts water should be used in application. Use the lower dilution when treating difficult-to-control species.

Frill or Girdle Treatments: Make a continuous cut or a series of overlapping cuts using an ax to girdle tree trunk. Spray or paint cut surface with the ORACLE/water mix.

Stump Treatments: Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

NOTE: For more rapid foliar effects, 2,4-D may be added to the ORACLE/water mix.

DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE

ORACLE can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of ORACLE should be applied directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying ORACLE directly to the soil. The use rate of ORACLE is dependent on the canopy diameter of the multiflora rose. Examples: Use ORACLE at 1/4, 1 or 2 1/4 fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2 qts. ORACLE per acre per year.

LO-OIL BASAL BARK applications of ORACLE should be applied to the basal stem region from the ground up to a height of 12-18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying ORACLE to the ground line. Refer to Mixing and Applications above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a Lo-Oil spray solution mixture: combine 1 1/2 gallons water plus 1 ounce emulsifier plus 1 pint ORACLE plus 2 1/2 pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

CONSERVATION RESERVE PROGRAM (CRP)

ORACLE is recommended for use on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION directions on page 4.

ORACLE treatment will cause injury or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

Newly Seeded Areas

ORACLE may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses

exceed the 3 leaf stage. Rates of ORACLE greater than 1 pint per treated acre may severely injure newly seeded grasses. Preplant applications – injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of ORACLE per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

Established Grass Stands

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species: bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass may be injured when treated with ORACLE at rates exceeding 1 pint per treated acre.

Weeds Controlled

ORACLE, when applied at specified rates, will control many annual and biennial weeds and provide control and suppression of many perennial weeds. (Refer to GENERAL WEED LIST on pages 7 and 8.)

Rates and Timings

Application rates and timing of ORACLE treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

Weed Type ¹ & Stage	Broadcast Rate Per Treated Acre	
	Amount of Formulated ORACLE pints	Equivalent lbs. ai
Annuals Small actively growing Established weed growth	1/4 to 1 1	1/8 to 1/2 1/2
Biennials ² Rosette diameter a) less than 3 inches b) 3 inches or greater c) bolting biennial	1/2 to 1 1 to 2 2	1/4 to 1/2 1/2 to 1 1
Perennials ² Suppression/Control	2	1

¹ For best results, treat Biennial weeds with ORACLE when they are in the rosette stage of growth. Retreatments may be made as needed; however, do not exceed a total of 2 applications of ORACLE® per year.

² Biennial and Perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

Tank Mix Treatments

To control grasses and additional broadleaf weeds, ORACLE may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate (Roundup®), paraquat (Gramoxone®), metsulfuron (Ally®) and others.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES, AND OTHER RESTRICTIONS.

ASPARAGUS

IMPORTANT

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Do not harvest prior to 24 hours after treatment.

Do not use in the Coachella Valley of California.

Multiple applications may be made per growing season. Do not exceed a total of 1 pint of ORACLE per treated acre per crop year.

Rates and Timings

Apply ORACLE to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

Weeds	Rate per Treated Acre
Mustard, Black Pigweed, Redroot (Carelessweed) Sowthistle, Annual ¹ Thistle, Canada Thistle, Russian	1/2 to 1 pt. (1/4-1/2 lb. ai)
¹ Bindweed, Field Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk	1 pt. (1/2 lb. ai)

¹ ORACLE may be applied in a tank mixture with either 2,4-D or Roundup Herbicide for improved control of Canadian thistle and field bindweed. READ AND FOLLOW 2,4-D AND ROUNDUP HERBICIDE PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

TURF AND LAWNS

For Use in General Farmstead (Non-Cropland) and Sod Farms

IMPORTANT

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

To avoid injury to newly seeded grasses, application of ORACLE should be delayed until after the second mowing. Furthermore, application rates in excess of 1 pint (1/2 lb. ai) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 1/4 pint (1/8 lb. ai) of ORACLE per treated acre on coarse textured (sandy-type) soils, or in excess of 1/2 pint (1/4 lb. ai) per treated acre on fine textured (clay-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of ORACLE have been activated in the soil by rain or irrigation.

Weeds Controlled

ORACLE, when applied at specified rates, will give control of many ANNUAL, BIENNIAL, and noted (*) PERENNIAL broadleaf weeds commonly found in turf. ORACLE will also give growth suppression of many other listed PERENNIAL broadleaf weeds and WOODY brush and vine species. (Refer to GENERAL WEED LIST on pages 7 and 8.)

Mixing and Application

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts. to 4 1/4 gals. per 1,000 sq. ft.) depending on density or height of weeds treated and on the type of equipment used.

Rates and Timings

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage And Type	ORACLE		
	Pints Per Treated Acre	lbs. ai Per Treated Acre	Teaspoons Per 1,000 sq. ft.
Annual			
Small, actively growing	1/2 to 1	1/4 to 1/2	1 to 2 1/4
Established weed growth	1 to 1 1/2	1/2 to 3/4	2 1/4 to 3 1/4
Biennial Rosette diameter			
less than 3 inches	1/2 to 1	1/4 to 1/2	1 to 2 1/4
3 inches or more	1 to 2	1/2 to 1	2 1/4 to 4 1/2
Perennials and Woody			
Brush and Vines	1 to 2	1/2 to 1	2 1/4 to 4 1/2

For best performance, apply when weeds are emerged and actively growing.

Retreatment may be made as needed, however, do not exceed a total of 2 applications of ORACLE per year. Do not exceed a total of 2 pints (1 lb. a.i.) ORACLE per treated acre during the growing season.

Tank Mix Treatments

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

Tank mix treatments of ORACLE may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product.

Apply 1/5 to 1/2 pint (1/10-1/4 lb. ai) of ORACLE per treated acre with 1/2 to 1 1/2 lbs. acid equivalent of 2,4-D, MCPA, or MCPP, or with 3/8 to 1/2 lb. ai of bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 2 pints (1 lb. ai) of ORACLE per treated acre during the growing season.

GRASS SEED CROPS: Grasses Grown for Seed such as Bermuda Grass, Bluegrass, Fescue and Ryegrass**IMPORTANT**

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

Refer to PASTURE, HAY, RANGELAND, AND GENERAL FARMSTEAD (Non-Cropland) section (pages 20-22) for possible grazing and feeding restrictions.

Do not use on bentgrass unless possible crop injury can be tolerated.

Weeds Controlled

ORACLE will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that ORACLE be applied in a tank mix with other herbicides.

Alfalfa ¹	Chamomile, Corn	Hemlock, Poison	Lettuce, Prickly
Bedstraw, Catchweed	Chickweed, Common	Knapweed, Russian ¹	Mayweed (Dogfennel)
Bindweed, Field	Chickweed, Mouseear	Knawel	Ragwort, Tansy
Buttercup, Corn	Clover	Knotweed, Prostrate	Sorrel, Red (Sheep Sorrel)
Buttercup, Creeping	Cockle, White	Kochia	Sowthistle, Annual
Buttercup, Western Field	Dock, Broadleaf	Ladysthumb	Starwort, Little
Catchfly, Nightflowering	Dock, Curly	Lambsquarters, Common	Thistle, Canada ¹

¹ Top growth only.

Rates and Timings

Apply 1/2 to 1 pint of ORACLE per treated acre on SEEDLING GRASS after the crop reaches the 3-5 leaf stage. Apply up to 2 pints of ORACLE on well-established Perennial grass. DO NOT APPLY AFTER THE GRASS SEED CROP BEGINS TO JOINT. For best performance, make applications when weeds are in the 2-4 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

Tank Mix Treatments

For control of grasses or additional broadleaf weeds, ORACLE may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions.

Broadcast Rate Per Treated Acre:

Apply 1/2 to 2 pints of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	1 - 4 pts. (0.5 - 2.0 lb ai/A) ¹
MCPA Amine	MCPA	4 lb/gal	1 - 2 pts. (0.5-1.0 lb ai/A) ¹
Buctril®	bromoxynil ²	2 lb/gal	1 - 2 pts.
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	1 3/4 - 4 pts.
Karmex®	diuron ²	80% DF	2 - 4 lbs.
Stinger®	clopyralid	3 lb/gal	1/4 - 1 pt.

¹ When using formulations other than 4 lb/gal, use pounds active/acre listed.

² Herbicides with the same common name and/or different formulations may be used.

Annual Grass Control

For suppression of ANNUAL GRASS WEEDS such as: Brome, Downy (Cheatgrass) Fescue, Rattail
Brome, Rippgut, Windgrass

Apply 2 pints of ORACLE per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

PREPLANT DIRECTIONS (POST HARVEST/FALLOW/CROP STUBBLE/SET-A-SIDE) FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS

IMPORTANT

Observe all precautions. Read and follow mixing and application instructions.

Weeds Controlled

ORACLE may be applied alone or in tank mix combinations with other herbicides registered for this use.

ORACLE can be applied either POST HARVEST in the fall, spring or summer during the FALLOW period or to CROP STUBBLE/SET-A-SIDE acres. ORACLE, when applied at the specified rates, will control many ANNUAL broadleaf weeds, see the WEEDS CONTROLLED section under small grains. In addition, ORACLE will control or suppress the following BIENNIAL and PERENNIAL broadleaf weeds:

Alfalfa ¹	Dandelion, Common ¹	Nightshade, Silver	Thistle, Milk
Artichoke, Jerusalem	Dock, Curly ¹	Redvine	Thistle, Musk
Bindweed, Field	Dogbane, Hemp	Smartweed, Swamp	Thistle, Plumeless
Bindweed, Hedge	Garlic, Wild ²	Sowthistle, Perennial ¹	Thistle, Scotch
Blueweed, Texas	Horsenettle, Carolina	Spurge, Leafy	Trumpetcreeper (Buckvine)
Bursage (Bur Ragweed),	Knapweed, Diffuse	Thistle, Bull	
Povertyweed, Lakeweed) ¹	Knapweed, Spotted	Thistle, Canada ²	

¹ These perennials may be controlled using ORACLE at rates lower than those recommended for other listed perennial weeds. (See RATES AND TIMINGS under this heading).

² See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control program for these weeds.

Rates and Timings

Apply ORACLE as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicides penetration of weed foliage. See CROP-PING RESTRICTIONS for specified interval between application and planting to prevent crop injury.

For best performance, make application when ANNUAL weeds are less than 6 inches tall, when BIENNIAL weeds are in the rosette stage and to PERENNIAL weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright PERENNIAL broadleaf weeds, such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets, after the effective period for ORACLE. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of ORACLE see the RATES AND TIMINGS section under the SMALL GRAINS heading for details.

Oracle Rates Per Treated Acre

Weed Type	Amount of Product Per Treated Acre
Annual	1/2 - 1 pt.
Biennial	1 - 2 pts.
Perennial Suppression Control - noted ⁽¹⁾ perennials	1 - 2 pts. 2 pts.

Retreatment may be made as needed, however, do not exceed a total of 2 applications of ORACLE per year.

Tank Mix Treatments

ORACLE may be tank mixed with one or more of , but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled and geographic or other restrictions.

ORACLE Broadcast Rate Per Treated Acre for Annual Weed Control:**Annual Weed Control**

Apply 1/4 to 1 pint of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Aatrex® 4L ¹	atrazine	4 lb/gal	0.5 - 6 pts.
Aatrex® Nine-O ¹	atrazine	90% DF	0.5 - 3.3 lbs.
Amber® ²	trisulfuron	75% DF	0.28 - 0.35 oz.
Ally® ²	metsulfuron-methyl	75% DF	0.1 oz.
Fallowmaster®	glyphosate + dicamba	2.0 lb/gal	22 - 44 fl. oz.
Gramoxone® Extra	paraquat	2.5 lb/gal	1.5 pts.
Kerb® ¹	pronamide	50-W	0.5 - 1.0 lb.
Landmaster ®BW	glyphosate + 2,4-D	2.4 lb/gal	27 - 54 fl oz.
Roundup®	glyphosate	3 lb/gal	8 - 48 fl oz.
Sencor® DF ¹	metribuzin	75% DF	0.5 - 1 lb.
Sencor® 4 ¹	metribuzin	4 lb/gal	0.75 - 1 1/2 pts.
2,4-D	2,4-D	4 lb/gal	1 -2 pts. (0.5 - 1 lb ai/A) ³

¹ Tank mixes of ORACLE with these products may be subject to special restrictions. See the Product Label of the tank mix partner for intended use rates, restrictions and other precautions.

² When tank mixing with sulfonylurea herbicides refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1 - 2 quarts/100 gallons of spray or not more than 0.25 - 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea resistant weeds may not be controlled by tank mixes of ORACLE and a sulfonylurea. Refer to the ORACLE tank mix section for alternative tank mixes.

³ When using formulations other than 4 lb/gal, use pounds active/acre listed.

ORACLE Broadcast Rate Per Treated Acre for Biennial and Perennial Weed Control:

Apply 1 to 2 pints of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	2 - 4 pts.
2,4-D	2,4-D	4 lb/gal	2 - 6 pts.(1.0 - 3 lb ai/A) ¹
Landmaster ®BW	glyphosate + 2,4-D	2.4 lb/gal	54 fl oz.
Roundup®	glyphosate	3 lb/gal	1 - 5 qts.
Tordon® 22K	picloram	2 lb/gal	1/2 - 1 pt.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

Special Tank Mix Treatments

For suppression of perennial weeds, apply 1/2 - 1 pint of ORACLE with 8 to 16 fluid ounces of Roundup Herbicide or Roundup RT per treated acre.

For wild garlic control, apply 1 pint ORACLE with 3 pints of 2,4-D LV Ester (4 lb/gal) per treated acre. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use ORACLE, ORACLE plus Curtail®, or ORACLE plus Roundup® Herbicide tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 1 pint ORACLE with 1/2 to 1 lb Kerb® 50W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply ORACLE plus Landmaster® BW or Fallowmaster® Herbicide to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle. Use 1/8 - 1/4 pint ORACLE plus 22 - 54 fluid ounces of Landmaster® BW or Fallowmaster® Herbicide for annual weed control or 1/4 - 1/2 pint ORACLE plus 22 - 54 fluid ounces of Landmaster® BW or Fallowmaster® Herbicide for perennial weed suppression.

Cropping Restrictions

The following recommendations are based on Oracle use rates up to 2 pints per treated acre.

CORN, SORGHUM and SOYBEANS may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of ORACLE per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of ORACLE per treated acre. Exclude days when ground is frozen.

WHEAT may be planted in the fall or spring following applications. Also, spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per pint of ORACLE per treated acre or 1.25 days per 1 ounce. Moisture is essential for ORACLE degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of ORACLE per treated acre or 3 days per ounce. Moisture is essential for ORACLE degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION ONLY) FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND WASHINGTON

IMPORTANT

Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

Do not treat subirrigated cropland or areas where the soil remains saturated with water throughout the year.

Make only one application of ORACLE per year.

Spot Application is defined as an area no greater than 1000 ft. sq. per acre.

Weeds Controlled

ORACLE, when applied at specified rates, will control or suppress many broadleaf weeds including:

Bindweed, Field
Dock, Broadleaf (Bitterdock)
Dock, Curly
Knapweed, Black
Knapweed, Russian
Ragwort, Tansy
Spurge, Leafy
Thistle, Canada

Rates and Timings

ORACLE may be applied at any time following a crop harvest to stubble fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost.

Apply 1 qt. (1 lb. ai) of ORACLE per treated acre. Application may be made up to one month prior to the planting of wheat.

NOTE: Do not use unless injury to wheat or rotated barley will be acceptable.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugarbeets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

WIPER APPLICATION USES

IMPORTANT: Observe all PRECAUTIONS on pages 3 to 5. Read and follow MIXING AND APPLICATION instructions on page 4.

ORACLE may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part ORACLE to 1 part water. Do not contact desirable vegetation with herbicide solution. Wiper application should only be made to crops (including pastures) and non-cropland areas described in this label with the exception of Grain Sorghum (Milo).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a well-ventilated area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

(Nonrefillable container 5 gallons or less): Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. If not, triple rinse emptied container and offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix

STORAGE AND DISPOSAL *(continued)*

tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(Refillable containers up to 250 gallons): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning the container before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC: 1-800-424-9300

NOTICE OF WARRANTY AND DISCLAIMER

Seller warrants that at the time of delivery the product in this container conforms to its chemical description contained hereon and is reasonably fit for its intended purpose under normal conditions of use. This is the only warranty made on this product. To the extent consistent with applicable law, seller expressly disclaims any implied warranties of merchantability or fitness for any particular purpose and, except as set forth above, any other express or implied warranties. Any damages arising from breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid for this product by Buyer, and shall not include incidental or consequential damages such as, but not limited to, loss of profits or values. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the Seller. To the extent consistent with applicable law, in no case shall Seller be liable for the consequential, special or indirect damages resulting from the use or han-

dling of this product. All such risks shall be assumed by the Buyer. Buyer acknowledges the use of its own independent skill and expertise in the selection and use of the product and does not rely on any oral or written statements or representations.

REGISTERED TRADEMARKS

Aatrex[®], Amber[®], Beacon[®], Bicep[®] II Magnum, Concep[®], Dual[®] II Magnum, Evik[®], Gramoxone[®], and Princep[®] are trademarks of Syngenta Group Company.

Karmex[®], Accent[®], Ally[®], Express[®], and Harmony[®] are trademarks of E.I. duPont de Nemours & Company, Inc.

Bronate[®], Buctril[®], and Sencor[®] are trademarks of Bayer CropScience.

Curtail[®], Garlon[®], Stinger[®], Surpass[®] and Tordon[®] are trademarks of Dow AgroScience.

Clarity[®], Frontier[®], Guardsman[®], Marksman[®] and Prowl[®] are trademarks of BASF Corporation.

Bullet[®], Fallow Master[®], Harness[®], Landmaster[®], Lariat[®], Lasso[®], Permit[®], Roundup[®], and Roundup RT[®] are trademarks of Monsanto Company.

Eradicane[®] is a Trademark of Gowan Co.

Sutan[®] is a Trademark of TRI-AG Inc.

Asulox[®] is a Trademark of United Phosphorus.

Registered: January 13, 2006

Amended: August 25, 2009

Revised by Notification: Storage and Disposal Statements/Add-back Use Directions Preplant to Cotton: April 5, 2010

Updated: (Address Change) May 24, 2017

NOTES

6.2500



6.2500



ORACLE® Dicamba Agricultural Herbicide

For weed control in corn, cotton, sorghum, small grains, pasture, hay, rangeland, general farmstead (non-cropland), rights-of-way, public utility and industrial areas, fallow, sugarcane, asparagus, turf and grass seed crops.

Active Ingredient:

Dimethylamine salt of dicamba
(3, 6-dichloro-*o*-anisic acid)* 49.77%

Inert Ingredients: 50.23%

Total: 100.00%

*This product contains 41.35% 3, 6-dichloro-*o*-anisic acid (dicamba) or 4 pounds per gallon (480 g/L).

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for complete *Precautionary Statements, Directions for Use and Conditions of Sale and Warranty.*

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed.

FIRST AID (Substituted Benzoic acid)

If swallowed:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Remove person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER (Substituted Benzoic acid)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: 1-(866)-359-5660

Personal Protective Equipment (PPE)

Some material that is chemical-resistant to this product is natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks
- Chemical-resistant gloves (except for applicators using groundboom equipment, pilots and flaggers)
- Protective eyewear

See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as directed on label.

EPA Reg. No. 93182-10

EPA Est. No. 2217-KS-1^P

EPA Est. No. 44616-MO-2^H

EPA Est. No.: 19713-TN-8

(Superscript designates first letter of lot number on jug.)

Oracle® Registered Trademark of Gharda Chemicals International, Inc.

Net Contents: 2.5 gal.

Distributed by:



Gharda Chemicals International Inc.

760 Newtown-Yardley Rd., Suite 110
Newtown, PA 18940
1-215-968-9474