



## L-Arginine Detection Report

**Protocol No.:** DET-PRO-016

**Name of the product:** Lumineux Oral Essentials clean and fresh

---

**Approved by:** *Nour Abochama* Date: 2019-03-01  
Nour Abochama  
Director QA/QC, American Testing Lab Inc.

**Approved by:** *Kouros Maddahi DDS* Date: 3/1/2019  
Kouros Maddahi DDS  
Founder, Oral Essentials Inc.

## Table of Contents

1	Objective: _____	3
2	Scope: _____	3
3	Responsibility: _____	3
4	Materials, Equipments and documents: _____	3
4.1	List of Equipment: _____	3
5	Analytical Method _____	4
5.1	Standers and Reagents: _____	4
5.2	Solutions: _____	4
5.3	standard: _____	4
5.4	Sample: _____	4
5.5	Chromatographic Conditions: _____	5
5.6	Injection Run: _____	5
5.7	System Suitability: _____	5
5.8	Calculations: _____	5
6	Protocol _____	6
8	Results: _____	6
9	Conclusion: _____	6

**1 Objective:**

To determine the effect of Lumineux Oral Essentials clean and fresh on L-Arginine concentrations.

**2 Scope:**

A 3T3 clone A31 Cell Line from mouse was exposed to a 5000 µg/ml solution of test product for 24 hours. A determination of L-Arginine concentration was performed using HPLC method.

**3 Responsibility:**

The persons responsible for the process will perform the and record the information. The responsible person will supervise the study, verify the completion of the records and write the report. Quality Assurance will review and approve the Report.

Name of the Personnel	Designation	Signature
__Nour Abochama__	__Director QA/QC__	<i>Nour Abochama</i>
__Reem Samman__	__Chemist__	<i>ras</i>

**4 Materials, Equipments and documents:**

List with Equipment No. and all Standard Operation Procedures (SOPs) for the normal processes under test.

**4.1 List of Equipment:**

Name of Equipment	Equipment No.
HPLC with UV detector	LAB-2426
Balance	LAB-2233
Pipette	LAB-2422
Incubator	LAB-2223
HPLC Column, Luna C18, 250 x 4.6 mm, 5µm	LAB-0175

Complied By: ras Date: 2019-02-28Reviewed By: Nour Abochama Date: 2019-02-28

## 5 Analytical Method

### 5.1 Standers and Reagents:

- 5.1.1 L-Arginine Reference Standard (or equivalent);
- 5.1.2 Phosphoric Acid H<sub>3</sub>PO<sub>4</sub>, ACS grade (or equivalent);
- 5.1.3 Acetonitrile HPLC grade (or equivalent)
- 5.1.4 Purified water USP grade (or equivalent);

### 5.2 Solutions:

- 5.2.1 *Mobile Phase:* 90% of Line A: 10% of Line B where: Line A: 0.1% Phosphoric Acid, Line B: Acetonitrile.

### 5.3 standard:

- 5.3.1 Accurately weight and transfer about 13 mg ± 1.0 mg of L-Arginine Reference Standard into 10 mL volumetric flask.
- 5.3.2 Dissolve in about to ¾ of volume with purified water.
- 5.3.3 Complete to volume with mobile phase.
- 5.3.4 Mix well.

### 5.4 Sample:

- 5.4.1 Transfer 1 ml of test sample into 10 mL volumetric flask.
- 5.4.2 Dissolve in about to ¾ of volume with purified water.
- 5.4.3 Swirl to homogenize the solution.
- 5.4.4 Let the solution to cool at room temperature.
- 5.4.5 Complete to volume with purified water and mix well.
- 5.4.6 Immediately filter through a 0.45 µm nylon filter and transfer into HPLC vial discarding the first 2 mL.

**5.5 Chromatographic Conditions:**

- 5.5.1 Injection Volume: 5  $\mu$ L;
- 5.5.2 Wavelength: 195 nm;
- 5.5.3 Mobile Phase: As per section 5.2.1 of this method;
- 5.5.4 Flow Rate: 0.5 mL/min;
- 5.5.5 Expected RT: 3.7 min;
- 5.5.6 Run time: 15 min.

**5.6 Injection Run:**

- 5.6.1 1 inj Mobile Phase BLK;
- 5.6.2 5 inj working standard STD;
- 5.6.3 1 inj samples;
- 5.6.4 1 inj control (working standard) QSTD

**5.7 System Suitability:**

- 5.7.1 From the 6 injections of standard, area RSD should not be more than 5.0%.
- 5.7.2 Recovery between standard control and mean area of standard should be within 95.0%-105.0%.

**5.8 Calculations:**

- 5.8.1 Calculate against the mean area of first 5 injection of standard.

## 5.8.1.1 Recovery standard

$$\% \text{ Recovery} = (\text{Area}_{\text{qstd}} / \text{Mean Area}_{\text{std}}) \times 100$$

Where:

- Area<sub>qstd</sub> = Area of peak of interest in standard or Control.
- Mean area<sub>std</sub> = Mean area of peak of interest in standard

## 5.8.1.2 Sample Calculation

$$\% \text{ of Arginine in Products} = (\text{Area}_{\text{spl}} / \text{Mean Area}_{\text{std}}) \times (\text{C}_{\text{std}} / \text{W}_{\text{spl}}) \times 100 \text{ mL} \times 100$$

Where:

- Area<sub>spl</sub> = Area of Arginine in Sample preparation
- Mean Area<sub>std</sub> = Mean Area of Arginine in working standard preparation
- C<sub>std</sub> = Concentration of standard in working standard (mg/mL)

$W_{spl}$  = Weight of sample (mg)

 Complied By: ras Date: 2019-02-28

 Reviewed By: Nour Abochama Date: 2019-02-28

## 6 Protocol

- 6.1 The test item is prepared y adding 10% (v/v) FCS and 1% antibiotics (v/v) to the artificial Sylvia  
 6.2 Prepare 5000 µg/mL solution of Lumineux Oral Essentials clean and fresh . Expose the test  
 6.3 item to the product solution for 24, 36, 48, 60, 72, 84 and 96 hours then prepare as described in section 5.4.

## 7 Approval:

- 7.1 Submit the Document to QA for review and approval.

## 8 Results:

Exposure Time	Blank (µg/ml)	Test Product (µg/ml)	Water (µg/ml)	Ref. Product (µg/ml)
0 hours	0.00	0.2116	0.211	0.223
24 hours	0.00	0.1692	0.211	0.222
36 hours	0.00	0.1269	0.210	0.223
48 hours	0.00	0.1058	0.211	0.221
60 hours	0.00	0.0846	0.211	0.02
72 hours	0.00	0.0423	0.211	0.02
84 hours	0.00	0.0212	0.211	0.02
96 hours	0.00	0.0211	0.211	0.02

## 9 Conclusion:

The Arginine-specific gingipain (Rgp) concentration have been 90% decreased in test item after 96 hours of exposure.

The solution at 5000 µg/mL is prepared form Lumineux Oral Essentials clean and fresh Lot# 17K13B2

 Approved By: Nour Abochama  
 Nour Abochama, M. Sc.,

 Date: 2019-03-01