

Usage Instructions

Suggested Use for:

Most Peptides

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Inject 10iu Sub-Q, Monday - Friday, upon waking
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

DSIP

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Inject 20iu 1 (one) hour prior to sleeping
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

PT-141

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Inject 20iu 1 (one) hour prior to sexual activity
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

Retratutide

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Weeks 1-4 Inject 50iu (2.5mg) Sub-Q weekly
- 3 - Weeks 5-8 Inject 100iu (5mg) Sub-Q weekly
- 4 - Titrate upwards weekly at 50iu increments, as tolerated, (not to exceed 200iu)
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

Selank

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Inject 10iu Monday - Friday
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

Semaglutide

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Weeks 1-4 Inject 10iu (.25mg) Sub-Q weekly
- 3 - Weeks 5-8 Inject 20iu (.5mg) Sub-Q weekly
- 4 - Titrate upwards weekly at 10iu increments, as tolerated, (not to exceed 50iu)
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting

Suggested Use for:

Tirzepatide

- 1 - Reconstitute with 2ml of bacteriostatic water
- 2 - Weeks 1-4 Inject 100iu (2.5mg) Sub-Q weekly
- 3 - Weeks 5-8 Inject 200iu (5mg) Sub-Q weekly
- 4 - Titrate upwards weekly at 100iu increments, as tolerated, (not to exceed 400iu)
- 3 - Refrigerate at 2° - 8° Celsius (36° - 46° Fahrenheit)
- 4 - Use within 30 days after reconstituting
- 5 - Refrigerate after reconstituting