



IMPORTANT INSTRUCTIONS

How to chew Nicotex Gum

The method of chewing the gum is NOT the same as that of an ordinary chewing gum. Use the following steps to ensure maximum benefit from the gum.







STEP 1

Chew it slowly till you get the taste of Nicotine

Place and Rest between your cheek and gums

1. What is Nicotex?

Nicotex is a gum containing nicotine that helps people who smoke or chew tobacco and gutka quit the habit. It works on the proven principle of "Nicotine Replacement Therapy (NRT)".

Nicotex does not contain tar and other harmful chemicals which are present in cigarettes, tobacco and gutka.

2. What is Nicotine?

Nicotine is a chemical present in tobacco leaves. When you smoke or chew tobacco, nicotine enters the blood and goes to the brain. It activates the part of your brain which produces a feeling of pleasure and happiness. After sometime, as the nicotine levels drop, the feel-good effect reduces, and your body starts craving for the next cigarette or tobacco. Thus, you get addicted to smoking cigarettes or chewing tobacco.

3. How does Nicotex work?

Each piece of Nicotex provides the nicotine that you normally get from cigarettes or chewing tobacco. The level of nicotine is lower than that in cigarettes or chewing tobacco. Nicotex allows your body to gradually adjust to having less nicotine until you no longer need any and thus, helps you quit smoking or chewing tobacco.

4. What is the indication for the use of Nicotex?

To reduce withdrawal symptoms including nicotine craving associated with quitting smoking/chewing tobacco and gutka containing tobacco.

5. Which is the right strength for you?Nicotex is available in two strengths, 2 mg and 4 mg. 2 mg is appropriate for those who smoke 20 cigarettes a day

4 mg is appropriate for those who smoke more than 20 cigarettes a day1.

6. What is the dosage and administration?

4 mg: As advised by the doctor.


2 mg: Consume a gum whenever you get an urge to smoke. Over a 12 week period, Nicotex allows your body to gradually adjust to have less nicotine.



Cipla **nicotex**®



Below dosage schedule is recommended:



Do not exceed more than 24 gums per day. Reduce 1 gum per week Recommended therapy is for 12 weeks.

NoteDo not eat or drink or use removable dentures with the gum in the mouth. Avoid having any drinks for 15 minutes before the $\,$ gum is used. The side effects which might be seen with Nicotex will be the same as nicotine given by any other route. The side effects are usually dose dependent and may not be experienced by all. Common side effects such as headache. nausea, hiccups, stomach discomfort, sore mouth or throat, jaw muscle ache and sleep disturbance may be seen particularly during the first few weeks.

Consult the Doctor

- \bullet If you are taking any other medicines especially, those containing Theophylline, Tacrine, Clozapine, Ropinirole or non-nicotine drugs to stop smoking. If you have had previous reactions to gums or other nicotine containing products. If you experience anything that is unusual after the use of Nicotex.
- \bullet If you have other medical conditions like uncontrolled high blood pressure, stroke, stomach ulcer, kidney or liver disease, heart or circulation problems, diabetes, allergic reactions, severe sore throat, inflammation of mouth, overactive thyroid, irritable bowel syndrome or adrenal gland cancer
- \bullet If you are pregnant or breast feeding, on sodium restricted diet, or need to use the gum for more than 12 weeks

Nicotex Overdose

Material No.: 21090126 Version: 01 Item: Leaflet Co-ordinator: Sibanti Artist: Chetan Date: 14/07/2021

If you have used more gums than recommended or in case a child has chewed or swallowed some, then contact a doctor immediately. The general symptoms of nicotine overdose include headache, sickness, diarrhoea and stomach pain.

Do not use Nicotex gum:

• If you are below 18 years of age • If you are a non-user of tobacco

Storage

STORE IN A COOL, DRY PLACE. PROTECT

FROM DIRECT HEAT AND SUNLIGHT.

Keep out of reach of children. Don't use after expiry date mentioned on the box.

cipla.health@cipla.com www.nicotex.in Call 1800 120 143143

No.1 Doctor Recommended Brand in India* * In 'NRT' category, IMS Feb 21 MAT. # J Behav Med. 1982; 5(3):43-51 (for 2mg); Arch Intern Med. 1989; 149:1818-21 (for 4mg) http://www.mayo.edu/issearch/documents/medication-handout-2015-02-pdf/doc-20140182

> We believe you can.

PACKAGING DEVELOPMENT Product Name: Nicotex Tin leaflet

Colours: BLUE WOOL TEST VALUE 5-8 (LIGHT FASTENING DATA) Pantone 363 U Pantone 2945 U Black						INK: Oil based Ink from DIC OR MICRO			
Design: Folded Refe				5315	Software: Illustrator CC				
Fonts: Links: NA									
Actual Size: 70 x 200 mm	Size after folding: 70 x 17 mm	Pharmacode/2D Code:			Grain Dire	ection : Parallel to length			Screen : #
Material: 60 GSM Maplitho Paper					Varnish: Artwork F				
Path: D:\Nicotex Leaflet Tin Container\21090126 Nicotex Leaflet Tin Container Common.ai									
• Instructions / Remark:			Checked by	Artist		Cordinator	file loaded in Server		Section Head
Any deviation must be brought to the notice of packaging development co-ordinator immediately. For any darification, please contact packaging development co-ordinator immediately. NO CHANGES IN ARTWORK SHOULD BE DONE BY THE PRINTER The printer should verify the e-proof against the approved artwork before submitting for approval and the e-proof should have printer details.			Pharma Co 2D Code QR Code Bar Code Artwork Spell check						