

Drug Facts

Active ingredient (per each tablet)

Purpose

Menthol 2.5 mg......Oral pain reliever

Indications for the temporary relief of pain associated with canker sores.

Warnings

Do not use this product for more than 7 days unless directed by a dentist or doctor. Stop use and ask a dentist or doctor if •sore mouth symptoms do not improve in 7 days •irritation, pain or redness worsens •swelling, rash or fever develops Do not exceed recommended dosage.

If pregnant or breast-feeding or taking prescription drugs, ask a health care professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions •adults and children 5 years of age and older, apply up to 3 tablets a day, as needed. Place the tablet on a clean, dry finger with the white side up. Place the white side on the sore and hold in place for 10 seconds. If sore is difficult to reach (in the fold between the cheek and gum or near the teeth or lip) break the tablet along the score and use half. Some discomfort may occur during the first few minutes, but will quickly subside, followed by hours of soothing relief. Within 30 minutes the tablet forms a clear, gel-like bandage that seals and protects the sore for hours before dissolving. Do not remove the tablet before it dissolves. In case of discomfort, the tablet may be removed by gently peeling the tablet from the sides while washing with warm water. Do not use any instrument to remove the tablet.

•children under 5 years, ask a doctor

Other information store in a cool, dry place.

Inactive ingredients annato, carbomer homopolymer type a, citrus oil, hydroxypropl cellulose, magnesium chloride, povidone K30, povidone K90, silicon dioxide, xylitol

Questions or comments? 833-363-2763 or consumeraffairs@doc-brands.com