

Cetuximab

Cat. No.:	HY-P9905
CAS No.:	205923-56-4
Target:	EGFR; Radionuclide-Drug Conjugates (RDCs)
Pathway:	JAK/STAT Signaling; Protein Tyrosine Kinase/RTK; Antibody-drug Conjugate/ADC Related
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.

BIOLOGICAL ACTIVITY

Description	Cetuximab (C225) is a human IgG1 monoclonal antibody that inhibits epidermal growth factor receptor (EGFR), with a K_d of 0.201 nM for EGFR by SPR. Cetuximab has potent antitumor activity ^[1] .		
IC ₅₀ & Target	EGFR 0.147 nM (K_d , Fixed A431 cells)	RDC Antibody	Soluble EGFR 0.201 nM (K_d)
In Vitro	Cetuximab (C225) is a monoclonal antibody that inhibits epidermal growth factor receptor (EGFR), with a K_d of 0.201 nM for soluble EGFR by SPR. Cetuximab also exhibits a K_d of 0.147 nM for EGFR in fixed A431 cells by ELISA ^[1] . Cetuximab (C225; 30 nM) time-dependently inhibits the proliferation of SCC-1, SCC-11B, SCC-38, and SCC-13Y cells after treatment for 8 d. Cetuximab (30 nM) causes G ₀ /G ₁ arrest, induces apoptosis, and reduces Rb, p27 ^{KIP1} , Bcl-2, and Bax expression in SCC-13Y cells. Cetuximab (30 nM) also enhances radiosensitivity and increases radiation-induced apoptosis in SCC-13Y cells ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.		
In Vivo	Cetuximab (1 mg/injection) has effect on the tumour volume but the effect is more pronounced on UT-SCC-14 xenografts. In UT-SCC-14 xenografts, Cetuximab significantly reduces the expression of EGFR, pEGFR and Ki67. Cetuximab significantly decreases the expression in the MCT1 and GLUT1 cells but differences are more pronounced in UT-SCC-14 xenografts ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.		

CUSTOMER VALIDATION

- Cell. 2023 Dec 7;186(25):5606-5619.e24.
- Cell Res. 2022 Oct 14.
- Cell Res. 2020 Dec;30(12):1063-1077.
- Nat Nanotechnol. 2024 Oct 28.
- Mol Cancer. 2021 Jan 18;20(1):17.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Goldstein NI, et al. Biological efficacy of a chimeric antibody to the epidermal growth factor receptor in a human tumor xenograft model. Clin Cancer Res. 1995 Nov;1(11):1311-8.
- [2]. Gustafsson H, et al. EPR Oximetry of Cetuximab-Treated Head-and-Neck Tumours in a Mouse Model. Cell Biochem Biophys. 2017 Jul 29.
- [3]. Huang SM, et al. Epidermal growth factor receptor blockade with C225 modulates proliferation, apoptosis, and radiosensitivity in squamous cell carcinomas of the head and neck. Cancer Res. 1999 Apr 15;59(8):1935-40.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA