





# Technical Bulletin for: San Jose Scale

*Quadraspidiotus perniciosus* (Comstock) Hemiptera, *Diaspididae* • QUAPER



<b>DISTRIBUTION</b>	Found in most areas of the world.
<b>HOSTS</b>	Apples, peaches, pears, plums and Rubus.
<b>DESCRIPTION</b>	
<b>ADULT</b>	About the size of a pinhead; gray, circular, and slightly raised. It often has a ringed appearance from molting. Females can reach 38 mm in diameter. The soft body underneath is yellow with no wings or legs.
<b>NYMPH</b>	Black, less than 1 mm in diameter.
<b>EGGS</b>	Female San Jose scale do not produce eggs, nymphs are born live.
<b>LIFE HISTORY</b>	Overwinters in nymph stage. Increased scale problems can be expected after mild winters. After developing through larval and pupal stages, the males mature and back out from their scales about 4 to 6 weeks after birth. Adult males fly for only a few days and are capable of mating immediately with the females, which remain under their scales. Each female produces several hundred live crawlers over a 6-week period. Timing of the different stages varies from year to year, depending on temperatures. Usually, crawlers of the first generation appear in early June and may continue to be produced until early August. The young crawlers can be carried to other trees by the wind, on the feet of birds, on the clothing of farm workers or on orchard equipment. Females of the first generation mature in late July, and second generation crawlers appear in August. Second generation crawlers continue to be produced until October or November.

## MONITORING INFORMATION

<b>LURE ACTIVE INGREDIENT, SUBSTRATE &amp; FIELD LIFE</b>	SJS Pheromone and Isogranyl propionate in a Red Rubber Septum. Lure Longevity: four (4) weeks.	
<b>TRAP TO USE</b>	Scale Card	
<b>MONITORING STRATEGY</b>	Use a minimum of 3-4 traps, irrespective of orchard size. Place traps in orchards at the pink stage of apple fruit bud development. Position traps on the northern or eastern side of the tree at a height of 6 to 7 feet. Check traps frequently (every day or 2) until the first male scale is caught. Check with Cooperative Extension or Master Gardener for local information and recommendations.	
<b>CULTURAL &amp; PHYSICAL CONTROLS</b>	Good pruning practices can help reduce scale problems. Pruning out infested limbs and water sprouts can help reduce spring and summer scale populations as well as open up the canopy to allow better spray coverage in tree tops where scales are often concentrated.	

**Alpha Scents, Inc.**  
insect monitoring systems

Alpha Scents, Inc., 360 S. Sequoia Pkwy. Canby, OR 97013  
TEL: 503-342-8611 • FAX: 314-271-7297 • sales@alphascents.com • www.alphascents.com