BABY BABY

COLICKY HIGHNEEDS

HAVE NORMAL SLEEPING & EATING PATTERNS, BUT OFTEN WAKE FROM **COLIC PAIN**

SLEEP

IRREGULAR SLEEPING & EATING PATTERNS. WAKE & FEED **FREQUENTLY**

NO NEGATIVE REACTION TO NOISE OR PEOPLE

NOISE

HAVE INTENSE NEGATIVE REACTIONS TO

CRYING FOR MORE THAN 3 HOURS A DAY.

CRYING EPISODES LAST LONGER & SEEM

TO HAPPEN FOR NO REASON

CRYING

CRY LESS THAN COLIC BABIES, BUT MORE THAN A **NORMAL BABY**

NOISE & COMMOTION

CRYING TYPICALLY LASTS FOR 3 MONTHS, BUT CAN LAST UP TO 6 MONTHS

DURATION

CRYING LASTS THROUGHOUT FIRST FEW YEARS

Colic Symptoms

GRIMMACED FACE

DIFFICULT TO SOOTHE

RED FACE

ARCHED BACK

GAS



EXCESSIVE CRYING

HIGH PITCHED PIERCING SOUND

CLENCHED FISTS

KNEES DRAWN TO CHEST

High Needs Symptoms

DIFFICULT

NEGATIVE REACTION TO NOISE

FUSSY

WAKE FREQUENTLY

DEMANDING



IRREGULAR SLEEPING & EATING PATTERNS

HAVE A HARD TIME ADJUSTING TO NEW ENVIRONMENT

HYPERACTIVE

FEARFUL OF NEW PEOPLE

AVERAGE CRYING EPISODE OF A COLICKY **BABY**





AVERAGE CRYING EPISODE OF A HIGH NEEDS





































OF BABIES BORN (OR 2 OUT 10) HAVE COLIC

OF BABIES BORN (OR 1 OUT 10) ARE HIGH **NEEDS**