

# GENERATIONS 20/20 LIFEGUARD MILK REPLACER NT 1600

PRODUCT CODE: 3000AUTO

FORM: Powder

MEDICATION: Oxytetracycline 1600 g/ton Neomycin 1600 g/ton

## **MIXING DIRECTIONS:**

Measure 2 quarts of clean, warm water (110°F) into a clean bucket. Sprinkle 10 oz (1 cup)\* of milk replacer onto the water surface and mix thoroughly. For larger batches, mix 1-1/4 lbs of milk replacer powder per gallon of warm water. Always make reconstituted milk replacer as needed for immediate feeding.

## **FEEDING DIRECTIONS:**

In any calf-raising program, it is important that calves receive colostrum for the first 3 days. Starting the 4<sup>th</sup> day, feed milk replacer according to the following directions:

 $\underline{\textit{Light Breed Calves}}$  - 1-1/2 to 2 quarts liquid milk replacer twice per day.

<u>Heavy Breed Calves</u> - 2 to 2-1/2 quarts liquid milk replacer twice per day.

Feed at a rate to provide 10 mg of Oxytetracycline and 10 mg of Neomycin per pound of bodyweight daily. Feed continuously for 7 to 14 days.

## **FEATURES & BENEFITS:**

- Medicated for the treatment and control of bacterial enteritis and for the treatment of bacterial pneumonia.
- Formulated with LifeGuard®, a proprietary blend of prebiotics and essential oils designed to support digestive health and immune function.
- Contains chelated and organic trace minerals to maximize absorption.
- Specifically formulated for auto feeder mixing to reduce build up.
- Formulated with all-milk proteins for increased availability.

# **GUARANTEED ANALYSIS:**

20.00%
20.00%
0.15%
0.75%
1.25%
0.70%
30,000 IU/lb.
10,000 IU/lb.
200 IU/lb.

# **INGREDIENTS:**

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHT), Yeast Extract, Dried Skimmed Milk, Lecithin, Dried Milk Protein, Dicalcium Phosphate, Calcium Carbonate, Magnesium Sulfate, l-Lysine, dl-Methionine, Vitamin A Acetate, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Copper Proteinate, Manganese Proteinate, Zinc Proteinate, Iron Proteinate, Selenium Yeast, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Ascorbic Acid, Biotin, Riboflavin, d-Calcium Pantothenate, Niacinamide, Choline Chloride, Menadione Sodium Bisulfite Complex (Source of Vitamin K<sub>3</sub>), Folic Acid, Pyridoxine Hydrochloride, Calcium Iodate, Zinc Sulfate, Ferrous Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Sulfate, Sodium Silico Aluminate, Polyoxyethylene Glycol (400) Mono and Dioleates, Natural and Artificial Flavor.

\*The measuring cup provided in the bag holds approximately 10 ounces of milk replacer powder. For greater accuracy, a suitable scale is recommended.

#### WARNING:

If symptoms persist after using for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 28 hours beyond remission of disease symptoms. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Withdraw 5 days before slaughter.

**CAUTION:** Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

# PRODUCT REFERENCE