



=medicated=

STARTER GROWER

For Chicks From Hatch Until Laying Age



For the prevention of coccidiosis where immunity to coccidiosis is not desired in broilers and replacement pullets.
For the prevention of coccidiosis in turkeys.

FEEDING DIRECTIONS

Feed free choice to pullet chicks from hatch to laying age.

Feed free choice to broiler chicks from hatch to harvest. May feed to turkeys at about 20 weeks of age as a finishing feed. Provide free choice clean, fresh water at all times.

ACTIVE DRUG INGREDIENT: Amprolium 113.5 g/ton (0.0125%)

WARNING: Use as the sole source of Amprolium.

SPECIAL CONSIDERATION: Do not use with feeds containing bentonite

GUARANTEED ANALYSIS:

Crude Protein (Min.)	18.00%
Lysine (Min.)	1.00%
Methionine (Min.)	0.40%
Crude Fat (Min.)	3.00%
Crude Fiber (Max.)	5.00%
Calcium (Ca) (Min.)	0.75%
Calcium (Ca) (Max.)	1.25%
Phosphorus (P) (Min.)	0.70%
Salt (NaCl) (Min.)	0.10%
Salt (NaCl) (Max.)	0.60%

18%

Higher Protein
to Support Healthy,
Happy Chicks



Medicated for
the PREVENTION
of COCCIDIOSIS



Complete, Balanced
Nutrition

Available in a 5 lb pouch or a 25 lb bag

INGREDIENTS:

Grain Products, Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Sodium Sesquicarbonate, DL-Methionine, L-Lysine, Vitamin A Supplement, Vitamin D Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Vitamin B-12 Supplement, Riboflavin, d-Calcium Pantothenate, Folic Acid, Pyridoxine Hydrochloride, Menadione Sodium Bisulfite Complex (source of Vitamin K activity), Thiamine Mononitrate, Biotin, and Ethoxyquin (a preservative), Copper Sulfate, Cobalt Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate, Ethylenediamine Dihydroiodide, and Sodium Selenite.