

Vitamin D May Play Key Role in Macular Degeneration

Mon, 08/31/2015 - 10:10am by University of Buffalo



rstock

Vitamin D has been studied extensively in relation to bone health as well as cancer. Now, a team led by a researcher at the University at Buffalo has discovered that vitamin D may play a significant role in eye health, specifically in the possible prevention of age-related

macular degeneration, or AMD, among women who are more genetically prone to developing the sight-damaging disease.

In a paper published in *JAMA Ophthalmology* online, Amy Millen, associate professor of epidemiology and environmental health in UB's School of Public Health and Health Professions, and her team found that women who are deficient in vitamin D and have a specific high-risk genotype are 6.7 times more likely to develop AMD than women with sufficient vitamin D status and no high risk genotype.

"Most people have heard that you should eat carrots to help your vision. However, there appear to be many other ways that adequate nutrition can support eye health. Having adequate vitamin D status may be one of them," said Millen, PhD, the study's lead author. "This is not a study that can, alone, prove a causal association, but it does suggest that if you're at high genetic risk for AMD, having a sufficient vitamin D status might help reduce your risk."

"To our knowledge, this is the first study that's looked at the interaction between genetic risk and vitamin D status in the context of age-related eye disease," adds Millen.

Macular degeneration is characterized by the deterioration of the macula, a small part of the central retina where the eye's photoreceptors (rods and cones) are most highly concentrated. The leading cause of legal blindness, macular degeneration affects more than 10 million Americans — more than cataracts and glaucoma combined — according to the American Macular Degeneration Foundation. The disease affects a person's central vision, which is needed for common tasks such as reading and driving. The effect is similar to that of a rain drop on the center of a camera lens.

Researchers analyzed data compiled on 1,230 women ages 54 to 74 who participated in the Carotenoids in Age-related Eye Disease Study (CAREDS), which is an ancillary study of the Women's Health Initiative (WHI) Observational Study (OS). The WHI OS is a major National Institutes of Health-funded research program aimed at addressing the most common causes of death, disability and poor quality of life in postmenopausal women. UB is one of 40 WHI centers nationally. CAREDS was conducted among participants at three of the centers: University of Wisconsin (Madison), the University of Iowa (Iowa City) and the Kaiser Center for Health Research (Portland, Oregon).

Researchers were able to determine participants' vitamin D status by analyzing serum samples for a vitamin D biomarker, 25-hydroxyvitamin D [25(OH)D], which provided a glimpse into vitamin D intake through all sources: diet, supplements and sunlight.

Human skin can synthesize vitamin D when exposed to ultraviolet light, Millen explains. However, for many people, 15 to 30 minutes a day with 10 percent of their skin exposed might be sufficient. In winter months, when there is a lower solar angle, sun exposure may not be sufficient to maintain blood level for people who live north of a line from about Washington, D.C., to Los Angeles. At these times and locations, dietary intake may be needed. Dietary sources of vitamin D include fortified foods such as milk and foods that naturally contain vitamin D such as fatty fish like salmon and mackerel.

"Macular degeneration has been found to be strongly associated with genetic risk," Millen said. Among many genes linked to AMD, one of the strongest is a specific genetic variant (Y402H) in the complement factor H gene, called CFH for short. This gene codes for the CFH protein that is involved in the body's immune response to destroy bacteria and viruses.

Inflammation is believed to be involved in the development of macular degeneration.

“People who have early stage AMD develop drusen, lipid and protein deposits that build up in the eye. Your body sees this drusen as a foreign substance and attacks it, in part via the complement cascade response,” explains Millen. “CFH is one of the proteins involved in this response. We see more AMD in people who have certain variants in the gene which encodes a form of this CFH protein that is associated with a more aggressive immune response.”

Vitamin D shows promise for protecting against macular degeneration because of its anti-inflammatory and antiangiogenic properties; antiangiogenic refers to slowing the growth of new blood vessels, often seen in late stages of AMD.

“Our thinking was, if a person’s vitamin D status is better, would it reduce the immune response to drusen? We wanted to understand if the association between vitamin D and AMD differed depending on a person’s genetic risk for AMD,” said Millen. “Our study suggests that being deficient for vitamin D may increase one’s risk for AMD, and that this increased risk may be most profound in those with the highest genetic risk for this specific variant in the CFH protein.”

The study results, however, shouldn’t prompt people to run to the nearest grocery store to purchase vitamin D supplements.

“Our message is not that achieving really high levels of vitamin D are good for the eye, but that having deficient vitamin D levels may be unhealthy for your eyes,” Millen said.

Although the odds of having AMD was higher in women who were deficient for vitamin D, with 25(OH)D levels below 12 ng/mL (30 nmol/L), increasing vitamin D

levels beyond 12 ng/mL did not further lower the odds of AMD to any meaningful extent, she explains.

“This study supports a role for vitamin D in eye health. That’s significant because when the Institute of Medicine’s report on the dietary reference intakes for vitamin D and calcium were released in 2011, the committee could only make conclusions about D related to bone health,” said Millen. “There wasn’t enough evidence at that time to make any recommendation based on D status and other outcomes beyond bone health.”

Source: [University of Buffalo](#)

RELATED READS



**\$2.4M
National Eye
Institute
Grant JAX**

for Retinal Pigment Epithelium Studies




**New Drug
Reduces
Rate of
Progression**

from Incurable Eye Disease



**Ophthalmolog
y's Diverse
Pipeline**

Could Rejuvenate Previously Stagnant Therapy Area



Key
Compounds
Identified to
Resolve

Abnormal Vascular Growth in AMD

Comments for this thread are now closed.



0 Comments

Drug Discovery & Development

Login

Recommend

Share

Sort by Best

This discussion has been closed.

ALSO ON DRUG DISCOVERY & DEVELOPMENT

FDA Permits Marketing of AI-based Device to Detect Certain Diabetes-

2 comments • 3 months ago

Mike_in_Durham — This comment by "Linda" is repeated virtually verbatim after several other articles about

Gene Discovery Unlocks Mysteries of Our Immunity

1 comment • 23 days ago

Monica — LOL. In July of 2017. it was discovered that I got type 2 diabetes, By the end of the July month. I was given a

A Novel 'Smart' Antibiotic May Target Most Common Bacterial Infection

1 comment • 20 days ago

GAD_tsg — While I applaud work in this area, I think its a bit early to be sending out a rather hype-high article like this.

FDA Permits Marketing of Artificial Intelligence-Based Device to Detect

1 comment • 3 months ago

Nicole — good device. My husband was diagnosed with diabetes 2 about 3 months ago. We immediately began

Subscribe Add Disqus to your siteAdd DisqusAdd

0 Dislikes 0 Dislikes 0 Dislikes 0 Dislikes