

# Dr. Scholl's® ClearAway® Wart Remover

The Drug labeling information on this website may differ from the product labeling on purchased product. Please consult your purchased package for information specific to that product.

## **Drug Facts**

Active ingredientPurposeSalicylic acid 40%Wart remover

**Uses** • for the removal of common warts. The common wart is easily recognized by the rough 'cauliflower-like' appearance of the surface

### Warnings

For external use only

#### Do not use

- if you are a diabetic
- if you have poor blood circulation
- on irritated skin or any area that is infected or reddened
- on moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes

Stop use and ask a doctor if discomfort lasts

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- wash affected area
- may soak wart in warm water for 5 minutes
- dry area thoroughly
- if necessary, cut medicated disc to fit wart
- apply medicated disc with sticky side adhering to skin
- cover medicated disc with enclosed cushion
- repeat procedure every 48 hours as needed (until wart is removed) for up to 12 weeks

# Other information

- store between 20° to 25°C (68° to 77°F)
- protect from excessive heat (temperatures above 40°C (104°F))

*Inactive ingredients* antioxidant (CAS 991-84-4), iron oxides, mineral oil, petroleum hydrocarbon resin, silicon dioxide, synthetic polyisoprene rubber, talc

Questions? 1-866-360-3226