

# ECO INSECT CONTROL SC

Reg. No. L7232 N-AR0750 Act No. 36 of 1947  
V130208

## A SUSPENSION CONCENTRATE CONTACT AND STOMACH INSECTICIDE FOR THE CONTROL OF INSECTS AS LISTED

IRAC INSECTICIDE GROUP CODE 5A

**Active Ingredient:**  
Spinosad..... 120 g/l

**Registration Holder:**  
Agro-Serve (Pty) Ltd, Reg. No. 1973/000868/07

PO Box 652147, Benmore, 2010  
15 Diesel Road, Isando, 1600

**CUSTOMER CARE:** Tel 086 133 3586 OH,  
info@efekto.co.za, www.efekto.co.za

A product from Dow AgroSciences (Pty) Ltd

**EMERGENCY NUMBER:** 082 446 8946    **UN NO:** 3082

### WARNINGS:

Withholding period - Minimum time between last application and harvest:

Chives .....	7 days	Onions .....	7 days
Cucurbits .....	7 days	Peas.....	7 days
Green beans.....	7 days	Potatoes.....	7 days
Leeks .....	7 days	Spinach .....	7 days
Lettuce .....	7 days	Tomatoes .....	3 days

- Handle with care.
- May cause skin and eye irritation.
- Moderately toxic to fish and toxic to aquatic organisms and honey bees.
- Store in a cool place.
- Store away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

### PRECAUTIONS:

- Avoid inhalation of spray mist.
- Wash with soap and water after use or accidental skin contact.
- Wear gloves when mixing and applying.
- Avoid contamination of eyes and in case of accidental contact with eyes, wash with running water for at least 15 minutes.



**CAUTION/  
VERSIGTIG**



October 2019

- Do not eat, drink or smoke whilst mixing and applying or before washing hands and face.
- Avoid spray drift onto other vegetables and fishponds.
- Clean applicators before using for other remedies and dispose of wash water where it will not contaminate vegetables, rivers and fishponds.
- Dispose of empty containers in a safe place and do not reuse for anything else.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

### DIRECTIONS FOR USE

### USE ONLY AS INDICATED

ECO INSECT CONTROL SC should be mixed with water and applied as a foliar spray on the foliage of plants with a sprayer except for lawn caterpillar control.

CROP/PEST	DOSAGE / 10 ℥ WATER	REMARKS
<b>CHIVES</b> <i>Thrips (Thrips tabaci)</i>	6 mL	Apply as a full cover spray once every 7 – 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>CUCURBITS</b> <i>Thripidae spp.</i> American (African) Bollworm ( <i>Helicoverpa armigera</i> ) American Leafminer spp. ( <i>Liriomyza trifolii</i> ) Tomato leafminer moth ( <i>Tuta absoluta</i> )	6 mL 8 mL 12 mL	Apply as a full cover spray once every 7 - 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pests. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>GREEN BEANS AND PEAS:</b> Potato leafminer ( <i>Liriomyza huidobrensis</i> )	12 mL	Apply as a full cover spray once every 7 – 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>LEAKS</b> <i>Thrips (Thrips tabaci)</i>	6 mL	Apply as a full cover spray once every 7 – 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>LAWNS</b> Lawn caterpillar	5 mL (10 ℥ spray mixture/ 20 m <sup>2</sup> )	Apply 10 ℥ spray mixture evenly over 20 m <sup>2</sup> lawn area with a fine rose watering can as a drench as soon as the lawn caterpillars appear. Repeat the application as soon as a new generation of lawn caterpillars appear.
<b>LETTUCE</b> American (African) Bollworm ( <i>Helicoverpa armigera</i> ) Tomato Leafminer moth ( <i>Tuta absoluta</i> )	8 mL 12 mL	Apply as a full cover spray once every 7 - 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>ONIONS</b> <i>Thrips (Thrips tabaci)</i>	6 mL	Apply as a full cover spray once every 7 - 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>ORNAMENTALS AND ROSES</b> Western flower thrip ( <i>Frankliniella occidentalis</i> )	15 mL	<b>DO NOT exceed the dosage.</b> <b>Corrective spray:</b> Remove all open flowers before treatment when thrips are present on Roses. Not essential on other plants but can improve efficacy. Apply as a full cover spray on foliage, open flowers and buds. <b>Preventive spray:</b> Apply as a full cover spray before buds open and apply at weekly intervals. A minimum of three applications 7 days apart should give sufficient control.
<b>POTATOES</b> Tomato leafminer moth ( <i>Tuta absoluta</i> )	12 mL	Apply as a full cover spray as soon as pest appears and repeat every seven days up to a maximum of 3 consecutive sprays in any 21-day period. Thorough wetting of plants is important.
<b>SPINACH</b> <i>Thripidae spp.</i> American (African) Bollworm ( <i>Helicoverpa armigera</i> ) American Leafminer ( <i>Liriomyza trifolii</i> ) Tomato leafminer moth ( <i>Tuta absoluta</i> )	6 mL 8 mL 12 mL 12 mL	Apply as a full cover spray once every 7 - 10 days. Use the seven-day intervals when heavy infestations occur or when favourable climatic conditions favour the pest. Start applications when the pest is first noticed and apply a minimum of two consecutive sprays to ensure effective control. Thorough wetting of plants is important.
<b>TOMATOES</b> American leafminer ( <i>Liriomyza trifolii</i> ) Tomato leafminer moth ( <i>Tuta absoluta</i> ) Africa(American) bollworm ( <i>Helicoverpa armigera</i> )	12 mL	Apply as a full cover spray as soon as pest appears and repeat every 7 days up to a maximum of 3 consecutive sprays in any 21-day period. Thorough wetting of plants is important.

# ECO INSECT CONTROL SC

Reg. Nr. L7232 N-AR0750 Wet Nr. 36 van 1947  
V130208

## 'N SUSPENSIE KONSENTRAAT KONTAK- EN MAAGINSEKDODER VIR DIE BEHEER VAN INSEKTE SOOS AANGEDUI

IRAC INSEKDODER GROEP KODE 5A

Aktiewe Bestanddeel:  
Spinosad..... 120 g/l

Registrasie Houer:  
Agro-Serve (Edms) Bpk, Reg. Nr. 1973/000868/07

Posbus 652147, Benmore, 2010  
15 Diesel Road, Isando, 1600

KLIENTEDIENIS: Tel 086 133 3586 KU,  
info@efekto.co.za, www.efekto.co.za

'n Produk van Dow Agrosciences (Edms) Bpk  
NOONDNOMMER: 082 446 8946 UN NO: 3082

### WAARSKUWINGS:

Onthoudingsperiode - Minimum tyd wat moet verstryk tussen die laaste toediening en oes:

Aartappels.....	7 dae	Pampoengewasse .....	7 dae
Blaarslai.....	7 dae	Preie .....	7 dae
Ertjes .....	7 dae	Spinasis .....	7 dae
Grasius .....	7 dae	Tamates.....	3 dae
Groenbone .....	7 dae	Uie .....	7 dae

- Hanteer versigtig.
- Kan vel en oë irriteer.
- Matig giftig vir visse en giftig vir waterorganismes en heuningbye.
- Berg in 'n koele plek.
- Berg weg van kos en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werk en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande; kwaliteit van verdunningswater; vereenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die plaag teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestatie van die betrokke middel as gevolg van die versuum van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorseen kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

### VOORSKERMMAATREELS:

- Vermy inseming van sproeinewel.
- Dra handskoene gedurende aanwending.
- Was met seep en water na gebruik van toevallige velkontak.



CAUTION/  
VERSIGTIG



Oktobter 2019

- Vermy besoedeling van oë en ingeval van toevallige kontak met oë, spoel met lopende water vir ten minste 15 minute.
- Moenie eet, drink of rook terwyl gemeng of toegeleid word, of voordat hande en gesig gewas nie.
- Vermy wegdrywing van spuitnevel na ander groentes en visdamme.
- Maak toerusting skoon voordat vir ander middels gebruik word en gooi spoelwater weg waar dit nie onder groentes, riviere en visdamme sal besoedel nie.
- Moet nie lêe houers vir enige ander doel gebruik nie en gooi die houers weg op 'n veilige plek.
- Voorkom besoedeling van kos, voedingstowwe, drinkwater en eetgerei.

### GEBRUIK SLEGS SOOS AANGEDUI

ECO INSEKBEEHEER SC word vooraf met water gemeng en as 'n loofbespuiting op plante toegediend met 'n spuit behulp van grasperkusre beheer.

PLAAG	DOSIS 10 ℥ WATER	OPMERKINGS
AARTAPPELS <i>Tomatella Blaarmyner (Tuta absoluta)</i>	12 mℓ	Dien as volle dekbespuiting toe sodra plaag opgemerk word en herhaal toediening sewed as met 'n maksimum van 3 opeenvolgende bespuitings gedurende 'n 21 dae periode. Deeglike benutting van plantie is baie belangrik.
BLAARSLAI Amerikaanse (Afrika) Bolwurm <i>(Helicoverpa armigera)</i>	8 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.
Tamatelle Blaarmyner <i>(Tuta absoluta)</i>	12 mℓ	
ERTJIES EN GROENBONE Aartappel Blaarmyner <i>(Liriomyza huidobrensis)</i>	12 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.
GRASUISE (CHIVES) Blaaspootjie <i>(Thrips tabaci)</i>	6 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.
GRASPERKE Grasperkruspe	5 mℓ (10 ℥ spuitmengsel /20 m²)	Dien 10 ℥ spuitmengsel per 20 m² grasoppervlakte as 'n grond deurdrenking toe met 'n gieter met 'n rooskop sodra die grasperkruspe opgemerk word. Herhaal die toediening sodra 'n nuwe generasie verskyn.
ORNAMENTALE PLANTE EN ROSE "Western flower thrip" <i>(Frankliniella occidentalis)</i>	15 mℓ	MOET NIÉ DIE DOSIS OORSKRIE NIÉ. KORREKTEWE BESPUITING: Verwyder alle oop blomme voor bespuiting as blaaspooties teenwoordig is op Rose. Nie nodig op ander plantte soos blomme maar dit kan effektwortel verbeter. Dien as 'olle dekbespuiting toe oor die blare, blomme en knoppe.
PAMPOENGEWASSE Blaaspootjie spp. Amerikaanse (Afrika) Bolwurm <i>(Helicoverpa armigera)</i>	6 mℓ	VOORKOMENDE BESPUITING: Dien as 'olle dekbespuiting wanneer blomknoppe open. 'n Minimum van drie bespuitings elke 7 dae behoef effektyf beheer te gee.
Amerikaanse Bladmyner <i>(Liriomyza trifolii)</i>	8 mℓ	
Tamatelle Blaarmyner <i>(Tuta absoluta)</i>	12 mℓ	
PREIE Blaaspootjies <i>(Thrips tabaci)</i>	6 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.
SPINASIE Blaaspootjie spp. Amerikaanse (Afrika) Bolwurm <i>(Helicoverpa armigera)</i>	6 mℓ	
Bladmyner <i>(Liriomyza trifolii)</i>	8 mℓ	
Tamatelle Blaarmyner <i>(Tuta absoluta)</i>	12 mℓ	
SPINASIE Blaaspootjie spp. Amerikaanse (Afrika) Bolwurm <i>(Helicoverpa armigera)</i>	12 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.
BLADMYNER <i>(Liriomyza trifolii)</i>	12 mℓ	
TAMATES Amerikaanse Blaarmyner <i>(Liriomyza trifolii)</i>	12 mℓ	
Afrika (Amerikaanse) bolwurm <i>(Helicoverpa armigera)</i>		
Tamatelle Blaarmyner <i>(Tuta absoluta)</i>		
UIE Blaaspootjies <i>(Thrips tabaci)</i>	6 mℓ	Dien as volle dekbespuiting elke 7 – 10 dae toe, met 'n minimum van twee bespuitings sodra die plaag vir die eerste maal opgemerk word. Gebruik die 7 dae interval wanneer swaar besmettings van die plaag voorkom of wanneer weerstoestande die plaag bevorderel. Deeglike benutting is van kardinale belang.