

# XTETIC Optimized Global Eyecon Protocol:

Professional Use Only



A professional treatment for the correction of multiple eye contour blemishes.

**IDEAL FOR POST INFLAMMATORY HYPERPIGMENTATION  
& MEDIUM DEPTH PIGMENT**

---

#### Periocular Peeling

##### ACTIVE INGREDIENTS:

*Azelaic Acid, Resorcinol, Phytic Acid, Salicylic, Lactic Acid, Citric Acid*

#### Periocular Solution

##### ACTIVE INGREDIENTS:

*Sweet Clover, Troxerutin, Glabra and Phytic Acid*

#### Crystal Fiber Eye Mask

##### ACTIVE INGREDIENTS:

*Tranexamic Acid, Hyaluronic Acid, Aloe Vera, Cucumber Extract*

# XTeTIC

## PROFESSIONAL

### Protocol

**Cleanse:** HydraMilk Cleanser.  
Remove with cotton pad.

**Degrease:** Gently wipe cotton pad moistened with degreasing solution around periorbital area.

**Buffer:** Protect the tear duct and outer corner of the eye with Mesolips Protector.

**Apply:** Using at most, 2ml for each eye, apply periorbital peeling solution with a cotton swab or carefully with a small fan brush uniformly on the upper lid, lower lid, and external periorbital area (expression lines).

**Time on Tissue:** Leave for 3 min. Carefully watch the client for any reactions or discomfort.

**Remove:** Wipe away excess product with a dampened cotton pad.

**Neutralize:** Spray the Post-Peel Neutralizing Spray onto a cotton pad and slide it over the treated area on both eyes. Leave for a few minutes.

### MODALITIES



ULTRA SOUND  
THERAPY



MICRONEEDLING

### Fitzpatrick

1 2 3 4 5 6

## Transepidermal Solution:

**Extract Solution:** Remove the white lid from the vial. Place the vial-syringe adapter on the vial neck exerting a slight downward pressure. Position the syringe in the adapter adjusting it with another turn of the screw. Place the vial upside down and pull the plunger backwards to empty the vial contents into the syringe. Dispense 2ml into a measuring beaker.

**\*Begin with one eye. Complete the microneedling and repeat on the other eye.**

**Microneedle:** With a small fan brush, apply to one section of the eye. Microneedle to enhance penetration of ingredients. Repeat on next section. Use maximum amount of 2ml of solution for the eye area (upper lid, lower lid, external periorbital area/expression lines). In the eye contour area, it is recommended not to exceed .25mm in depth.

**Add-On Option:** Apply HA Densimatrix, one drop per eye, to soothe and replenish treated area.

**Mask:** Apply Post-Procedure Crystal Fiber Mask. Place and adapt to the shape of the face. Leave for 15-20 minutes

**Protect:** Apply Mesoprotech Sun Protective Repairing Stick 100+ on every area treated.

### Product

- Periorbital Peeling Solution
- Crystal Fiber Mask

### Home Care

- Energy C Eye Contour
- Mesoprotech 100+ Stick
- Mesocaps Eye Contour Recovery

