



### **Optimized Protocol**

# cosmelan + melanostop tran3x

### **Professional Use Only**

A depigmenting method with an intensive corrective effect that regulates over production of melanin in the melanocytes, inhibiting and controlling new darkspots.

### **ACTIVE INGREDIENTS**

### cosmelan

Complex formula that uses 51 active ingredients, including numerous Tyrosinase inhibitors

5 main Tyrosinase inhibitors and lightening agents: Kojic Acid, Azelaic Acid, Arbutin, Phytic Acid, and Licorice.

\* MD formula is compounded with 8% Hydroquinone.

### melanostop tran3x

20% Azelaic, 10% Resorcinol, 6% phytic, 3% Tranexamic Acid





### **PROTOCOL**

### Prep- mesoestetic® hydra milk cleanser or mesoestetic® gel cleanser

- Cleanse the skin with mesoestetic® hydra milk cleanser or mesoestetic® gel cleanser using circular motions, and remove cleanser with warm water using facial sponges
- 2. Degrease the skin with mesoestetic @ oil removing solution using gauze
- 3. Protect the mucus membranes on the face (tear ducts, nostrils, and lips) using a protector balm

## Method - mesoestetic® mesopeel® melanostop tran3x & mesoestetic® cosmelan mask

- 4. Pour 2-2.5ml of **mesoestetic® mesopeel® melanostop tran3x** into measuring beaker
- 5. With a fan brush apply an even layer of **mesoestetic® mesopeel® melanostop tran3x** on entire face. Time on tissue 3-5 minutes.
- 6. Spray mesoestetic® post-peel neutralizing spray directly onto the face until it is fully absorbed (rinse with cold water to further neutralize, if needed)
- 7. With a spatula dispense 3/4 of **mesoestetic® cosmelan mask** and apply an even thick layer on entire face. Avoid the eye contour area
- 8. Determine time on tissue according to professional recommendation. (Refer to time on tissue chart)
- 9. Patient goes home with recommended masked application protocol

### Time on tissue chart

FITZPATRICK	DRY SKIN	COMBINATION/ NORMAL SKIN	OILY SKIN	MD
I & II	8 HOURS	8 HOURS	10 HOURS	4-6 HOURS
III & IV	8 HOURS	10 HOURS	12 HOURS	6-8 HOURS
V & VI	10 HOURS	12 HOURS	12 HOURS	8-10 HOURS

### **NOTES:**

Consultation plays a huge role, professional must educate patient on treatment protocol and home care protocol.

It is recommended to use at least  $\frac{1}{2}$  of the jar for the initial mask application. Save  $\frac{1}{2}$  for "spot treatment" which should be done 21 days after initial mask application.

During application time and after removing the mask, the patient must avoid exposure to sources of light and heat.

Spritz water on face to prevent mask from cracking, without removing the product.

For Asian skin types, it is recommended not to leave the mask on longer than 5 hours. These are recommended application times only.

 $Consideration \ should be \ taken \ in \ accordance \ to \ skin \ sensitivity, \ evolution \ of \ pigment, \ and \ professional \ expertise \ with \ the \ product.$ 

Different degrees of peeling vary per patient and the integrity of the skin.

Provide patient with detailed home care maintenance protocol.

Results may vary from patient to patient.

Mild to moderate erythema, stinging sensation, and tightening of the skin are to be expected after 24 to 48 hours of the cosmelan  $^{\circ}$  1 mask application.

If excess stinging sensation occurs while applying cosmelan  $^{\circ}$  2 cream, this may be adjusted.

See the cosmelan2® Home Care Protocol for more detailed home care instructions.

office 818. 432. 9500 eFax 818. 396. 3754 email info@xtetic.com

# mesoestetic® PRODUCTS

### **PROFESSIONAL**

- Cosmelan 1 Mask
- Hydra Milk Cleanser or Gel Cleanser
- Oil Removing Solution
- Mesopeel® Melanostop Tran3x
- Post-Peel Neutralizing Spray

### **HOME CARE**

- Cosmelan 2 Cream
- Melan Recovery Cream
- Melan 130+ Pigment Control Mesoprotech

### **FITZPATRICK**

All

### **SKIN CONDITIONS**

- Pigmentation
- Melasma
- Ephelides
- · Solar lentigo

### **TREATMENTS**

- Recommended: cosmelan® may only be performed twice a year. Six months after initial treatment
- As an option, you may prep skin with mesopeel® mandelic 40% as well

