

Lead-X Check Sheet

What You Should Know

- Research shows that nearly 40 percent of Americans have toxic levels of lead in their body. Unless symptoms are acute or sudden, lead poisoning can hide in the fat and bone tissue of the body.
- If you suffer from headaches, insomnia, irritability, low sex drive, nervous tremors, mood problems, nausea, depression, memory difficulties, trouble concentrating, poor coordination, or constipation you may suffer from toxic levels of lead in your system.
- The suggested safe blood level of lead is 10 micrograms/deciliter. Blood levels of lead over 2 micrograms/deciliter, however, have dramatically increased the risk of heart attack and stroke. Heart attack remains the most common cause of death in the United States.
- Researchers found that the risk of death from all causes in people with a blood level of lead over 2 micrograms/deciliter increased by 25 percent. Deaths from heart disease increased by 55 percent while risk of heart attacks increased by 151 percent and risk of stroke increased by 89 percent.
- Lead toxicity is linked to the epidemic of children with ADD and ADHD, developmental and learning problems, and autism.
- Symptoms of lead poisoning can occur below specified normal levels. Those with chemical sensitivity are more susceptible. Many people will notice an increase in joint stiffness and muscle soreness along with a decrease in energy and mental acuity.

How Lead-X Can Help

- Lead-X naturally chelates heavy metals out of the body.
- Lead-X has been proven to take significant amounts of lead out of the bloodstream.
- Lead-X is far more cost effective than the usual medical treatments for heavy metal removal.
- Lead-X is all natural and formulated by a naturopathic doctor.
- No prescription is required. Purchase now and stock up!

Lead Poisoning Symptoms

Symptoms of lead toxicity vary from patient to patient and can be difficult to diagnose properly because blood levels of lead can be misleading. As lead is fat soluble, toxic levels of lead can hide within the fat cells of the body, especially in the abdomen, joints, bones and brain.

Lead poisoning symptoms in adults include

- High blood pressure
- Abdominal pain
- Constipation
- Joint pains
- Weight loss or sudden weight gain
- Muscle pain
- Declines in cognitive ability
- Pain, numbness or tingling of the extremities
- Headaches
- Forgetfulness/memory loss
- Mood swings/disorders
- Decreased sex drive
- Reduced sperm count, abnormal sperm
- Miscarriage or premature birth

Lead poisoning symptoms in children include

- Developmental delay
- Learning difficulties
- Irritability
- Loss of appetite
- Weight loss
- Sluggishness and fatigue
- Abdominal pain
- Vomiting
- Constipation
- Hearing loss

Legal Disclaimers

This information presented is not evaluated by the FDA and is not intended to diagnose, treat, cure, or prevent any disease. Please visit your health practitioner for specific information. Herbs might alter prescription drug doses.

Unless an article or item is someone else's material, it is © 2015 by Quantum Fusion Nutraceuticals, LLC. You may make and distribute copies of this material for non-commercial purposes if you credit the source.

Do not use this product if you are pregnant, breastfeeding or taking any medication for a health condition. To avoid or minimize interferences, please take different supplements separately, if applicable. As with all other dietary supplements, please consult your physician before using this product. Please keep out of reach of children.

Actual product packaging and materials may contain more and different information than what is shown on our website. We recommend that you do not rely solely on the information presented and that you always read labels, warnings, and directions before using or consuming a product.