

NUTRI-KEY BORON SHUTTLE™

Super available liquid boron

The relationship between boron and pollen viability is so intimate that all crops should receive a foliar application of boron prior to flowering. Boron is the most leachable of all trace elements and consequently deficits are common. Boron is required as a calcium synergist and is very important in the early reproductive period of the crop cycle. Nutri-Key Boron Shuttle™ offers complexed boron in a balanced formulation that covers the full spectrum of background nutrition.

KEY PERFORMANCE

- 3.6% B plus background nutrition
- Superior Shuttle® chelation for enhanced uptake
- Improve flowering and fruit-set

GROWTH STAGE

- ✓ Vegetative Growth
- ✓ Pre-flower, Flowering, Fruit Set & Fill

SIZES:

1L, 5 L, 20 L, 200 L, 1000 L

PRODUCT CODE:

NKBS

TYPICAL ANALYSIS W/V:

Boron (B) (as sodium octaborate)	3.61%
Total Nitrogen (N) (as organic)	0.29%
Total Potassium (K)	0.51%
<i>Potassium (as organic)</i>	0.33%
<i>Potassium (as hydroxide)</i>	0.14%
<i>Potassium (as silicate)</i>	0.04%
Carbon (C) (as Shuttle® chelator)	3.38%
Sulfur (S)	0.53%
Iron (Fe)	0.31%
Zinc (Zn)	0.19%
Manganese (Mn)	0.15%
Magnesium (Mg)	938 mg/L
Copper (Cu)	490 mg/L
Molybdenum (Mo)	98 mg/L
SG	1.2
pH	6.4 – 7.4
Conductivity	20 – 40 mS/cm
Appearance	Brown liquid



NUTRI-KEY BORON SHUTTLE™

APPLICATION RATES

Vegetables

FOLIAR SPRAY RATE	0.5 L per 100 L water up to 2.5 L/ha equivalent
FREQUENCY	Twice prior to flowering or as required

Orchards & Vineyards

FOLIAR SPRAY RATE	0.5 L per 100 L water up to 5 L/ha equivalent
FREQUENCY	Twice prior to flowering or as required

Ornamentals

FOLIAR SPRAY RATE	0.5 L per 100 L water up to 3 L/ha equivalent
FREQUENCY	Twice prior to flowering or as required

Turf

FOLIAR SPRAY RATE	0.5 L per 100 L water up to 3 L/ha equivalent
FREQUENCY	Every 4 weeks or as required

Spot-Spraying

FOLIAR SPRAY RATE	25 mL per 10 L water
FREQUENCY	As required

Home Garden Application

Dilute 7.5 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre. Do not apply more than 6 times per year without demonstrated trace element requirement.

*Disclaimer: Any recommendations provided by Nutri-Tech Solutions Pty Ltd (NTS) or its Distributors are advice only. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (all of which may affect the performance of our program), no responsibility for, or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise), arising from such storage, mixing, application, or use will be accepted under any circumstances whatsoever. NTS recommend you contact an Agronomist prior to product application. The Buyer assumes all responsibility for the use of any NTS products.

Nutri-Tech Solutions P/L (NTS)

7 Harvest Road, Yandina, Queensland 4561, Australia

P +61 7 5472 9900 | E info@nutri-tech.com.au

www.nutri-tech.com.au

INSTRUCTIONS

Stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application.

Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application.

Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

IMPORTANT

This product contains 3% boron. Boron can be toxic to plants at high levels. Consult your NTS Agronomist for application advice.

Avoid spraying close to harvest if produce is susceptible to staining.

Note: Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

STORAGE & HANDLING

Read safety directions before use.

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 5°C. Crystallisation or sedimentation may occur below 5°C. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

YOUR LOCAL NTS DISTRIBUTOR: