

# DOMINATE-F™

**Microbe brew adjuvant for stabilisation & fungal domination**

**Specialised food formulation for microbial brews**

LMF™ contains the essential ingredients proven to promote crucial microbial diversity during the multiplication process. It ensures rapid microbe multiplication and reduces contamination potential.

## KEY PERFORMANCE

- Reduce contamination issues
- Optimise fungal-brew numbers
- Improves brew stability
- Fungal brews are ideal for bio-balancing

## SIZES:

1L, 5 L, 20 L

## PRODUCT CODE:

DOMFUN

**ACO Certification No. 456**

## TYPICAL ANALYSIS W/V:

SG	1.4
pH	4.4 – 5.4

\* This product is allowed for use in organic production systems.



# DOMINATE-F™

## BREWING INSTRUCTIONS

All brewing equipment must be thoroughly cleaned before and after use (e.g. using Path-X™ at 100 mL/10 L).

When using potentially contaminated water sources (e.g. dam water) treat water first using bleach (4% available chlorine) at a rate of 50 mL per 100 L of water.

Proportional amounts of bleach will be required if the available chlorine % varies. Aerate for at least one (1) hour, until the smell of chlorine dissipates.

## IMPORTANT

This is a brewing adjuvant, not a microbial inoculum.

## APPLICATION RATES

### Hydroponics

---

**RATE** Use 1 L Dominate-F™ together with 1 L LMF™ (Liquid Microbe Food) per 100 L brew

---

Refer to inoculums for specific brewing instructions.

**Note:** Dominate-F™ should be used in conjunction with LMF™.

---

## INSTRUCTIONS

### Shake or stir well before use.

Ensure compliance with your quality assurance code of practice regarding the use of microbial products before use.

**Note:** Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

## STORAGE & HANDLING

Read safety directions before use.

Store in ambient conditions out of direct sunlight.

This is not a microbial product but is used in the production of microbial brews.

While handling, brewing and applying microbial products protective clothing, gloves and face masks should be worn.

Avoid aerosols generated by aeration.

\*Disclaimer: Rates are dependent on biomass and water rates and should be used as a guide only. Application rates and frequencies should be adjusted according to crop monitoring and demand. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare as listed on label. Sensitive plants, or foliage/fruit easily susceptible to staining, should be test sprayed prior to full scale application. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use. Refer to product label or PIS for alternate application rates.

### Nutri-Tech Solutions P/L (NTS)

7 Harvest Road, Yandina, Queensland 4561, Australia

P +61 7 5472 9900 | E info@nutri-tech.com.au

[www.nutri-tech.com.au](http://www.nutri-tech.com.au)

---

## YOUR LOCAL NTS DISTRIBUTOR:

---