

NUTRI-CARB-N™

15% N with humic acid.

This convenient combination of liquid nitrogen complexed with humic acid effectively addresses high nitrogen requirements whilst conditioning the soil, promoting healthy root growth and stimulating beneficial soil microbes.

KEY PERFORMANCE

- Plant available N in amine form for all crop stages
- Complexed with humic acid to prevent leaching & condition the soil
- Stimulates beneficial soil microbes

GROWTH STAGE

- ✓ Vegetative Growth
- ✓ Pre-Flowering, Flowering, Fruit Set & Fill

SIZES:

1 L, 5 L, 20 L, 200 L, 1000 L

PRODUCT CODE:

NCN

TYPICAL ANALYSIS W/V

Total Nitrogen (N) (as urea)	15.18%
Total Potassium (K) (as organic)	1.10%
SG	1.1
pH	9.0 – 11.5
Conductivity	10 – 15 mS/cm
Appearance	Black Liquid



NUTRI-CARB-N™

APPLICATION RATES

Vegetables

FOLIAR SPRAY RATE

1L per 100 L water up to 5 L/ha equivalent

FERTIGATION RATE 10 – 40 L/ha

FREQUENCY Weekly or as required

Orchards & Vineyards

FOLIAR SPRAY RATE

1L per 100 L water up to 7 L/ha equivalent

FERTIGATION RATE 10 – 40 L/ha

FREQUENCY Every 2 – 4 weeks or as required

Ornamentals

FOLIAR SPRAY RATE

1L per 100 L water up to 5 L/ha equivalent

FERTIGATION RATE 10 – 30 L/ha

FREQUENCY Every 2 – 4 weeks or as required

Turf

FOLIAR SPRAY RATE

1L per 100 L water up to 5 L/ha equivalent

FERTIGATION RATE 10 – 40 L/ha

FREQUENCY Every 2 – 4 weeks or as required

Broadacre Crops

FOLIAR SPRAY RATE

3 – 5 L/ha in 80 – 120 L water, depending on canopy closure

FREQUENCY As required

Pasture

FOLIAR SPRAY RATE

3 – 5 L/ha in 80 – 120 L water

FREQUENCY As required

Spot-Spraying

FOLIAR SPRAY RATE

50 mL per 10 L water

FREQUENCY As required

INSTRUCTIONS

Shake or stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application.

Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, windspeed), prior to foliar application.

Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

IMPORTANT

Contains maximum 0.15% biuret. Not compatible with acidic inputs. Avoid spraying close to harvest if produce is susceptible to staining.

Note: Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

STORAGE & HANDLING

Read safety directions before use. Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 5°C. Crystallisation or sedimentation may occur below 5°C. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

YOUR LOCAL NTS DISTRIBUTOR:

*Disclaimer: Rates are dependent on biomass and water rates and should be used as a guide only. Application rates and frequencies should be adjusted according to crop monitoring and demand. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare as listed on label. Sensitive plants, or foliage/fruit easily susceptible to staining, should be test sprayed prior to full scale application. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application. Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use. Refer to product label or PIS for alternate application rates.

Nutri-Tech Solutions P/L (NTS)

7 Harvest Road, Yandina, Queensland 4561, Australia

P +61 7 5472 9900 | E info@nutri-tech.com.au

www.nutri-tech.com.au